



Heat transfer during high wind episodes in the marine atmospheric surface layer

Sørensen, L.L.; Sempreviva, Anna Maria; Pryor, Sara

Published in:
Abstracts (CD-ROM)

Publication date:
2008

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Sørensen, L. L., Sempreviva, A. M., & Pryor, S. (2008). Heat transfer during high wind episodes in the marine atmospheric surface layer. In *Abstracts (CD-ROM) EMS. EMS Annual Meeting Abstracts 2008 No. 5*

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Heat transfer during high wind episodes in the marine atmospheric surface layer

L.L.Sørensen (1), A.M.Semprevia (2,3), S.C.Pryor (3,4)

(1) National Environmental Research Institute, University of Aarhus, Denmark (2) Institute of Atmospheric Science and Climate - CNR Italy, (3) Department of Wind Energy, Risø-DTU, Denmark. (4) Indiana University, Bloomington, USA, (Contact Email: lls@dmu.dk)

The exact estimate of heat fluxes is of fundamental importance for applications of models at different scales. In the last few years, many scientists have addressed the limitation of the fluxes parameterisation using the Monin Obhukov (M-O) theory in the surface layer especially in the marine environment. Casual factors include non-local effects, i.e. entrainment from large eddies Johanson (2003) or interaction with sea spray particles (Andreas and De Cosmos, 2002). In the latter paper, reanalysis of the HEXOS (Humidity Exchange Over the Sea) data in open sea, indicate sea spray may contribute up to 40 % of the total latent flux for wind speed between 15 ms⁻¹ and 18 ms⁻¹. We first addressed this problem considering the basic issue of the correlation coefficient between humidity q' and temperature T' fluctuations defined as $R_{qT} = \langle q'T' \rangle / (s_q s_T)$, where $\langle q'T' \rangle$ is the q, T mean covariance, and s_q and s_T the standard deviations of q and T respectively. Both q and T are scalars, which are transported by turbulence, therefore it should be $R_{qT} = 1$. Sempreviva and Gryning (1996, 2000) and Sempreviva and Højstrup (1998), found that R_{qT} seldom reaches such a value in the surface layer. Here, we examine correlation coefficients for the scalars q , T and CO_2 in the marine surface layer using half hour average turbulent fluxes of moisture ($\langle w'q' \rangle$), heat ($\langle w'T' \rangle$), CO_2 ($\langle w'CO_2' \rangle$) and marine aerosols concentrations collected during an intensive campaign performed under the EU MEAD (Marine Effects of Atmospheric Deposition) project at a Swedish coastal site and during the Danish Galtæa III Cruise, where different sonic anemometers and fast sensors to measure humidity, CO_2 and particles have been used. We have chosen periods with high wind speed, estimated spectra and co-variances from different sensors and

considered the effect of sea spray and boundary layer height.