



Live-Fish-Facs-Droplet Cultivation: A Strategy to Explore Recalcitrant Acidobacteria

Nita, Iuliana; Buijs, Yannick; Bentzon-Tilia, Mikkel

Publication date:
2023

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Nita, I., Buijs, Y., & Bentzon-Tilia, M. (2023). *Live-Fish-Facs-Droplet Cultivation: A Strategy to Explore Recalcitrant Acidobacteria*. 250-250. Abstract from 16th Symposium on Bacterial Genetics and Ecology, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

[P159] LIVE-FISH-FACS-DROPLET CULTIVATION: A STRATEGY TO EXPLORE RECALCITRANT ACIDOBACTERIA

Juliana Nita¹, Yannick Buijs, Mikkel Bentzon-Tilia

¹*Technical University of Denmark, Department of Biotechnology and Biomedicine, Kgs. Lyngby, Denmark*

Antibiotics from few culturable microorganisms have prevented millions of deaths over the past many decades. While initially effective, their efficacy has dramatically decreased due to the emergence and spread of antibiotic resistance. As antibiotic resistant microorganisms are predicted to cause 10 million deaths by 2050, we are in dire need of novel antimicrobial molecules. The potential of natural microbiomes to produce diverse secondary metabolites (SMs) with therapeutic properties is an attractive solution to mitigate this crisis, and the abundant acidobacterial phylum has proven to harbor a vast repertoire of biosynthetic gene clusters encoding potentially bioactive SMs. However, most clades of the Acidobacteria are recalcitrant to cultivation, which impedes our ability to harness the potential of these SMs. To circumvent this limitation, we present a strategy aiming at coupling Fluorescence in situ Hybridization within living bacteria (live-FISH) with Fluorescence-Activated Cell Sorting (FACS) and droplet cultivation to selectively isolate these bacteria in order to gain access to previously uncultured and biosynthetically talented acidobacterial strains. Employing DNA probes hybridizing to the 16S rRNA of Acidobacteria subdivision 1 in a conventional FISH experiment allowed us to identify members of this group, including the uncultivable *Angelobacter* clade, from soil samples. Following this, we utilized *Acidobacterium capsulatum* as a model organism to demonstrate the feasibility of a live-FISH methodology targeting Acidobacteria. Despite a severe reduction in cell viability, we were able to improve cell survival through a modified hybridization approach. Separate from the Live-FISH procedure, we encapsulated *A. capsulatum* cells in minuscule droplets, and after three days of incubation, we observed proliferation, demonstrating the capacity of *A. capsulatum* to thrive in 200 pL-droplets. Further refinement of the live-FISH method and its integration with droplet encapsulation and cultivation will lead to targeted isolation of previously uncultured acidobacterial strains, allowing us to tap into the biosynthetic repertoire of this promising phylum.