



Environmental sustainability of food production and consumption in the nordic and baltic region - a scoping review for nordic nutrition recommendations 2023

Harwatt, Helen; Benton, Tim G.; Bengtsson, Jan; Birgisdóttir, Bryndis Eva; Brown, Kerry Ann; Van Dooren, Corné; Erkkola, Maijaliisa; Graversgaard, Morten; Halldorsson, Thorhallur; Hauschild, Michael

Total number of authors:
17

Published in:
Food and Nutrition Research

Link to article, DOI:
[10.29219/fnr.v68.10539](https://doi.org/10.29219/fnr.v68.10539)

Publication date:
2024

Document Version
Early version, also known as pre-print

[Link back to DTU Orbit](#)

Citation (APA):

Harwatt, H., Benton, T. G., Bengtsson, J., Birgisdóttir, B. E., Brown, K. A., Van Dooren, C., Erkkola, M., Graversgaard, M., Halldorsson, T., Hauschild, M., Høyer-Lund, A., Van Oort, B., Saarinen, M., Tuomisto, H. L., Trolle, E., Ögmundarson, O., & Blomhoff, R. (in press). Environmental sustainability of food production and consumption in the nordic and baltic region - a scoping review for nordic nutrition recommendations 2023. *Food and Nutrition Research*, 68. <https://doi.org/10.29219/fnr.v68.10539>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Review article

Received: 15 November 2022; Revised: 12 June 2023; Accepted: 24 January 2024

Title: Environmental sustainability of food production and consumption in the Nordic and Baltic region - a scoping review for Nordic Nutrition Recommendations 2023

Authors: Helen Harwatt¹, Tim G. Benton¹, Jan Bengtsson², Bryndís Eva Birgisdóttir³, Kerry Ann Brown⁴, Corné van Dooren⁵, Maijaliisa Erkkola⁶, Morten Graversgaard⁷, Thorhallur Halldorsson⁸, Michael Hauschild⁹, Anne Høyer-Lund¹⁰, Jelena Meinilä⁶, Bob van Oort¹¹, Merja Saarinen¹², Hanna L. Tuomisto^{6,12}, Ellen Trolle⁹, Olafur Ögmundarson⁸, Rune Blomhoff¹³

Affiliations: ¹ Royal Institute of International Affairs, Chatham House, London, UK ² Department of Energy and Technology, Swedish University of Agricultural Sciences, SLU, Uppsala, Sweden, ³ University of Iceland, Reykjavik, Iceland ⁴ University of Exeter, Devon, UK ⁵ World Wildlife Fund, Zeist, Netherlands, ⁶ University of Helsinki, Helsinki, Finland ⁷ Aarhus University, Aarhus, Denmark ⁸ University of Iceland, Reykjavik, Iceland ⁹ National Food Institute, Technical University of Denmark (DTU), Kgs. Lyngby, Denmark, ¹⁰ The Norwegian Directorate of Health, Oslo, Norway, ¹¹ Centre for International Climate Research, Oslo, Norway, ¹² Natural Resources Institute, Helsinki, Finland, ¹³ Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway

Corresponding author:

Helen Harwatt
Royal Institute of International Affairs
Chatham House
London
UK
e-mail: harwatt@chathamhouse.org

Abstract

This scoping review examines environmental impacts related to food production and consumption in Nordic and Baltic countries. The overarching advice to all Nordic and Baltic countries, in line with the current body of scientific literature, is to shift to a more plant based dietary pattern and avoid food waste. Taking into account current consumption patterns, there is a high potential and necessity to shift food consumption across the countries to minimize its environmental impact. More specifically, a substantial reduction in meat and dairy consumption and increased consumption of legumes/pulses, whole grains, vegetables, fruits, nuts and seeds is suggested as a priority intervention. Reducing the environmental impacts of seafoods is also key and suggestions include a shift to seafoods with lower environmental impacts such as seaweed and bivalves. As part of the suggested transition to a more plant-based diet, the scope for increasing the provision of plant-based foods through increasing the cultivation of legumes/pulses, vegetables and grains and through feed-to-food shifts within the region should be explored.

Popular summary

- The environmental impacts related to food production and consumption have been assessed
- The paper take a predominantly global perspective, while discussing the context and implications across the eight focal Nordic and Baltic countries
- The overarching advise to all eight countries is to shift to a more plant-based dietary pattern

Introduction

The global food system currently exerts substantial environmental burdens, from harvesting fish and seafood from nearly every river, lake and ocean; using around 50% of habitable land and 70% of available freshwater; causing major losses of nitrogen and phosphorous; being the main source of global anthropogenic methane emissions and accounting overall for around a third of anthropogenic greenhouse gas emissions – driving biodiversity loss, deforestation, climate change and eutrophication of fresh water and coastal ecosystems.¹ On its current trajectory, food production is likely to cause more losses of biodiversity and carbon sinks such as forests, and further global temperature rise.² Such impacts are already, and will increasingly, threaten human health and planetary health. For example, widespread and rapid changes in the atmosphere, ocean, cryosphere and biosphere have occurred; and climate change is affecting many weather and climate extremes in every region across the globe – causing adverse impacts to ecosystems and human populations.³


The planetary boundaries framework brings these major issues together to assess the current situation in relation to a set of interrelated and interacting biogeochemical boundaries at the global level – limits that if exceeded could result in large-scale abrupt (and potentially irreversible) ecosystem and climate destabilisation.⁴ The limits are based on evidence from earth system science in relation to how much of a given action or substance can be tolerated at the global level until regime shifts are likely triggered. Some boundary limits (e.g. climate) are based on evidence of large-scale (i.e., global, continental or regional) threshold behaviours, whereas other boundary limits (e.g., biodiversity and freshwater) are based on their interaction with sub-global or regional processes and boundaries. Assessments suggests that 6 of 9 planetary boundaries have exceeded a safe limit (climate change, land use change, biodiversity loss, biogeochemical flows (nitrogen and phosphorus), freshwater and novel entities (including plastics)).⁵ A more recent analysis of spatial variability in both ecosystems' sensitivity to nitrogen pollution and agricultural nitrogen losses found that the aggregated global surplus boundary for nitrogen is far exceeded by the current nitrogen surplus.⁶ Although not without its limitations,⁷ or criticisms – including those relating to the difficulty in defining global ecosystem thresholds for local environmental impacts⁸, the planetary boundaries concept uniquely and visually provides an understanding and broad framing regarding key environmental limits within which food systems should operate, beyond the limited scope of climate change. We use the planetary boundaries framework in this paper to explore a range of environmental impacts in a comparative way across the Nordic and Baltic countries rather than as a predictive or prescriptive tool, and provide background data and methodologies to enable comparisons with other measures of sustainability (see box 1 and appendices 1-3). The difference between countries in terms of impacts should not be interpreted as a lack of need to reduce impacts in countries with relatively small impacts, or to focus attention solely on countries with relatively large impacts.

No region in the world is on course to meet the food related portion of global environmental targets, indicating that the global food system is exceeding a safe trajectory to stay within the planetary boundaries (as indicated in Figure 1, where all 2018 impacts are above 100% in each region. In this test, regional diets in 2010 and 2018 are globally adopted and compared to global environmental targets).⁹ Despite a pressing need to reduce environmental burdens from all sectors, global impacts from the food system have increased by up to 14% over the past decade alone and have contributed to the exceedance of five global environmental thresholds.¹⁰ For example, in 2018, food-related greenhouse gas emissions at the global level exceeded the limit consistent with keeping warming below 2°C by 74% (Figure 1). Cropland use was 60% above the value aligning with limiting the loss of natural habitat according to Aichi Biodiversity Targets (SDG 15). Freshwater use exceeded

sustainable withdrawals by over 52% (SDG 6.4). Nitrogen and phosphorus application exceeded values (by 113% and 67% respectively) that would limit marine pollution to acceptable levels (as defined in SDG 14.1). Hence, major changes are required across all parts of the food system. This includes addressing the environmental impacts of international trade, which causes impacts in supplier regions that are not accounted for by importing regions e.g., trade of agricultural commodities between Latin America and Europe. Figure 2 shows the relative proportion of global environmental impacts from each food type, with beef accounting for the majority of land use and the largest proportion of GHGs, and grains accounting for the largest proportion of freshwater, nitrogen and phosphorus use.


It is important to include the environmental impacts in exporting countries of imports, and not limit sustainability considerations to local production as this can skew 'rankings' or perceptions of sustainability. For example, the Sustainable Development Report shows a very different performance score when 'spillover' impacts are included in an overall score that measures the total progress towards achieving all 17 SDGs and the impact of each country's actions on other countries' abilities to achieve the SDGs. When factoring in spillovers including 'environmental and social impacts embodied into trade', Finland, Denmark, Sweden, Norway shift from their top 4 ranking positions globally on progress toward meeting SDGs, to 124, 137, 139, and 146 global rankings respectively.¹¹ The impacts of imports could differ substantially depending also on agricultural production standards in exporting countries, such as nitrogen use and types of permitted plant protection products. Politically however, there could be a preference to import goods as their environmental impacts are not included in national inventories of importing countries (for example, in the reporting of national GHGs under the UNFCCC). Such accounting preferences are not beneficial in terms of reducing overall environmental burdens from the food system.

Figure 1: Global sustainability test comparing impacts of the food system with five global environmental targets¹²



Source: *Global Nutrition Report 2021*¹³ The analysis utilizes country-specific food consumption and environmental footprint data and relates them to the food portions of global environmental targets i.e. a global test to assess the impacts if everyone in the world consumed at the given rate. The methods and data are described in Box 1, and presented in appendices 1-2.

Figure 2: Environmental impacts of the global food system by type of impact and food group as a proportion of total impact



Source: *Global Nutrition Report 2021*¹⁴ The analysis utilizes country-specific food consumption and environmental footprint data. The methods and data are presented and described in Box 1 and appendix 2.

Tackling the planetary health crisis requires extraordinary levels of action across all sectors, at an unprecedented speed. According to the latest assessment from the Intergovernmental Panel on Climate Change (IPCC), “Rapid and far-reaching transitions across all sectors and systems are necessary to achieve deep and sustained emissions reductions and secure a liveable and sustainable future for all.”¹⁵ Taking transformative action rapidly this decade, including in most cases immediate emissions reductions in all sectors, would not only reduce projected adverse impacts on humans and ecosystems but would deliver many co-benefits, especially for human health. Taking such actions in the near-term typically requires a range of enabling policies to provide high up-front investments and lessen potentially disruptive changes.¹⁶ Transforming food systems is a critical part of meeting planetary health goals. A substantial departure from business-as-usual expectations is required, and a shift in the roles and responsibilities of public sector actors versus businesses in shaping dietary demand.¹⁷ This is even more apparent if taking into account safe *and* just earth system boundaries (ESBs) that minimize humans’ exposure to significant harm from Earth system change (by comparison, the PBs identify only safe biophysical boundaries). A recent analysis at global and sub-global scales found that seven of eight globally quantified safe and just ESBs and at least two regional safe and just ESBs in over half of the global land area are already exceeded. Ensuring human well-being thus requires systemic transformations across energy, food, urban and other sectors. Addressing the drivers of Earth system change is also required (including the economic, technological, and political), together with an increased understanding of the role of justice, economics, technology and global cooperation in creating a safe and just future.¹⁸

This paper is the result of an expert elicitation developed as a collaboration between the NNR2023 project, Chatham House and Nordic and Baltic scientists. Helen Harwatt and Tim Benton, Chatham House, are lead authors. The contributing experts (JB, RB, BEB, KAB, CvD, ME, MG, TH, MH, AH, JM, BvO, MS, ET and OÖ) have given significant scientific input relevant to the local context, while the members of the NNR Committee (RB, ET, TH, ME, and AH) have ascertained that the content is relevant to and within the scope of the NNR project. This paper, in addition to several forthcoming papers and other major reports, will assist the NNR Committee when formulating science advice to the authorities.

Aims of the paper

This paper examines environmental impacts related to current food production and consumption from the perspective of identifying problems and solutions. Because demand and supply are inextricably connected, for useful context we provide an overview of food production (including imports and exports), and consumption in relation to each major food group. We take a predominantly global perspective, while discussing the context and implications across the 8 focal Nordic and Baltic countries (Denmark, Finland, Iceland, Norway, Sweden, Estonia, Latvia, and Lithuania). The global perspective is mostly relevant to high income country settings, such as those across the Nordic and Baltic region. An overview of environmental sustainability considerations is provided for each food group within the Nordic Nutrition Recommendations (NNR), from these two perspectives (global and regional). From this assessment, we provide guidance for reducing the environmental impacts of food consumption. We identify considerations for reducing environmental

impacts in each food group, however the overarching consideration is to reduce the absolute environmental burden at the food system level, and where possible to reduce the absolute environmental impacts of food groups while recognizing that some need to increase (e.g., fruits and vegetables) to cover nutritional needs and improve population health. The provision of prescriptive and detailed advice for each individual country is beyond the scope of this paper, as are recommendations on achieving sustainable food production. Similarly, we are aware that some of the focal countries place an importance on self-sufficiency in terms of food production, and a number of analyses have explored its prospects.^{19,20,21} While it might be possible to meet both environmental and self-sufficiency targets in some contexts and countries, self-sufficiency is not synonymous with sustainable food consumption. Also, for some countries (e.g., Iceland), growing enough food to feed the entire country is not a realistic option, even if it was desirable from an environmental sustainability perspective. The complexity of reconciling self-sufficiency goals with national diets and environmental sustainability from the perspective of planetary boundaries is beyond the scope of this paper.

Research approach

Both between and within the countries, it is difficult to fully consider the great variation in agricultural structure and production, topographic and climate conditions, and land use.²² Given this consideration, combined with the unavailability of a comprehensive dataset on the spectrum of environmental impacts of food types consumed in the Nordic and Baltic region (e.g., biodiversity impacts across all food groups), we approached this assessment partly as an expert elicitation to ensure that the rich body of existing data on environmental impacts of foods and diets could be best interpreted within the context of the Nordic and Baltic region, and hence, within the project constraints, the contents would be as relevant as possible to the NNR2023. We collected input from 20 regional experts through a series of three workshops and via multiple reviews of the manuscript. The workshops were used to discuss the paper and its content in detail, and environmental sustainability issues within the region of relevance to the NNR food groups, including those global in scope, with focussed breakout sessions on meat, dairy, fruits and vegetables, pulses, fish and seafood, and ultra processed foods. The experts also contributed a range of specialist knowledge in relation to environmental impacts of food production and consumption across the region during three subsequent reviews of the paper. A workshop with all co-authors was also convened before the paper went to the public hearing. To maximize the confidence of our recommendations, in addition to the expert elicitation method we draw from existing analyses and reviews from the scientific literature and refer to research conducted on the Nordic and Baltic region.

The 20 regional experts (see acknowledgements) were appointed by the NNR2023 Committee based on a public call and after careful evaluation of each expert's competence and experience. To supplement the call, experts were also recruited after invitation from the NNR2023 Committee and from Chatham House. A fair distribution of experts among the Nordic and Baltic countries, in addition to experts with global competence, were sought when appointing experts. In addition, experts were recruited to bring a wide range of specialist knowledge from different research areas. While the input of the regional experts is highly appreciated, the final text in this paper is the sole responsibility of the authors.

Background information regarding food production and consumption in the 8 focal countries is provided in section 1, in addition to the aspects of environmental sustainability that apply across food groups and can therefore be considered as more general in application. Section 2 provides an overview of environmental sustainability considerations from a global and regional context for each NNR food group that we consider to be most significant in terms of consumption and environmental

sustainability. Section 3 provides an overview of environmental sustainability considerations from a global and regional context for each NNR food group that has less overall environmental impact as a proportion of total diets compared to the focal food groups in section 2, but which are still considered. Section 4 provides key considerations for the environmental sustainability of food consumption in Nordic and Baltic countries. Given the urgency of tackling a range of environmental issues, including climate change, we focus on options that have potential to reduce environmental impacts substantially in the immediate to short term, without reliance on supply side technologies currently under development, or not yet in widespread use. Hence, we focus our guidance on demand side shifts in consumption.

Data sources for food availability, food consumption and environmental impacts of food

The environmental analyses of food used in this paper represent the best available sources to demonstrate the impacts of food consumption in consistent units of analysis at the aggregate level across a range of food groups and across the Nordic and Baltic countries. As with all models that are analysing aggregate level data and impacts, the estimates provide an indication of impacts and a relative comparison between countries, foods and food groups. They do not calculate the impacts of food production at the granular, detailed level within countries i.e., specific amounts of the numerous foods produced using a number of specific agricultural techniques. Hence, the environmental impact data presented in this paper do not constitute a national food system assessment that might take consider the detailed impacts of food production and consumption.

All models for country-specific environmental footprint of foods, food groups and food systems have specific data sources and handling methods. In Box 1 we describe data sources and handling that constitute the basis for the main environmental analysis and impact data used and cited in this paper.

Box 1: Environmental data sources and handling methods

- Data on food production, imports, exports and food supply quantity (kg per person per year) for all 8 countries are provided in most food group chapters, in tables 2.2, 2.3, 3.2, 3.3, 3.4, 3.5, 4.2, 4.3, 4.4, 4.5, 5.2, 5.3, 6.2, 7.1, 7.2, 8.2, 8.3, 8.4, 9.2, 10.2, 11.2, and 11.3. The data were sourced from FAOSTAT Food Balance Sheets (FBS) for the year 2019. The FAO provide the following description: “FBS are compiled every year by FAO, mainly with country-level data on the production and trade of food commodities. Using these data and the available information on seed rates, waste coefficients, stock changes and types of utilization (feed, food, processing and other utilization), a supply/utilization account is prepared for each commodity in weight terms. The food component of the commodity account, which is usually derived as a balancing item, refers to the total amount of the commodity available for human consumption during the year. Besides commodity-by-commodity information, the FAO FBS also provide total food availability estimates by aggregating the food component of all commodities including fishery products. From these values and the available population estimates, the per person dietary energy and protein and fat supplies are derived and expressed daily. In the FBS production data refer only to primary products while data for all other elements also include processed products derived there from,

expressed in primary commodity equivalent.”²³ Food production is reported at the farm level for crop and livestock products (i.e. in the case of crops, excluding harvesting losses) and in terms of live weight for fish items (i.e. the actual ex-water weight at the time of the catch). Data are expressed in terms of dressed carcass weight, excluding offal and slaughter fats. Production of beef and buffalo meat includes veal; mutton and goat meat includes meat from lambs; pig meat includes bacon and ham in fresh equivalent. Poultry meat includes meat from all domestic birds and refers, wherever possible, to ready-to-cook weight. Per capita supply figures represent only the average supply available for the population as a whole and do not necessarily indicate what is actually consumed by individuals. Even if they are taken as approximation to per capita consumption, it is important to note that the amount of food actually consumed may be lower than the quantity shown in the FBS supply figures, depending on the degree of losses of edible food and nutrients in the household, e.g. during storage, in preparation and cooking.²⁴ For this reason, where we use FBS food supply data in this paper, we refer to it as food supply, rather than food consumption.

- Every year, authorities in over 245 countries and territories submit national food and agriculture statistics, as well as micro datasets collected through farm and household surveys, to FAO. In the FAOSTAT database, national food supply is estimated as average per capita foods available for consumption based on domestic production adjusted for exports, imports, and non-food uses. National authorities, who annually supply data to FAOSTAT, are not required to use standardised methodologies when collecting, categorizing and grouping data, which is a limitation of using FAOSTAT data in country-specific studies.
- To analyse the environmental impact of food availability, country-specific data from FAOSTAT are often paired with comprehensive country-specific databases of environmental footprints or life cycle assessment data that, to various degrees, include primary production, farmed animal feed requirements, processing, transport, and packaging, and may also consider food loss and waste. These models build on numerous assumptions and theoretical considerations. Despite commonly held limitations with all types of complex modelling endeavours, the models of environmental footprints that we refer to in this paper are innovative and represent the state-of-the-art at a level suitable for providing an indicative assessment of the environmental impacts of food consumption across countries and cross-country comparisons.
- In epidemiological national dietary surveys 24-hour recalls, food records, and Food Frequency Questionnaires are commonly used as evaluated methods of dietary assessment. In this paper, the country-specific food consumption, based on such methodologies, is compared to national dietary guidelines in Table 2. However, due to the differences in methodologies, food consumption data in table 2 sourced from dietary surveys is not necessarily directly comparable with food supply data used in this paper, sourced from the FAOSTAT FBS. There are advantages and disadvantages with both datasets.
- In this paper, we use datasets from 4 analyses to demonstrate the environmental impacts of foods and food consumption at global, regional and national levels – and in some cases, how they relate to the food portion of planetary boundaries: i) Springmann et al (Nature, 2018), ii) Springmann et al. (BMJ, 2020), iii) Global Nutrition Report 2021 and iv) Poore and Nemecek (Science, 2018).²⁵ Food supply corrected for waste (from FAO food balance sheets) is used as an estimation of food consumption. When interpreting these data, it is important to understand that food consumption estimated from food supply in food balance sheets may vary from the food consumption assessed by food records or 24-hour diet recall methodologies. In relation to i) and ii) above: Springmann et al (Nature, 2018) and Springmann et al. (BMJ, 2020) used the same environmental footprints and data sources.

Country-specific environmental footprints were calculated by combining national food consumption data with data on country-specific environmental footprints for crop, livestock and fish production. Specifically, food consumption was calculated from the FAOSTAT FBS database as country-specific food supply data adjusted for waste (see appendices 1-3). The environmental footprint data were calculated as follows:

- GHG emissions:
 - Data on crop related GHG emissions were taken from Carlson et al. (Nature Climate Change 2017). Carlson et al. compiled crop harvest and management data at the national and subnational level from > 15,000 units in the world, ranging from countries, states and countries, as well as agricultural data from FAOSTAT (Monfreda et al. Glob. Biogeochem. Cycles 22, GB1022 (2008) and Ramankutty et al. Glob. Biogeochem. Cycles 22, GB1003 (2008)). These data provide a crop-specific national and subnational assessment of how agricultural management practices interact with biophysical characteristics to generate heterogeneous patterns of resources used for crop production. Country-specific GHG emissions from crops were then calculated from these data by using standard IPCC Tier 1 methodology for GHG emissions.
 - Data on livestock related GHG emissions (including feed related emissions) were taken from Tubiello et al (Environ. Res. Lett 2013). They compiled a global emissions database with country level details based on an inventory-based, bottom-up accounting of activity data from FAOSTAT (2012) and GHG emissions using Tier 1 IPCC methodology.
 - GHG emissions from farmed and wild-caught fish were calculated from data on fish production from Troell et al (PNAS 2014), Chen et al (WorldFish Center and Int. Food Policy Res. Inst. 2017) and Troell et al (WorldFish Center and Int. Food Policy Res. Inst., 2014) and GHG emissions using IPCC methodology (Rosegrant et al, Int. Food Policy Res. Inst, 2017).
- Country specific data on cropland and blue water use were adopted from the IMPACT model (Robinson et al IFPRI 2015). These data include feed requirements for farmed animals in terrestrial and aquatic systems.
- iii): The Global Nutrition Report 2021 uses estimates of food demand calculated from FAO Food Balance Sheets and a database of country and food group-specific environmental footprints from i) Springmann et al. Nature 2018 and iv) Poore & Nemecek. Science 2018 (see appendices 1-3). Data on food demand for each country from the FAO was paired with a comprehensive database of environmental footprints, differentiated by country, food group, and environmental impact. The footprints take into account all food production, including inputs such as fertilisers and feed, transport, and processing e.g. of oil seeds to oils and sugar crops to sugars.
- In this paper we use data/analyses from i) Springmann et al (Nature, 2018), ii) Springmann et al. (BMJ, 2020), and iii) Global Nutrition Report 2021, in figures 1, 2, 3, 3.2, 4.2, 5.3, 6.2, 7.6, 8.2, 9.2, and 10.2 (see appendices 1-3). These data represent food consumption (not food supply). Figures 3.2, 4.2, 5.3, 6.2, 7.6, 8.2, 9.2, and 10.2 were created using data on planetary boundary impacts for each of the 8 countries from the GNR 2021 excel sheet (tab 'country environment').²⁶ Figure 3 was created using supplementary information from Springmann et al. (BMJ, 2020). The excel sheet ('global targets' tab), contains an overview of the country-level results in terms of current food consumption and various dietary shift scenarios in relation to planetary boundaries.²⁷

- iv): Most of the food group chapters contain environmental impact data for relevant foods from Poore and Nemecek 2018.²⁸ Figures 2.1, 3.1, 4.1, 5.1, 6.1, 8.1, 9.1, 10.1, 11.1, and 12.1 are drawn from Figure S13, which provides the proportion of GHG emissions, acidification, and eutrophication by stage of the supply chain by product. The supply chain in this dataset begins with the extraction of resources needed to produce inputs for agricultural production, the initial impact of choice by farmers, and ends at the retail store, the point of choice for consumers. The data set covers ~38,700 commercially viable farms in 119 countries and 40 products representing ~90% of global protein and calorie consumption. It covers five important environmental impact indicators: land use; freshwater withdrawals weighted by local water scarcity; and GHG, acidifying, and eutrophying emissions. Land use was calculated from inverse yield and occupation time. Occupation time is reduced by multiple cropping but increased by fallow requirements. Land use is seed, on- and off-farm arable and permanent crops, fallow land, temporary pasture, and permanent pasture. GHGs are CO₂, CH₄, N₂O to air, using IPCC AR5 100-year factors with climate-carbon feedbacks. Acidification is SO₂, NH₃, NO_x to air, and eutrophication is NH₃, NO_x to air, NO₃⁻, NH₄⁺, P, N to water; both using CML2 Baseline for characterisation factors²⁹. Freshwater withdrawals and scarcity-weighted freshwater withdrawals are both irrigation, drinking, pond, and processing water. These 5 environmental impacts are provided per kg of retail weight in most food chapters, in tables 2.1, 3.1, 4.1, 5.1, 6.1, 8.1, 9.1, 10.1, 11.1, and 12.1 (drawn from supplementary data file (aaq0216_datas2.xls), using mean quantities from tab 'results – retail weight'.³⁰

SECTION 1: BACKGROUND CONTEXT

Food production in the Nordic and Baltic countries

All 8 countries use varying amounts of land area for agriculture. Denmark uses the largest proportion of land for agriculture (66%), followed by Lithuania (48%), Latvia (32%), Estonia (23%), Iceland (19%), Finland (7%), Sweden (7%), and Norway (3%).³¹ There are some large variations in terms of land used for agriculture across the 8 countries (table 1). In all countries except Iceland and Norway, the largest proportion of agricultural land is used for temporary crops (defined by the FAO as crops with a less-than-one-year growing cycle, which must be newly sown or planted for further production after the harvest).

Table 1: Agricultural land use by country and type (2019)³²

	Cropland ³³ (1000 ha)			Land under permanent meadows and pastures ³⁴ (1000 ha)*
	Permanent crops ³⁵	Temporary crops ³⁶	Temporary meadows and pastures ³⁷	
Denmark	23.3	1794.4	524.6	206.7
Estonia	7.0	500.0	180.0	292.0
Finland	5.0	1237.0	786.0	24.0
Iceland	**	5.0	115.0	1751.0
Latvia	9.0	999.6	263.4	632.0
Lithuania	35.1	1886.4	258.5	728.0 ³⁸
Norway	3.1	318.2	482.5	179.1 ³⁹
Sweden	3.4	1320.7	1084.5	461.3

*Unless indicated by a footnote, land area under permanent meadows and pastures was not disaggregated into 'cultivated' and 'natural growing'. ** Data missing or unavailable in the FAOSTAT database.

Sweden has the largest area of farmland under organic production (613,964 ha), and Estonia has the largest share of total farmland under organic production (22%), followed by Sweden (20%), Latvia (15%), Finland (13%), Denmark (11%), Lithuania (8%), Norway (5%) and Iceland (0.4%).⁴⁰ Except for Iceland and Norway, all countries are part of the European Union, which has a goal as part of its farm to fork strategy for every member state to have a minimum of 25% of agricultural land under organic production methods by 2030.⁴¹ Member states can go beyond this target.

In some countries, temporary pastures and meadows occupy much greater areas in comparison to permanent pastures and meadows, as shown in table 1. As a comparison to the total area under permanent pastures and meadows, Denmark uses 2.5 times more land for temporary pastures and meadows, Finland uses 33 times more, Norway 2.7 times, and Sweden 2.4 times (Estonia, Iceland, Latvia and Lithuania all use less land for temporary pastures and meadows compared to permanent pastures and meadows) (table 1). In other analyses (i.e. not including the FAOSTAT used in table 1), some of the land area used for food production in the Nordic and Baltic region is categorized as semi-natural grassland – this is usually an area of land cleared of native forest and maintained as low productivity grassland through farmed animal grazing on a proportion of native vegetation species and cultivated species. One estimate indicates that such areas tend to be much smaller than those used for permanent grazing, and also smaller than the areas used for temporary grazing. Sweden has the largest area of semi natural grassland, occupying an area equivalent to 10% of its total agricultural land – and potentially more than twice the area used for permanent grazing.⁴² However, the process of defining semi-natural habitats and assigning them to land use classes in national registers differs among the Nordic and Baltic countries,⁴³ and as the FAOSTAT database does not include semi-natural pastures as categories of agricultural land, or natural growing permanent pastures and meadows for every country, it is not possible to obtain comparable land areas or food production outputs for such lands. Hence, we do not include semi natural grasslands as a category in our table of agricultural land (table 1).

Additional food production and supply data (including imports and exports) from the FAOSTAT database in relation to each food group and country are provided in each major food group chapter, in as much detail as possible taking into account the limitations of data collection and availability (see box 1 for more information on data sources and handling).

Food consumption in the Nordic and Baltic countries

The mean daily consumption of selected food groups for adults derived from national dietary surveys in the Nordic and Baltic countries is presented in table 2. The intakes differ between countries and also within countries, with large variances as described by Warensjö Lemming and Pitsi.⁴⁴ Although differences between countries might be due to differences in the definitions of the food groups, and also that the years and methods of data collection differ (between 2007 and 2020), the countries share not only similarities but also differences in food preferences.⁴⁵ The current Food Based Dietary Guidelines (FBDGs), which focus on population health outcomes, vary more or less between the countries when it comes to specific amounts or frequencies but in general all are partly based on the NNR 2012.⁴⁶ None of the Nordic countries are close to full adherence of their FBDGs. The surveys in different countries were conducted over a long time period (2007–2022) and might not be accurate reflections of current intakes. The observed intakes may have been influenced by changing food markets.⁴⁷ In Denmark for example, intake of whole grains has increased considerably compared to the amounts shown in table 2.⁴⁸

Table 2: Current food consumption (g/adult/day) per country and food group⁴⁹


	Den mark	Esto nia	Finl and	Icel and	Lat via	Lithu ania	Nor way	Swe den	NNR2012 advice*
Cereals#	218		130	211	221	223	259	190	Depends on wholegrain content
Bread	138	71	92	81	83		184	87	
Whole grains	54			56			68	42	Increase (replace refined cereals)
Fruits and berries**	184	217	162	98	146	161	178	128	Increase
Vegetables* *	199	145	184	114	216	185	155	176	Increase
Pulses***	1	6	13	5	17		4	12	Increase
Potatoes	92	94	74	72	105		67	88	-
Fish and seafood#	37	25	32	45	31	28	67	39	Increase
Total meat#	161	81	145	117	178	140	147	110	
Red meat	134	44	105	82	88	89	119	90	Limit
Pig meat	87		28	7		85			Limit
Poultry	27	19	39	35	49	49	28	22	-
Dairy (without cheese)	304	266	397	245	195	119	314	245	Replace with low fat types
Cheese	44	17	41	42	9	12	44	25	Replace with low fat types
Eggs	24	22	24	20	35	28	26	14	-
Nuts\$	5	4	8	6	5		5	5	Increase

* Qualitative advice from NNR 2012 (table 1.1).⁵⁰ **The food group 'Fruits and berries' does not include juice, while 'Vegetables' includes pulses, but excludes potatoes. *** dry weight, # The food group Cereals contains both food ingredients and prepared foods such as bread (for Denmark also cakes and biscuits). Meat and fish are reported in raw weight, except sausages, cold cuts, and canned/smoked fish products. Total meat includes red meat and poultry – both uncooked and processed. Red meat and poultry in Estonia does not include processed meat. Red meat in Latvia does not include sausages. \$ incl. seeds - if not included in bread.

Food consumption in relation to environmental sustainability targets

From an environmental sustainability perspective, the impacts of current diets in each of the Nordic and Baltic countries mostly exceed the levels that would be required to meet a range of global targets related to greenhouse gas emissions, cropland use, blue water use, nitrogen use, and phosphorus use (Figure 3). A value above 1 (indicated by the red dashed line) exceeds the environmental target (or is consistent with the target being exceeded in future) and can be considered disproportionate in the context of an equitable distribution of environmental resources and mitigation efforts, and in effect, represents a country outsourcing its responsibility towards fulfilling the target⁵¹, whereas a value or 1 or less is consistent with remaining inside planetary boundaries. For example, these calculations indicate that, if everyone in the world consumed the same average diet (indicated by “benchmark”) as currently consumed in Iceland, the food related component of the global GHG target (consistent with keeping warming below 2°C) would be exceeded by more than 5-fold. Shifting to current FBDGs would mostly improve the outcomes, but not enough. For example, if everyone in the world consumed the FBDG from 2013 for Denmark, the food related component of the global GHG target (consistent with keeping warming below 2°C) would still be exceeded by almost 4-fold (Fig. 3). Hence, not only are current consumption patterns largely incompatible with global environmental targets, but current FBDGs could also be improved to align with global environmental targets. It should be noted that some of the environmental impacts demonstrated in figure 3 occur in exporting countries, as the data represent consumption in the Nordic and Baltic countries.

Figure 3: Impacts of current (benchmark) diets and adopting current national dietary guidelines in relation to the food-related portions of global environmental targets: by country and environmental metric⁵²



The analysis utilizes country-specific environmental footprints described in Box 1 and appendices 1-3.

Despite the varying approaches to assessing the environmental sustainability of current and future dietary patterns, the common conclusion is that environmental impacts generally decline as the amounts of animal products in the diet also decline – with vegan diets having the smallest environmental impacts, and diets highest in products from ruminant animals (cows, sheep and goats) having the largest environmental impacts.⁵³ This conclusion is also supported by systematic literature reviews⁵⁴ and original studies from the Nordic countries.⁵⁵ Production locations and methods, and food selection may add some nuance, for example in terms of the importance of blue water use related to some imported vegetables, fruits, and nuts in some of the Nordic countries.⁵⁶ The comparative approaches of the various dietary analyses of environmental impacts (i.e. Life Cycle Assessment, thresholds and systems) are described in the previous paper in this series and hence will not be described again here.⁵⁷ In this paper, we focus exclusively on the environmental sustainability aspects of individual foods within food groups, and not dietary patterns. A combined assessment of nutritional aspects and environmental aspects of dietary patterns is considered in an upcoming paper of the series, where nutritional adequacy is assessed in the context of a whole diet rather than individual foods or food groups.⁵⁸

General points across the food groups

A number of general, cross-cutting points apply across the foods and food groups and are therefore described in the following sections to avoid repetition in the food chapters.

Factors shaping environmental impacts

Environmental impacts are influenced by location and production methods, which can be determined by national contexts and policies (e.g., whether irrigated/rainfed, fertilizer types and amounts, plant protection types and amounts, land clearance, mono/poly cultures, crop rotations, manure management, tillage), processing (e.g., slaughter, cooking), packaging, and transportation (which can include refrigeration) type and distance (e.g., freight by road, rail, sea or air). Hence, there can be significant variation in environmental impacts within the same food product depending on different practices and choices, primarily during production. For example, per unit of food production, due to their higher energy requirements, trawling fisheries and recirculating aquaculture can result in 3 times more GHGs than non-trawling fisheries and non-recirculating aquaculture; due to lower macronutrient densities and digestibility of feeds used in grass-fed systems, grass fed beef requires more land and tends to emit more GHGs than grain-fed beef; and, in comparison to open field production, greenhouse production can have lower land requirements, higher yields, but almost 3 times higher GHGs (depending on heating and lighting usage, and energy source).⁵⁹ However, the greatest differences in terms of environmental impacts tend to occur between foods and food groups, regardless of production practices.⁶⁰ For example, in terms of GHGs, land use, energy use, acidification potential, and eutrophication potential, ruminant meats can have impacts 3–10 times higher than other animal-based foods and 20–100 times higher than plant-based foods on a per unit of weight basis.⁶¹

The denominator used to express environmental impacts of a food can influence interpretation. For example, due to their high protein content, tree nuts, oils, pulses, rice, soybeans and wheat are more efficient when measured by protein rather than energy, whereas cassava and sugarcane are more efficient by energy rather than protein content.⁶² While food waste is a major issue globally

that must be addressed, it is not covered in this paper as it is assumed to be a known problem that applies across food types.

Technology

We are aware that there are numerous technologies designed to decrease the environmental impacts of agriculture, including livestock production. For example, several initiatives designed to reduce the environmental impacts of ruminant meat production are underway in the Nordic countries and elsewhere, including feed additives to reduce methane emissions and improved manure management. Technologies under development include cell culture and precision fermentation. For example, microbial proteins⁶³ could provide environmental benefits compared to animal-based proteins, and recombinant proteins synthesised by microbes could reduce environmental impacts compared to milk⁶⁴ and egg white proteins.⁶⁵ Cell-culturing technologies can also be explored as a way to produce fats synthesized by microbes.⁶⁶ While such technologies are all likely necessary to some extent, they are insufficient to deliver the scale and pace of transformation needed to meet planetary health goals – this necessitates changes in food consumption, and reducing food loss and waste.⁶⁷ Therefore, we concentrate our assessment on consumption shifts that are achievable in the short term (with appropriate reshaping of markets and political and public support⁶⁸).

Supply side interventions

While the majority of environmental impacts of a food group or product often occur during the production (on farm) stage, there can be options for reducing impacts throughout the supply chain, including the processing and transport stages of the life cycle.⁶⁹ For example, reducing GHGs from transport through improved loading logistics and reducing packaging amounts. In most food chapters we demonstrate the variation in impact type and where it occurs along the supply chain for a range of foods. However, we assume that options for reducing the environmental impacts of production, processing and transport (such as precision agriculture and other technological approaches) apply across the food groups and will therefore not necessarily be discussed in each individual food group. Existing technologies that are not currently implemented uniformly across food production include precision agriculture. Precision agriculture, a method used to apply inputs in a direct rather than diffuse way in appropriate amounts at the correct times, could be important for minimizing inputs such as nitrogen and water use. The environmental benefits of increasing input efficiency through such technologies would be greatest in the least efficient systems.⁷⁰

Minimizing environmental impacts in a relative way i.e., per unit of food produced, does not necessarily equate with environmental sustainability – an assessment will still be required to ensure the impacts of production are consistent with environmental goals. In addition, impacts will increase if relative impacts are reduced but absolute impacts increase due to an increase in production. Ultimately, while reducing environmental impacts throughout the food supply chain is necessary, this alone cannot achieve the absolute reductions required at the system level to align with environmental goals. Changing the types and quantities of food grown and consumed is also necessary.⁷¹

Production methods

Using crop rotations and intercropping are considered important aspects of more environmentally sustainable food production, and directly overlap with sustainable food consumption in some cases. For example, combining grains and pulses during production and consumption is beneficial for farming practices and human health.⁷² Conservation tillage and cover cropping, particularly with

nitrogen fixing crops, reduces the requirement for nitrogen inputs and increases fertilizer input efficiency by reducing nutrient loss from agricultural systems.⁷³

Organic food production can result in beneficial outcomes, including higher levels of associated biodiversity in comparison to non-organic production.⁷⁴ For example, on-farm and near-farm biodiversity tends to be higher in organic agricultural systems, likely a result of lower fertilizer, herbicide, and pesticide use as well as through creating a more diverse landscape.⁷⁵ In addition, the use of manure as fertilizer can promote higher soil organic carbon in organic systems. However, due to land use and land clearing requirements, applied on a large spatial scale organic agriculture would likely have a net negative impact on biodiversity and soil organic carbon.⁷⁶ Therefore, while a shift to organic production at the global level could provide sufficient food availability, it is only feasible with a reduction in food waste and a shift to mostly plant-based diets to accommodate higher land use requirements (due to a reduction in yield, which is 34% lower when conventional and organic systems are most comparable.⁷⁷), without extending agricultural area.⁷⁸

Comparing the environmental impacts of conventional and organic food production at a local scale, a meta-analysis of LCAs found that, per unit of food produced, organic systems have higher land use requirements and eutrophication potential, lower energy use (due to less reliance on synthetic fertilizer and pesticides), and no difference in GHGs and or acidification potential. The differences are largely a result of differences in nutrient management. Manure (which organic systems largely depend on for nitrogen) releases nutrients in response to environmental conditions rather than crop nutrient demand, which can reduce nutrient assimilation by plants, and lead to reduced growth and yields – in turn increasing land use requirements. Nutrients not taken up by plants can also result in eutrophication and acidification.⁷⁹ However, there are techniques that can halve the land use difference between organic and conventional systems, including rotational farming, cover cropping, multi-cropping, and polyculture.⁸⁰

For some products, the difference in environmental impacts between organic and conventional production is relatively small, for example GHGs from potatoes⁸¹ and land use requirements for legumes and perennial crops.⁸² However, yield can be substantially lower in organic production – and can be further lowered if large portions of the harvest is lost or rejected (for example, if it does not conform to certain size, appearance or shape standards, or due to higher vulnerability to pests and diseases). For example, organic banana plantations give around 50% lower yield compared to conventional production.⁸³ Variety can also have an impact – for example, specialist varieties of tomato can have 3 times higher GHGs per kilogram of product than conventional varieties due to lower yields.⁸⁴

The solution might not be so simple as ‘organic or conventional’ or ‘pesticides or no pesticides’. Intercropping, agroforestry, and integrated pest management could play important roles – although the evidence for how this would operate in the Nordic region, with cold temperatures, needs to be further explored.⁸⁵ Crop rotations with brassicas to sanitise soil for cereals is considered important – but more knowledge about crops is needed, to identify which grow best together and which crop rotations are most successful – taking into account the practicalities for food producers i.e. establishing what can actually be grown. The choice between organic and non-organic also relates back to the choice of, or weighting of, metric. For example, organic farming has more benefits for biodiversity (e.g., underground biodiversity⁸⁶) – but requires more land to produce the same amount therefore, widespread uptake of lower-yielding, but nature-positive farming, requires commensurate changes in dietary composition.⁸⁷ It is likely that a sustainable food system will need to integrate the benefits of conventional, organic, and other agricultural systems.⁸⁸

Location and context

Due to a lack of comprehensive datasets on the environmental impacts of food production or consumption for each of the Nordic and Baltic countries for the range of foods covered in this paper, to demonstrate the environmental impacts of the foods within the NNR food groups, we used a global dataset of food Life Cycle Assessments (LCA) that has standardized boundaries to indicate a range of environmental impacts per unit of food production⁸⁹ (see box 1 for more details on data usage).

The spatial distribution and concentration of different pressures related to food production varies on land and in aquatic environments. A recent analysis found that while global impacts of food production on habitat disturbance, GHGs, nutrient pollution and freshwater use are dominated by land-based animal agriculture (with the greatest burden from pig meat closely followed by cattle meat), an estimated up to >10-fold variation in cumulative environmental efficiencies exist among countries for many livestock, fisheries, and crop products. For example, the efficiency of producing the same crops can vary 4 to 18 times among countries due to differences in water consumption, fertilizer/pesticide use, and farming practices. Similarly, efficiencies for marine fisheries vary up to 22-fold among countries depending on the specific species fished and equipment used within a country.⁹⁰ Therefore, finer-scale analyses are useful to identify where environmental pressures are located and how the environmental efficiency of production might vary among regions. Spatial cumulative footprint assessments explore where and how much each type of food contributes to food's total environmental footprint. The impacts depend on what is being displaced (for example, forest, wetlands), the sensitivity of systems to specific pressures, and local biophysical and socioeconomic conditions.⁹¹ For example, if food production is spatially located in areas with the most suitable climatic and soil conditions for a crop, this can increase agricultural input efficiency and decrease environmental impacts but may have implications for how far food is transported.⁹²

Another issue related to location and context is the use of agricultural wastes and by-products as animal feeds, which has the potential to reduce the environmental impacts of livestock production by 20%⁹³ – but is dependent on food safety/environmental regulations in each country or location.

Biodiversity impacts

Generally, biodiversity can be considered as a broad concept that includes diversity at the genetic, species and ecosystem level, and an essential component of functional ecosystems. The available comparative data for LCA used in this paper does not incorporate metrics for the impact of production on biodiversity, partly because different studies are highly heterogeneous in what they assess as “biodiversity”, and partly because of very significant context- and scale-dependency. For the former, different taxonomic groups (birds, plants, insects, microbes) may respond very differently to field-, farm- and landscape drivers.⁹⁴ For the latter, there are significant geographic and climatic effects that constrain a farm's biodiversity, which interact with the nature of the surrounding landscape to determine what may be found there and how agricultural practices may impact.⁹⁵ Biodiversity impacts are therefore difficult to measure and indicators and metrics largely rely on value judgements (for example, depending on which species or landscapes are considered of most value to humans), and the availability of a baseline comparison. The decisions around conservation or protection of certain species could differ depending on the values of the decision maker/s.

Farming impacts biodiversity in different ways and is very complex to assess at the system level and more so at the product level, making comparisons with other environmental metrics difficult. Monocultures reduce biodiversity both in terms of crop types, and by limiting the types of habitats

and food availability needed to support a diverse range of wildlife.⁹⁶ Areas where the same crops are grown every year lack the beneficial decontaminating effects of crop rotations, resulting in an increased requirement for plant protection products. Monocultures at a landscape level can have a range of impacts, including on the fauna and flora.⁹⁷ For some crops, genetic diversity within the crop itself is important for reducing disease, e.g., potatoes.⁹⁸ Producing food with less pesticide use would be positive for biodiversity. Since the 1950s pesticide use has had major impacts on soil and above ground biodiversity. Across Europe, pesticide and fertilizer use has been identified as the main pressure for bird population declines across the vast majority of common birds - especially invertebrate feeders but also farmland species, long-distance migrants and woodland birds.⁹⁹ Taking a cautious approach and applying pesticides and fertilizers only when absolutely necessary could help – via precision agriculture methods for example. However, to maximise biodiversity gains, it is important to provide the necessary space and habitat for wildlife in the landscape – reducing the use of pesticides and fertilizers alone will not be sufficient.¹⁰⁰

That biodiversity is not covered in the LCA data presented in the food groups is not to detract from its local and global importance. Biodiversity underpins a wide range of supporting and provisioning ecosystem services (from soil fertility, carbon storage, pollination, and natural pest control) as well as having cultural and social value. Across Europe – and globally – wildlife has been in significant decline for decades, and the bulk of this is caused by agriculture interacting increasingly with climate change.¹⁰¹ Reducing the demand for, and pressure on, land through changing the composition of diets may allow more environmentally beneficial farming systems to be adopted, and help to protect and restore globally important carbon and biodiversity repositories.¹⁰² The Kunming-Montreal Global Biodiversity Framework¹⁰³ was adopted in December 2022 (including by all 8 Nordic and Baltic countries). The framework contains 23 targets for 2030, a number of which have significant implications for food production and consumption and could thus raise the political importance of such issues in the coming years.

SECTION 2: FOCAL FOOD GROUPS

This section contains a chapter for each of the following NNR food groups: cereals; vegetables, fruits, and berries; pulses (legumes), fish, fish products and seafood, meat and meat products, milk and dairy products; and eggs.


NNR Food Group 3: Cereals

Global context

Globally, maize is the type of cereal produced in the largest quantity (1,162 million tonnes (mt)), followed by wheat (761 mt), rice (757 mt), barley (157 mt), oats (25 mt), and rye (15 mt).¹⁰⁴ In terms of use, cereals are a major source of livestock feed. Currently 42% of global cereal production is dedicated to feeding farmed animals, which alone uses 43% of global cropland.¹⁰⁵ As a proportion of global production, animal feed currently accounts for 17% of wheat, 4% of rice, 56% of barley, 61% of oats, 46% of rye, and 59% of maize production.¹⁰⁶ It should be noted that the ratio of cereals allocated to farmed animal feed versus direct human food is not uniform each year – variations can occur due to, for example, weather conditions and fertilizer prices.

Across the impacts shown in figure 3.1, most impacts for cereals occur during production i.e. on farm. Maize has the most land-use change associated with production, followed by wheat and grains¹⁰⁷. All impacts from losses are very similar across the cereals.

Figure 3.1: Proportion of environmental impacts from different life cycle stages: Cereals (global)¹⁰⁸



Under business-as-usual food consumption and expected population growth to 2050, the absolute environmental impacts of cereal production will increase, even if the efficiency per unit of production improves. For example, it has been estimated that a further two- to three-fold increase in nitrogen (N) supply will be required to support global food production for the anticipated population of ~9.7 billion by mid-century. Synthetic N applied to cereal production (wheat, rice and maize) accounts for more than 50% of the total fertilizer used globally for crops. Three countries consume more than half of this amount across the 3 cereals: China (22.7%), India (18.5%) and the US (14%).¹⁰⁹ However, there is substantial scope to reduce N use, given that only 35% of the 115 million tonnes of N applied annually to crops is actually taken up by them (the remaining 75 million tonnes of N, or ‘excess N’ discharges into the surrounding environment including waterways and atmosphere). Excess N varies significantly by region, with China having the largest share (33% excess N) of the global total, followed by India (18%), and the US (11%).^{110,111} Applying N in the correct quantities, at the correct time and in the correct place is an important method for reducing excess N. Such precision agriculture techniques can be assisted with the use of technology, and also require training and knowledge among farmers.

In terms of individual crops/products, maize meal tends to have the lowest environmental impacts across the cereals shown in table 3.1, particularly in relation to freshwater use. Paddy rice has the highest GHG footprint, which is primarily a result of methane emissions from waterlogged rice fields during the production stage. The majority (80%) of global rice production is cultivated in waterlogged fields, which are either irrigated or rainfed. While rice can be grown in dry conditions, the yield is much lower in comparison (around 33% of the yield from paddy rice).¹¹²

Table 3.1: Average environmental impacts per kg of retail weight: Cereals (global)¹¹³

	Land use (m ² /kg)	Greenhouse gas emissions (kg CO ₂ eq/kg)	Acidifying emissions (g SO ₂ eq/kg)	Eutrophying emissions (g PO ₄ ³⁻ eq/kg)	Freshwater (L/kg)
Wheat & Rye (bread)	3.9	1.6	13.4	7.2	648
Maize (meal)	2.9	1.7	11.7	4.0	216
Oatmeal	7.6	2.5	10.7	11.2	482
Rice	2.8	4.5	27.2	35.1	2,248

Substituting one cereal for another with lower environmental impacts could be an option for improving environmental sustainability outcomes. However, the intended use is important to consider. For example, where high protein and/or gluten content is required (e.g., wheat for bread making and pasta), this characteristic would need to be matched. In terms of environmental impact, high protein content tends to be associated with high application of N fertilizer¹¹⁴ – although not necessarily if grown with legumes due to their nitrogen fixing properties.¹¹⁵

Thus, it might be more appropriate to consider alternative food products rather than seek like-for-like replacements. Another potential option is to grow cereals in rotations with legumes to benefit from their biological N fixation and in turn reduce reliance on synthetic N fertilizer (i.e. obtained through the Haber-Bosch process).¹¹⁶


An expansion of cereal production is not inevitable, even taking into account the expected growth in the human population. The major route for reducing all environmental impacts of cereals is in conjunction with a shift to more plant-based diets (particularly in high- and middle-income countries), thus reducing the amount of cereals required for farmed animal feed. The demand for animal feed is partly a result of high levels of livestock production, but largely due to the inefficiency of converting crop nutrients to animal products. For example, an analysis based on US production estimated that to deliver 1 calorie of beef for human consumption requires 37 calories of plants, 1 calorie of pork requires 12 calories of plants, 1 calorie of chicken requires 9 plant calories, 1 calorie of eggs and 1 calorie of dairy each require 6 plant calories.¹¹⁷ Global averages for protein feed conversion efficiency (i.e. the % of protein in feed converted to protein in product) range from 4% for beef to 25% for eggs. The equivalent range for calories (i.e. the % of energy in feed converted to energy in product) is 2% for beef and 24% for whole milk.¹¹⁸ At the global level, 36% all calories produced from crops are fed to farmed animals with only 12% of those calories returning as livestock products for human consumption, such as meat and milk¹¹⁹ - equating to a loss of 32% of all crop calories produced due to this conversion process¹²⁰). The world's croplands could potentially feed billions more people by shifting from animal feed to producing food for human consumption.¹²¹ It would also be important to reconfigure crop production, to enable human edible crops to be

grown in place of animal feed crops. It has been estimated that by taking this approach in the US (i.e., replacing feed crops with healthy alternatives for human consumption, such as legumes, whole grains, fruits and vegetables) could feed an additional 350 million people without increasing the cropland area.¹²²

Nordic and Baltic context

Figure 3.2 shows the variation in terms of impacts of cereal consumption across the Nordic and Baltic countries and how they relate to global environmental limits for food consumption. For example, if everyone in the world ate the same amount of cereals as Lithuania it would use over 70% of the global nitrogen amount allocated to food consumption.

Figure 3.2: Impacts of cereal consumption across the Nordic and Baltic countries in relation to the food portion of global environmental limits (%)



Source: *Global Nutrition Report 2021*¹²³ The analysis utilizes country-specific food consumption and environmental footprint data and relates them to the food portions of global environmental limits i.e. a global test to assess the impacts if everyone in the world consumed at the given rate. The methods and data are described in Box 1, and presented in appendices 1-3.

Denmark is the biggest grain producer of the Nordic and Baltic countries, followed by Sweden and Lithuania (table 3.2). Grain production is sufficiently high to satisfy national consumption in every country, except Iceland and Norway, where there is some reliance (a strong reliance for Iceland) on imports to meet demand. Lithuania and Latvia are the largest exporters of cereals, followed by Denmark. It should be noted that production amounts and import demand is not uniform each year – instead, annual variations can occur.

Lithuania is the biggest wheat producer, followed by Denmark and Sweden. Denmark is the biggest producer of rye, followed by Sweden and Latvia. Finland is the biggest producer of oats, followed by Sweden and Denmark. Denmark is the biggest producer of barley, followed by Sweden and Finland. Lithuania is the biggest producer of maize, followed by Denmark and Sweden. None of the countries currently produce rice.¹²⁴ Globally, Finland and Sweden are within the top 10 largest oat producers (4th and 10th respectively), and Denmark is the 4th largest producer of rye.

Table 3.2: Cereal production and supply across the Nordic and Baltic countries in 2019¹²⁵

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance* (1000 tonnes)	Food supply (Kg/person/year)
Denmark	9,518	1,348	1,822	9,044	118
Estonia	1,630	165	1,162	633	96
Finland	4,036	327	716	3,647	119
Iceland	8	103	0	111	80
Latvia	3,163	861	3,223	801	115
Lithuania	5,252	796	4,298	1,750	147
Norway	1,324	885	41	2,168	124
Sweden	6,148	1,087	1,552	5,683	110

*The amount remaining for national use taking into account production, imports and exports (balance = production + imports – exports).

In every country except Estonia, Iceland and Sweden – more than half of cereal supply is used to feed farmed animals, and in every country animal feed accounts for a substantial share of cereal supply. Allocation is highest in Denmark which uses 75% of cereal supply for farmed animal feed, followed by Latvia (63%) and Finland (59%) (table 3.3).

Table 3.3: Cereal allocation to animal feed and direct human food across the Nordic and Baltic countries in 2019¹²⁶

	Feed (1000 tonnes)	Food (1000 tonnes)	Feed as % of total supply ¹²⁷
Denmark	6,791	683	75
Estonia	315	127	50
Finland	2,149	659	59
Iceland	44	27	40
Latvia	505	219	63
Lithuania	963	405	55
Norway	1,103	670	51
Sweden	2,662	1,102	47

Given that animal feed occupies a large proportion of supply in every country coupled with the general need to reduce livestock production and consumption of animal products, there is scope to explore reducing overall cereal supply and/or reallocating supply from animal feed to human food for direct consumption. Depending to some extent on local conditions and cropping systems, it might be possible for all countries to shift supply of cereals from feed to food. Table 3.4. describes the relationship between cereals/grains used as feed and that used as human food. The larger the number the larger the proportion of cereals used for feed compared to use for food. For example, in Lithuania 114 times more oats are used for farmed animal feed compared to human food. The feed used to food supply ratios shown in table 3.4 could guide the exploration, taking the size of the cereal food supply in table 3.5 and the production in table 3.2 into account. The option to shift cereal supply to human food for direct consumption would depend on the quality/grade of the cereal. Where the grade is not high enough for human consumption, there could be potential to increase the grade during production with additional resources, that might include skills and training for cereal producers.

Table 3.4: Feed to food supply ratio across the Nordic and Baltic countries in 2019¹²⁸

	Wheat	Rye	Barley	Rice	Oats	Maize	Cereals, other
Denmark	6.8	8.5	-	0.2	8.8	10.6	61.0
Estonia	3.0	0.5	16.1	0.0	3.2	1.0	27.0
Finland	0.7	0.0	21.6	0.0	36.6	-	1.9
Iceland	0.8	-	4.9	0.0	0.0	-	-
Latvia	1.5	0.8	4.1	0.0	15.6	14.5	2.6
Lithuania	0.5	1.5	-	0.1	114.0	101.0	49.6
Norway	0.4	2.7	38.7	0.0	3.5	-	-
Sweden	1.0	0.0	-	0.0	21.4	3.2	71.3

- Data unavailable to make the calculation.

Wheat is by far the main grain consumed in each country, followed by rye – except for Iceland where more rice and oats are consumed in comparison to rye, and in Norway where more oats and rice are consumed in comparison to rye. In comparison to wheat and rye, there is generally a relatively low consumption of oats and barley across the countries (table 3.5). Hence, there could be potential to increase the consumption of oats and barley in most if not all countries. Increasing/shifting to barley consumption could be a way to improve the environmental sustainability of cereals given that barley generally has the lowest environmental impacts across a range of metrics (substantially lower for some metrics, e.g., freshwater use) (table 3.1). While the current availability of freshwater across the 8 countries is generally not a limiting factor for food production, it is possible that some recently observed conditions (e.g., drought) could intensify in frequency and magnitude in the future due to the impacts of climate change. Also, the subsequent impacts (e.g, low availability of hydro power and high export leading to high energy prices) could increasingly impact food production. Shifting a portion of oats and barley currently used for animal feed to human food for direct consumption could particularly benefit Iceland which is currently dependent on cereal imports to meet supply (table 3.2), in turn helping to increase consumption of such cereals and potentially reducing import dependency.

Table 3.5: Cereal food supply (not including cereals produced for feed) across the Nordic and Baltic countries in 2019 (kg/person/year) by cereal type¹²⁹

	Wheat	Rye	Barley	Rice	Oats	Maize	Cereals, other
Denmark	85	18	0	5	4	5	0
Estonia	55	16	5	3	5	10	3
Finland	79	26	5	5	3		1
Iceland	71	1		4	3		0
Latvia	62	25	18	3	3	1	4
Lithuania	113	15	14	3	0	1	1
Norway	102	3	3	5	12	0	0
Sweden	87	10	2	6	2	2	0

Exploration of new crop options/reintroductions such as quinoa and millet could emerge as local growing conditions change and opportunities to cultivate different crops arise. For example, quinoa is now grown in Finland, quinoa and chickpeas are now grown in Denmark, and there is potential to resurrect millet production. There is scope to learn from similarly positioned countries that have experience growing such crops on a commercial scale (e.g., the UK¹³⁰). Breeding of locally adapted varieties and development of locally applicable cultivation practices are also needed to support the production.

Considerations for reducing environmental impacts of cereal consumption in Nordic and Baltic countries

From the overview of cereal production, consumption, and trade provided in this section, we identify a number of opportunities to explore as potential ways to reduce environmental impacts of the cereal food group:

- **Opportunities with major benefits:**
 - Cereals have among the lowest relative impacts, from an environmental sustainability perspective, which makes increased consumption an option for increasing environmental sustainability of diets.
 - Shift portions of animal feed to direct human food if possible (differences in cereal quality need to be determined and adjusted if needed, and crop reconfiguration to best support human health might be required). This method would allow cereal/grain consumption to increase, without increasing production (and the associated environmental impacts) and would need to be supported by a reduction in animal agriculture (see chapter 8). Increased consumption of whole grains is in general recommended for humans based on health evidence, including for nutrient supply such as fibre.¹³¹
 - Increase the proportion of cereals grown using less environmentally damaging methods (assisted by an overall reduction in cereals grown for animal feed). This could include organic methods (or similar), and increasing and improving crop rotations with legumes to reduce the requirement for synthetic N.¹³² Precision agriculture techniques could further reduce 'excess nitrogen' and other key resource requirements, including water.

Opportunities with relatively minor benefits:

- Explore new crop options and reintroductions (such as millet). As local growing conditions change, more opportunities to cultivate different crops might arise.
- Explore differences in environmental impacts from locally (i.e., within each country), regionally (i.e. within the 8 countries) and internationally sourced cereals to potentially identify further options for reducing environmental impacts, taking into account potential changes in environmental conditions and increasing occurrence of environmental shocks. Environmental impacts of imported cereals might vary depending on the types and amounts of pesticides and fertilizers that are permitted in the country of origin, the methods of production such as intensive farming with tilling versus no till

and organic (or similar) methods, and the conditions in the country of origin such as water shortages and soil degradation. Trade stipulations might be a useful tool for reducing the environmental impacts of imports.


NNR Food Group 4: Vegetables, fruits, and berries

Global context

At the global level, in terms of fruit, bananas are currently produced in the largest quantity (118 mt), followed by oranges (114 mt) and apples (87 mt). In terms of vegetables, potatoes (which are considered a separate food group in the NNR context) are produced in the largest amount (370 mt), followed by casava (300 mt) and tomatoes (180 mt).¹³³ Approximately 10% of global potato production and 30% of cassava production are used as farmed animal feed.¹³⁴

Figure 4.1 demonstrates the proportion of environmental impacts associated with different fruits and vegetables throughout their product life cycles. For some products, the production stage accounts for the majority of impacts e.g., brassicas and cultivated berries, while for other products and impacts, other stages are more important e.g., transport of bananas in relation to GHGs. Citrus production has the largest carbon sequestration benefits (through tree growth) per unit of product, followed by apples and bananas.

Figure 4.1: Proportion of environmental impacts from different life cycle stages: Vegetables, fruits and berries (global)¹³⁵



Root vegetables (including potatoes) and onions generally have the lowest environmental impacts per unit of weight, particularly in comparison with salad vegetables such as tomatoes. Root vegetables (including potatoes) and onions can be most easily stored, with relatively small inputs and little waste. In terms of fruits, apples and citrus tend to have the lowest associated environmental impacts, and also store well (table 4.1).

Table 4.1: Average environmental impacts per kg of retail weight: Vegetables, fruits and berries (global)¹³⁶

	Land use (m ² /kg)	Greenhouse gas emissions (kg CO ₂ eq/kg)	Acidifying emissions (g SO ₂ eq/kg)	Eutrophying emissions (g PO ₄ ³⁻ eq/kg)	Freshwater (L/kg)
Potatoes	0.9	0.5	3.9	3.5	59
Cassava	1.8	1.3	3.4	0.7	0
Tomatoes	0.8	2.1	17.2	7.5	370
Onions & Leeks	0.4	0.5	3.6	3.2	14
Root Vegetables	0.3	0.4	2.9	1.6	28
Brassicas	0.6	0.5	8.2	5.0	119
Citrus Fruit	0.9	0.4	4.0	2.2	83
Bananas	1.9	0.9	6.4	3.3	115
Apples	0.6	0.4	3.5	1.5	180
Berries & Grapes	2.4	1.5	12.3	6.1	420

There are a range of factors that influence the environmental impacts of fruits and vegetables. Production practices can vary highly within products and even within countries (for example, use of pesticides and fungicides varies significantly even across EU countries). The scale of production (from small holder to industrial level) can have a significant influence. Generally, larger scale production has higher yields and more efficient production, which can be used as a metric of environmental impact per unit of production (but not on biodiversity). However that is not always the case and is potentially short-term if for example soil degradation prevents the maintenance of such yield levels. For example, highly mechanized production, such as in orange groves, can increase fuel use and GHGs, and a high application of synthetic fertilizer can result in pollution of nearby waterways.

Some crops are typically produced under a certain method. For example, bananas, citrus and tomatoes are largely produced in intensive monoculture plantations for the export market, which have a range of associated issues such as use of herbicides to remove other vegetation from the land (leading to soil erosion and run-off), and larger requirements for plant protection products (such as chemical pesticides) and fertilizer. Potato production tends to involve more intensive soil tilling compared with other crops, which releases GHGs and reduces soil organic matter content, making organic fertilisers particularly important for crop rotations involving potatoes.¹³⁷

In general, more chemical plant protection products are used in the production of fruits and vegetables than other types of agricultural production (in terms of per hectare and kg of harvested product), and tends to be higher in intensive fruit and berry production compared with vegetables.¹³⁸ While pesticide use is mostly concentrated during the production stage, for some fruits it is also applied at other stages e.g., fungicides applied to bananas for transportation, and some are applied to the soil e.g., soil disinfectant for strawberries to prevent mould. Production in warmer locations can also require higher levels of pesticide application. Some crops require plant protection to enable sufficient yields, for example fungicides are applied to control potato blight.¹³⁹ Hence, yields from organic potato production are significantly lower – potentially up to 50%. The use of climate-controlled greenhouses can reduce pesticide requirements, as can crop rotations (which can also help reduce the occurrence of potato blight).¹⁴⁰

In comparison to outdoor production, greenhouses can produce greater quantities of food on a smaller area over a shorter time period. Food production in greenhouses can also reduce the reliance on having land available with appropriate soil quality, which is a limiting factor in some

locations e.g., Norway, and can be more efficient with respect to water use. There can be substantial differences due to production methods, even with greenhouse production. For example, GHGs from organically produced tomatoes can be 40% higher due to a reduced yield in comparison to conventional production (everything else being equal in terms of energy to heat and light the greenhouse).¹⁴¹ A major trade off compared with outdoor production is the energy requirements for heating (and lighting and sometimes cooling) the greenhouse and the subsequent GHGs,¹⁴² versus a greater requirement for plant protection products outdoors. However, this trade off could be reduced if renewable energy sources or waste heat are used to power the greenhouse - or with greater energy efficiency - or if crops with lower demands for heating/cooling were cultivated. In addition, food production in greenhouses generally requires less resource input in the form of fertilizer and pesticides compared to outdoor production¹⁴³ and can protect crops to a certain degree from climate extremes, thereby reducing crop losses in extreme weather events.


The impacts related to harvesting, processing and losses incurred during the life cycle stages can also vary substantially. For example, higher levels of waste result from consuming orange juice compared to whole oranges. The types of equipment used during harvesting can also be important – for example, energy use in apple production tends to be highest where hydraulic ladders/platforms are used. Minimising losses during storage is particularly important.¹⁴⁴

While the environmental impacts of transport tend to account for the smallest portions of overall impact, the proportion of overall impact can be more significant for fruits and vegetables compared to animal products for example. In general, transport via rail and sea are less impactful compared to air and road freight, however the actual impacts of all types of food also depend on travel distance, load and requirements for refrigeration. Seasonal consumption of locally produced fruits and vegetables can help to reduce the requirements for longer distance transport of fruits and vegetables with a short shelf life, and/or that require refrigeration. For some crops, it is possible to extend the season through selection of multiple varieties e.g., strawberries that fruit at different times, in turn reducing import requirements. Conversely, where growing conditions are particularly challenging, importing fruits and vegetables such as tomatoes, lettuces and cucumbers from warmer regions could be the lower-impact option in terms of GHG emissions for example, if the full life cycle is considered (i.e., not just comparing the transport impacts). For other impacts, such as water and pesticide use, production in the Nordic/Baltic countries could be more sustainable than imported products.

Nordic and Baltic context

Figure 4.2 shows the variation in terms of impacts of fruit and vegetable consumption across the Nordic and Baltic countries and how they relate to global limits for food consumption. For example, if every country in the world consumed the same as Estonia, it would use 30% of the global limit in terms of nitrogen allocated for food consumption. Given that the consumption of this food group needs to increase across the Nordic region, reducing the environmental impacts of fruits and vegetables would be beneficial, particularly in relation to nitrogen use and pesticides.

Figure 4.2: Impacts of fruit and vegetable consumption across the Nordic and Baltic countries in relation to the food portion of global environmental limits (%)



Source: *Global Nutrition Report 2021*¹⁴⁵ The analysis utilizes country-specific food consumption and environmental footprint data and relates them to the food portions of global environmental limits i.e. a global test to assess the impacts if everyone in the world consumed at the given rate. The methods and data are described in Box 1, and presented in appendices 1-3.

Generally, there is significant potential to increase fruit and vegetable production across the Nordic region (including Iceland). In terms of fruit – the production of apples, pears, cherries, currants and plums could potentially be increased. Strawberries, raspberries, bush blueberries, buckthorn and other berries grow well in the local climate. The potential to harvest more wild berries might be substantial (most wild berries in Norway, Finland and Sweden¹⁴⁶ are unpicked, for example). However, there are no data on the amount of wild berries that are accessible and could be harvested sustainably, the potential consumer demand for wild berries, or the potential impact on wildlife that might utilise the berries as a food source. Incentivising the harvest of wild berries could also be difficult (some countries such as Finland currently rely on seasonal labour from migrant workers) – but there is potential to link this with health and recreation activities rather than having a sole purpose of food harvesting. Wild herbs and plants such as nettles might also have some potential within the region. Root vegetables, including potatoes, and brassicas could be increased. The main limitations across the region are availability of arable land, a short growing season and the need for a long storage period (with methods that reduce losses). Despite this, there are estimates to suggest that in Norway, under crop rotations and optimal use of agricultural soils, vegetable production could be increased by around 5 times and potatoes could be increased by around 7 compared to current production levels.¹⁴⁷

Increasing climate-controlled greenhouse production could reduce the requirement for plant protection, and help to overcome limits of outdoor fruit production across the region.¹⁴⁸ However, energy use could become an issue but this could evolve if the input from renewable energy sources increases and the energy efficiency of climate-controlled greenhouse gas production improves, making greenhouse production more viable (although reducing energy demand overall rather than replacing all fossil fuel energy with renewable sources should be a key step). Energy and food prices have been identified as important variables in the Norwegian context, and the provision of government support (investment, electricity costs), that is not unlike other aspects of government

support (e.g., import protections; subsidised livestock production) to enable adoption.¹⁴⁹ Iceland already uniquely benefits from substantial climate-controlled greenhouse production, facilitated by geothermal heat and hydropower.¹⁵⁰

Identifying the extent to which increasing crop diversity and agroforestry can reduce pesticide use and improve other environmental sustainability outcomes requires more study across the region. Based on a number of trial farms in Sweden, it could be viable to implement agroforestry to grow nuts, berries and fruits in agricultural systems.¹⁵¹

There is a good potential for rain fed agriculture in the Nordic and Baltic region, with some possible requirement for irrigation during the summer season. While there are currently no major water stress issues in the Nordic and Baltic countries, there are several other considerations regarding increasing fruit and vegetable production. An increase in labour would be required, particularly for methods such as spatial intercropping on a small scale. Proximity to markets and/or processing facilities could be important for some crops, and profitability for farmers could be an issue. Competition with cheaper imported products, lack of agreement between producers and retailers, and toll barriers are also an issue – for example, domestic production needs to compete with imported products. The variation in trade support could therefore influence crop types. Also identified as important in the Nordic context is the preservation of certain types of food (from a normative or cultural perspective), which could play a role in influencing the viability of different crop types. There are some important locational perspectives, such as the policies to take peatland out of agricultural production in Denmark, Norway and Finland. Although policy effectiveness is in turn a large determinant of whether such locational considerations impact food production – for example, a law to prevent peatland conversion in Norway has been found to be largely ineffective.¹⁵²

Currently all Nordic and Baltic countries have a substantial dependency on imports to meet domestic demand for fruits and vegetables. For some countries, such as Iceland, a high dependency has been placed on air freight. Increasing consumption of vegetables and fruits is necessary to obtain healthy and sustainable plant rich diets, and this requires either an increase in local production, a change in consumption (i.e., to eat more seasonally available products), an increase in imports, or a combination of those factors. There could be a cultural expectation among consumers to continue importing food in large proportions to maintain purchasing possibilities (such as strawberry availability during the winter), and to see a continuation of current land use nationally (such as high levels of forest cover), rather than more conversion to agriculture. One possible factor that might help to maintain such expectations is that the environmental impacts of imports are not seen or experienced directly by consumers in importing countries. However, this should be contrasted against concerns regarding preparedness for trade disruptions and resilience to environmental shocks. Some level of domestic food security considerations should be integrated into food system planning, to avoid an expectation among rich nations that they will always be able to ‘buy themselves out of food supply problems’.¹⁵³ Such considerations could also be extended to imported supplies – for example, decreasing imports from water-scarce regions (e.g., in Spain) and regions that are likely to become water stressed.

Given the various trade-offs and variation in impacts, identifying which crops or production methods or locations are ‘best’ for environmental sustainability depends on the metric or metrics given most weight – which might be determined, for example, by a pressing environmental issue such as drought, or a value judgement. There is a lack of knowledge about the production systems and their environmental impacts for imported fruits and vegetables, which further complicates the process. Including issues, such as social impacts, in addition to environmental sustainability could also change the weighting/priority order. Furthermore, considering production systems over the long term is also

important – for future generations and to account for environmental changes. For example, climate change is expected to shift pest and disease ranges suggesting that future farming systems, and crop suitability, will change in the Nordic region. More frequent periods of drought could enhance water pollution and create water scarcity issues. Therefore, a focus on groups of fruits and vegetables that are more resilient to climate and environmental shocks, and store well (such as potatoes and apples) could be most appropriate.

Fruits and berries

All countries have a substantial dependency on imports to meet domestic fruit and berry supply. Lithuania is the least dependent on imports, with a 4-fold ratio of production compared to balance (taking into account imports and exports) (table 4.2).

Table 4.2: Fruit and berry production and supply across the Nordic and Baltic countries in 2019¹⁵⁴

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	47	546	157	436	60
Estonia	4	103	5	102	75
Finland	31	420	14	437	73
Iceland	0	31	0	31	87
Latvia	17	184	76	125	49
Lithuania	42	345	213	174	47
Norway	33	397	2	428	77
Sweden	41	824	91	774	59

For all countries, other fruits (a group which includes pears, quinces, apricots, cherries, peaches, plums, strawberries, raspberries, gooseberries, currants, blueberries, melons, figs and mangos) are the most highly consumed. Except for Estonia, this is followed by oranges, bananas and apples, which are the most consumed single fruit in Estonia (table 4.3).

Table 4.3: Fruit and berry food supply across the Nordic and Baltic countries in 2019 (kg/person/year)¹⁵⁵ by fruit type

	Other fruits	Bananas	Oranges	Apples	Grapes	Pineapple
Denmark	22	11	10	3	7	3
Estonia	21	14	15	16	5	2
Finland	21	18	16	9	4	3
Iceland	28	9	14	13	5	3
Latvia	17	11	9	4	2	1

Lithuania	15	8	11	6	2	1
Norway	22	15	17	13	5	2
Sweden	21	7	16	7	3	2

Vegetables

All countries have a substantial dependency on imports to meet domestic vegetable supply. Lithuania is the least dependent on imports, with a 1.6 fold ratio of production compared to balance (taking into account imports and exports) (table 4.4).

Table 4.4: Vegetable production and supply across the Nordic and Baltic countries in 2019¹⁵⁶

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	261	420	79	602	96
Estonia	25	97	9	113	81
Finland	263	253	8	508	85
Iceland	5	23	0	28	77
Latvia	62	175	52	185	89
Lithuania	178	232	116	294	96
Norway	195	244	1	438	75
Sweden	310	699	55	954	84

Except for Latvia and Lithuania, other vegetables (a group which includes cabbages and other brassicas, artichokes, asparagus, lettuce, spinach, pumpkins, peppers, carrots, mushrooms and frozen, dried and preserved vegetables), are the most highly consumed across all countries, followed by potatoes (which are the most consumed vegetable in Latvia and Lithuania) (table 4.5).

Consumption across the countries is generally focused on very few products (e.g., cucumber and salad vegetables), hence there is scope to expand the diversity of vegetable consumption.

Table 4.5: Vegetable food supply across the Nordic and Baltic countries in 2019 (kg/person/year)¹⁵⁷ by vegetable type

	Other vegetables	Tomatoes	Onions	Potatoes
Denmark	76	14	7	61
Estonia	62	13	7	57
Finland	62	15	8	58
Iceland	59	12	6	39
Latvia	69	12	8	112
Lithuania	69	18	9	84
Norway	62	10	3	49
Sweden	61	16	8	55

Considerations for reducing environmental impacts of fruit, berry and vegetable consumption in Nordic and Baltic countries

Even though vegetables, fruits and berries have among the lowest relative impacts, from an environmental sustainability perspective, based on the overview of production, consumption and trade, we identify a number of opportunities to explore as potential ways to reduce environmental impacts of the fruit, berry and vegetable food group. As consumption of this food group would need to increase to meet dietary guidelines (table 2), the opportunities relate to reducing environmental impacts per unit of production, or relative impacts, rather than reducing total, or absolute, impacts from this food group. Our suggestions span across production and consumption aspects and could have relatively major or minor impacts depending on the level of uptake/implementation:

- Diversifying consumption of fruits and vegetables could reduce the dependency of imported salad vegetables during the winter months and import of fruits during the summer months. This would include a decreased consumption of tomato, cucumber, pepper and lettuce during the winter and restricting consumption of these products more to the summer and autumn. During the colder months (winter and spring), a greater consumption of root vegetables (e.g., carrot, parsnip, celeriac, swede and beetroot), brassicas (e.g., cabbage) and onions could potentially reduce the need for salad vegetables, in addition to using vegetables that preserve well to enable longer storage and use. Even where imports of vegetables are necessary during the winter months, importing Chinese cabbage, onions and root vegetables has a lower GHG impact than importing salad vegetables, due to increased storability and reduced risk of waste during transport and post-retail. In terms of fruit, this could involve an increase in tree fruit (e.g., apples, pears, cherries, plums) during the colder months and wild berries during the summer months. However, the potential impacts on local wildlife that rely on wild berry consumption should be considered. Estimated harvest levels are low across the region.¹⁵⁸ However, the estimated harvest levels are variable and uncertain, and if berry utilization should increase drastically, the impacts on biodiversity may need to be assessed.
- Increasing the supply of tree fruits (such as apples and pears) produced within the region could help reduce the import of citrus, bananas and grapes. Increasing the domestic cultivation of a wider variety of apples would also help diversify the agricultural landscape, and extend the season due to variations in harvesting, ripening and storage times (which could potentially be shortened). Tree fruits have the additional environmental benefit of carbon sequestration and storage and could be used in agroforestry systems.
- Increasing the use of climate-controlled greenhouses (or underground, thermally insulated or vertical agriculture) could help to increase production of fruits and vegetables throughout the year within the region and reduce the requirement for plant protection products and water use in water scarce regions (e.g., in importing countries such as Spain). The potential for this measure could be highest in countries where the proportion of renewable energy is the highest and could potentially increase over time if the proportion of renewables in the energy mix increases.

- Root vegetables (including potatoes) and onions generally have the lowest environmental impacts per unit of weight and can be most easily stored, with relatively small inputs and little waste. To reduce the requirement for potato blight treatment and related loss of crop, crop rotation and genetic diversity are known options.
- Increasing the proportion of fruits and vegetables (of both imports and domestic supply) grown using less environmentally damaging practices, such as organic methods, and intercropping, would help to reduce the use of plant protection products and synthetic fertilizer, and potentially provide biodiversity benefits by for example increasing landscape heterogeneity.
- One way to reduce the overall impact of fruit and vegetable production is to increase the proportion that reaches the market by reducing the stringency of shape/size/quality requirements i.e., increasing the proportion of 'ugly' fruits and vegetables available for consumer purchase. The practice of removing products from the market that do not conform to pre-set standards is greater for fruits and vegetables compared to cereals and dried legumes – hence there is most potential to utilise this measure as a way to reduce the environmental impacts of fruits and vegetables.
- Some environmental issues such as soil health and biodiversity, are not captured by LCA. Hence, additional assessments would be required to avoid soil damage and biodiversity loss and ensure for example that the richest soils are utilised for the highest yields, using methods that protect soil and allow for sustained production over time.


NNR Food Group 5: Pulses (legumes)

Global context

Pulses are dried seeds from legume plants and include chickpeas, lentils, dried peas and dried beans. Legumes include soya beans, fresh beans and fresh peas.¹⁵⁹ At the global level, 336 mt soya beans, 26 mt dry beans, 23 mt green beans, 14 mt chickpeas, 14 mt dry peas, 9 mt dry cow peas, 6 mt lentils, 5 mt dry broad beans, 4 mt dry pigeon peas, and 2 mt green broad and horse beans.¹⁶⁰

For most pulses and legumes, the majority (~80%) of environmental impacts occur during the production stage (figure 5.1). Transport generally accounts for a small proportion of impacts for all products. Two thirds of global soybean production occur almost equally in the US and Brazil, followed by Argentina which produces 11%.¹⁶¹ Hence, imports of certain products, such as soy beans, to the European market for example will tend to involve long distance transport (but not necessarily air freight). Drying and preserving pulses and legumes extends their availability throughout the year and reduces waste. Dried products also have a smaller energy requirement in relation to transportation, compared to frozen or canned products (due to a reduced weight and no refrigeration).¹⁶²

Figure 5.1: Proportion of environmental impacts from different life cycle stages: Pulses (legumes) (global)¹⁶³



In general, pulses and legumes are among the foods with the lowest relative environmental impacts – particularly peas and beans (table 5.1 - note that other pulses and peas are in dry weight).

Table 5.1: Average environmental impacts per kg of retail weight: Pulses (legumes) (global)¹⁶⁴

	Land use (m ² /kg)	Greenhouse gas emissions (kg CO ₂ eq/kg)	Acidifying emissions (g SO ₂ eq/kg)	Eutrophying emissions (g PO ₄ ³⁻ eq/kg)	Freshwater (L/kg)
Tofu (soybeans)	3.5	3.2	6.7	6.2	149
Other Pulses (dry)	15.6	1.8	22.1	17.1	436
Peas (dry)	7.5	1.0	8.5	7.5	397

Growing practices greatly influence the environmental impacts of pulse and legume production, in terms of both scale and type. Grown as part of crop rotations with cereals for example can provide numerous benefits including increasing the yield of subsequent cereal crops grown in the same area (as they use the nitrogen supplied by the pulses and legumes) and less requirement for plant protection products (for example, crop rotations reduce the risk of fungal disease such as pea root rot), as well as increasing landscape-scale heterogeneity and its associated biodiversity benefits. As legumes and pulses fix nitrogen in the soil, they do not require nitrogen fertilizers which is a major environmental benefit. There is less requirement for tilling of the soil for legume and pulse production.¹⁶⁵


Despite their nitrogen fixing properties, there are production practices that use high amounts of nitrogen fertilizer to increase yields – for example, soya bean cultivation in the USA.¹⁶⁶ Cultivating soya beans in monocultures requires the use of chemical plant protection products. In terms of pesticide residue, soybeans can have the same proportion of non-compliance with permissible limits as for fruit and vegetables.¹⁶⁷

Land use change associated with pulse and legume production can substantially add to their environmental impacts. For example, adding the CO₂ emissions from deforestation and burning of crop residues more than doubled the impact of soybeans grown in Brazil to 1.6 kg CO₂ equivalents per kg soya beans.¹⁶⁸ However, even with such additions, the climate impacts alone are relatively much smaller than most animal products such as beef, pork, chicken and cheese (see table 8.1). Land use change, such as deforestation, and the installation of monocultures with fertilizer and pesticide application can adversely impact the landscape and surrounding biodiversity. For example, heavy expansion of soybean cultivation on the Brazilian Cerrado is endangering important biodiversity.

Despite being considered the most species rich savannah in the world, the Cerrado is the least protected ecosystem in Brazil. This process of land clearing for soybean cultivation has also been introduced to the Amazon rainforest region (though 48% of soy production occurs in the Cerrado region, and 13% from the Amazon).¹⁶⁹ One consequence of such large-scale soybean cultivation is the infrastructure created to enable harvests to be transported, which in turn opens up the region for further exploitation. Soybean cultivation in the Amazon region has largely resulted in the conversion of existing small-scale agriculture into large-scale farming with high use of machinery (and thus much less requirement for farmers), a dominance of monocultures, a lack of crop rotation, and ongoing soil erosion.¹⁷⁰

While soy production is a substantial driver of deforestation in the Brazilian Amazon, the expansion of pasture land for beef production is the leading driver. In addition, 95% of Brazilian soy is used for animal feed.¹⁷¹ Globally the picture is similar - soy production is estimated to be the 3rd largest driver of deforestation (with cattle and palm oil being the 1st and 2nd largest drivers, respectively).¹⁷² Only 7% of global soy production is used to produce tofu, soy milk, and tempeh directly for human consumption, and around 13% is used to produce oil for human consumption. Most soy (76%) is used to feed farmed animals – largely chickens and pigs (figure 5.2).¹⁷³ However, an analysis of soy embedded in food consumption across the EU found that farmed fish had the largest amounts per unit of product, followed by chicken meat.¹⁷⁴ Globally, a small portion of soy production is certified as deforestation-free.

Figure 5.2: Allocation of global soy production to its end uses by weight (2017-2019).¹⁷⁵




Around a quarter (23%) of pulses produced in 2019 were used for farmed animal feed.¹⁷⁶ Reducing the consumption of animal products, particularly chicken and pork, is therefore the most effective way to reduce the environmental impacts of soy production, and also, but to a lesser extent and

concern, pulse production. Replacing portions of beef consumption with pulses can also reduce environmental impacts considerably – for example, by reducing GHGs to make a substantial contribution to national climate goals, while also sparing large areas of agricultural land.¹⁷⁷ Due to the inefficient conversion of plant nutrients to animal nutrients, the environmental impacts of humans consuming pulses directly are substantially lower.¹⁷⁸

Nordic and Baltic context

Pulses (and legumes) account for low environmental impacts within the Nordic and Baltic diets. Due to the nitrogen fixing process during production, pulses account for little nitrogen use in relation to consumption across the Nordic region. Cropland and freshwater use tend to account for the majority of impacts, particularly for consumption in Norway (figure 5.3).

Figure 5.3: Impacts of pulse (legume) consumption across the Nordic and Baltic countries in relation to the food portion of global environmental limits (%)



Source: *Global Nutrition Report 2021*¹⁷⁹ The analysis utilizes country-specific food consumption and environmental footprint data and relates them to the food portions of global environmental limits i.e. a global test to assess the impacts if everyone in the world consumed at the given rate. The methods and data are described in Box 1, and presented in appendices 1-3.

Pulses and legumes are an under-consumed food group across the region. The most promising process for increasing consumption is in conjunction with a reduction in meat consumption i.e., a replacement of portions of meat with legumes/pulses, which would maximise environmentally beneficial outcomes and potentially health benefits. Such a shift is considered a strong motivator for exploring options to increase local production. In addition, there are issues related to imports such as deforestation from soy. For example, soy imports to Denmark have been estimated to use a land area equivalent to the combined land area of Zealand and Lolland, and are considered to pose the greatest risk (in comparison to timber, pulp and paper), as 65% of the land used is located in countries that are high or very high risk for deforestation and social challenges.¹⁸⁰ In general, there is potential to increase pulse/legume production across the region. For some countries, current plans revolve around increasing pulse production as part of a crop rotation for the purpose of farmed animal feed (e.g., Norway). There are limits regarding the extent that pulses and legumes can be grown for direct human consumption in terms of variety, as lentils and chickpeas might be limited to

certain areas and even the production of broad beans is limited in Finland. There are ongoing chickpea trials in Denmark for example, with yields being weather dependent.¹⁸¹ There is also potential for learning from other regions with similarities in weather and climatic conditions that have recently started to produce chickpeas on a commercial scale (e.g. the UK¹⁸²).

Some countries are more favourably positioned geographically for pulse production (e.g., greater potential in southern Sweden compared to Norway or Finland). Generally, peas and various types of beans grow well in the Nordic region (peas were historically grown on a larger scale for direct human consumption before shifting agricultural production to favour animal agriculture). Hence, there is a legacy of pulse and legume production in the region, which can be utilised to guide the expansion of current production. It is important to increase research in support of increasing pulse and legume production, including breeding and field trials (which might also involve increasing the grade of crops from animal feed to a grade high enough to enable direct human consumption) - and also providing appropriate training for practitioners to overcome a lack of experience in cultivating legumes, including the use of technologies to increase protein content and crop rotations techniques.¹⁸³ In Sweden, production is currently limited to one area and certain types of legumes/pulses (fava/broad beans and yellow peas), although it is possible to grow hardy varieties of soy beans in southern Sweden.¹⁸⁴ However changes in climatic conditions could also change the opportunities for pulse and legume production (for example, in Sweden and southern Finland). The use of greenhouses and/or tunnels could increase the production of soy and broad beans. Therefore, the growth of pulses and legumes in the region could be considered dynamic rather than being limited to current opportunities. There is also an issue of scale. For example, in Sweden production tends to be niche and limited to local gardens, and hence would require scaling up (this is also an aspect where learning from other countries such as the UK could be particularly useful for informing how to scale from niche to commercial level). Due to the relatively small scale of production, pulses are generally lacking compared to other crops in terms of investments to increase yields, and research on pests and diseases.

An increasingly prominent area of research regarding pulse production and consumption across the region relates to their use in various meat-replacement products, or meat alternatives. For example, in Sweden the broad beans and yellow peas are currently used to produce tempeh. Increasing plant-protein production for such products is a growing field of emerging research in Finland. For example, in Finland there are large research projects such as ScenoProt¹⁸⁵ and 'Leg4Life'¹⁸⁶ which aim to develop, with stakeholders, cultivation methods to increase production of legumes that flourish in Finnish conditions (pea, faba bean, lupins, clovers) and processing methods to create legume products for animal feed and for direct human consumption. Such an initiative could help overcome a reliance on soy beans in processed products, which is apparent due to a lack of processing opportunities for domestically produced legumes.¹⁸⁷ Research in this field has also increased in Denmark where old pea varieties grown for human consumption are being investigated in the project Peas & Love (covering pea yield stability, taste, and quality), and organic broad beans cultivation in the project ØkoFaba. In Norway, projects exploring the opportunities for growing legumes in a warming climate have yielded promising results (e.g., FoodProFuture and VOM).

For some countries (such as Iceland), a dependency on imports is likely to remain due to the limited potential for cultivating legumes and pulses, and for others (such as Norway) a reconfiguration of existing cropland, in addition to fixing the current yield gap,¹⁸⁸ would likely be required due to a limited availability of cropland. For example, replacing a proportion of crops currently cultivated for animal feed with pulses or legumes for direct human consumption.¹⁸⁹ If implemented as crop rotations with cereals, such a reconfiguration would help to tackle a number of fungal diseases that

impact cereals. It is possible that some domestically grown pulses and legumes could at least partly replace imports, for example brown beans are a good alternative to kidney beans imported from the US.¹⁹⁰ In some regions and for some crops, there is also an element of cultural heritage related to legume production (e.g., brown beans on Öland, where farmers can claim environmental subsidies for conserving local varieties and farming methods¹⁹¹). Legume production can have positive impacts on the landscape and biodiversity and contribute to a varied agricultural landscape.¹⁹² The specific opportunities to cultivate pulses and legumes within the region might be influenced to some extent by permissible plant protection products, as this is not the same in every country.

Currently, all countries produce pulses – with Latvia being the highest producer and Lithuania the lowest producer. Finland, Lithuania and Norway depend to an extent on imports to meet supply (table 5.2).

For comparison, pulse/legume consumption in line with a flexitarian diet that aligns with planetary boundaries (if implemented in combination with medium-ambition technological measures to reduce inputs during food production, and reductions in food loss and waste of 75%), recommends a minimum daily pulse/legume intake of 75g, or 27.4 kg per year (as part of an average daily intake of 2,100 kcal).¹⁹³ Based on the annual supply data in table 5.2 and consumption data in table 2, all countries consume far below this recommended minimum intake (on a per person basis).

Table 5.2: Pulse¹⁹⁴ production and supply across the Nordic and Baltic countries in 2019¹⁹⁵

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	85	77	45	117	1
Estonia	111	4	67	48	5.7
Finland	64	28	0	92	2.6
Iceland	(no data)	0	(no data)		1.1
Latvia	102	59	132	29	0.4
Lithuania	306	19	270	55	2.6
Norway	30	584	2	612	12.6
Sweden	131	49	22	158	2.5

Peas are generally the most highly consumed across all countries, except for Iceland where beans are the most highly consumed (table 5.3).

Table 5.3: Pulse (legume) food supply across the Nordic and Baltic countries in 2019 (kg/person/year)¹⁹⁶ by pulse type

	Beans	Peas	Pulses, other	Soybeans
Denmark	0.1	0.5	0.3	0.2
Estonia	0.05	5.5	0.01	0.2
Finland	0.03	2.3	0.08	0.2
Iceland	0.5	0.02	0.2	0.4
Latvia	-	-	0.2	0.2
Lithuania	0	1.6	0.9	0.1
Norway	0.1	7.4	4.8	0.3
Sweden	0.2	1.7	0.3	0.3

Considerations for reducing environmental impacts of legume and pulse consumption in Nordic and Baltic countries

From the overview of legume and pulse production, consumption, and trade provided in this section, we identify a number of opportunities to explore as potential ways to reduce environmental impacts of the legume and pulse food group:

- Pulses and legumes have among the lowest relative impacts, from an environmental sustainability perspective, and have much lower impacts across the board in comparison to meat for example, whether the pulses/legumes are domestically produced or imported. Reducing the consumption of animal products and replacing portions of them with legumes and pulses is the major route to reducing overall environmental impacts while simultaneously increasing pulse/legume consumption. Reducing the absolute amount of beef, pork and chicken production in particular could reduce the substantial environmental impacts (including deforestation) of soy bean production given that most (76%) soy beans are used for animal feed.
- Reducing imports of soy beans for farmed animal feed is considered in section 8 on meats and meat products.
- Crop rotations with legumes/pulses and cereals would reduce the need for plant protection products such as fungicides, and nitrogen fertilizers. Increasing production of legumes/pulses for direct human consumption within the Nordic region would also help to diversify agricultural production and landscapes, and reduce reliance on imports.
- Reducing the overall land requirements of agriculture at the global level (by reducing livestock and feed crop production, and increasing legume/pulse consumption) could have additional major benefits for biodiversity and climate change targets if native ecosystems and vegetation cover were allowed to recolonise spared land. This shift would also allow space for more environmentally friendly farming methods that typically have lower yields compared to conventional production methods (e.g., organic).¹⁹⁷
- Traceability within the supply chain of imported pulses/legumes is not always possible. Hence, the environmental impacts of imported products are not always known. A precautionary approach would be to adopt a default organic (or similar) procurement policy, and adopt approaches to enable and incentivise organic (or similar) methods with crop rotations for production within the Nordic region.
- Innovation in legume/pulse production, manufacture and processing, and consumer behaviour could facilitate an increased production and consumption of legumes/pulses for direct human consumption.

NNR Food Group 7: Fish, fish products and seafood

Global context

Globally, around 200 million tonnes (mt) of fish and seafood are produced every year (176,592,630 tonnes in 2019¹⁹⁸). In recent decades, growth in aquaculture has supplied most demand growth while growth from wild caught fish has been limited due to protective measures taken in light of the well-established scientific consensus on the state of the world's wild fish stocks. Farmed fish and seafood now contribute 53% to total global production of fish and seafood and is expected to continuously increase due to limited growth potential in the capture sector.¹⁹⁹ The majority (84%) of wild-caught fish are consumed directly by humans, and 16% are used to feed farmed animals (including 11% for farmed fish). China produces the largest amounts of fish and seafood (60 mt in 2017), followed by Indonesia, India, Vietnam and the US.²⁰⁰


Capture fisheries and aquaculture both have a range of environmental impacts, which vary depending on the habitat types, production method and equipment type - ranging for example from intensive and higher-input shrimp and salmon farming to lower-input mussel farming in aquaculture; and from lower bycatch and lower fossil fuel use in pole and line fisheries, to benthic trawl fisheries with significant bycatch, damage to habitats and fossil fuel use. There are also dependencies between the two – for example, capture fisheries providing feed inputs to aquaculture; and various forms of pollution from aquaculture impacting capture fisheries.²⁰¹ However, some impacts of capture fisheries are particularly difficult to measure – for example, in terms of catch, only the fish brought back to land ('landings') – are recorded in most databases, such as those of the UN fisheries. Discards are not reported, making it impossible (or very difficult and with uncertainties) to assess fish stocks.²⁰² Despite producing only 1% (by weight) of food and feed for farmed animals globally, aquatic systems (wild and farmed) have been estimated to account for 10% of the footprint of all food produced in terms of greenhouse gas emissions, freshwater use, habitat disturbance, and nutrient pollution.²⁰³

In terms of GHGs, the main impact from capture fisheries is fossil fuel use for fishing vessels (although emissions from shipping of exports can be higher e.g., seafood from Norway to Asia²⁰⁴), while the main GHG impact for aquaculture comes from feed production (although air freighted exports can have a higher impact e.g., farmed salmon from Norway to Asia²⁰⁵). The impacts of climate change are expected to affect capture fisheries and aquaculture. For example, high-latitude regions will experience an average increase of 30 to 70% in terms of overall catch potential, while a decreased catch of up to 40% is likely to be experienced in tropical areas.²⁰⁶ Warming seas due to global temperature rise could reduce the nutritional content of fish – for example, plankton living in cold (-2°C) regions contain 3 times more unsaturated fatty acids than those in warm (29°C) waters.²⁰⁷ In terms of aquaculture, even small temperatures can impact productivity of many farmed species, in addition to creating new opportunities for diseases and parasites. Increased frequency and intensity of adverse weather could impact farm infrastructure, and land-based operations could experience freshwater shortages. Supply of crop-based feeds could also be threatened.²⁰⁸ Climate change could therefore negatively impact food webs and fisheries.²⁰⁹

Figure 7.1 provides a range of GHG footprints associated with fish and seafood production, based on data from 1690 fish farms and 1000 unique fishery records. The 23 represented species groups cover over 70% of global 'blue food' production.²¹⁰ Generally, there is no clear size difference between GHG footprint in terms of capture vs farmed – both production types have a range of footprint sizes depending on species (although GHGs from some methods have been difficult to quantify, e.g.,

bottom trawling, and require further investigation²¹¹). Small pelagic fishes (herring, sardines) generate lower emissions than all fed aquaculture, and flatfish and crustaceans generate the highest amounts. When comparing the same species, farmed fish tend to have lower GHG footprints compared to wild caught (e.g., salmon and shrimp) – with the difference being most substantial for bivalves. However, there are examples where wild caught salmon have lower GHG footprints compared to farmed salmon. For example, a recent assessment of Norwegian salmon production found GHGs from wild caught to be up to 86% lower than farmed – which also had higher land use requirements and marine ecotoxic and eutrophying emissions.²¹² Seaweed has the smallest GHG emissions across the range of seafood (figure 7.1).


Figure 7.1: Greenhouse gas emissions per kilogram of edible fish and seafood



Source: Gephart et al. (2021). Environmental performance of blue foods. Nature. OurWorldInData.org/fish • CC BY
 Note: Includes on-farm and off-farm, but does not include post-farmgate emissions. This means it does not include impacts such as transport to retail, packaging, processing or cooking.

The land requirements for aquaculture result mostly from feed production, and less so from the conversion of terrestrial areas to fish farms, which account for 60% of global aquaculture production (10% of production is from coastal ponds and the remainder is from open-water cages and ropes, in seas and in lakes). Around 70% of farmed fish are given supplemental feed inputs derived from agriculture. Farming of unfed fish species like filter feeding molluscs accounts for around 24% of aquaculture production and put less pressure on land, however this type of production is growing at a slower rate compared to fed species.²¹³ Figure 7.2 shows a range of land use footprints from aquaculture, including land used for farms and feed production. The variation in land use between species can be substantial (e.g., milkfish vs trout). Bivalves and seaweed have the smallest land use requirements. One key consideration regarding land use is the opportunity cost i.e. other competing land uses, or uses that provide greater levels of social and environmental benefits in comparison. Growing crops for fish feed rather than crops for direct human consumption is a growing tension.²¹⁴

Figure 7.2: Land use per kilogram of edible fish and seafood ²¹⁵




Source: Gephart et al. (2021). Environmental performance of blue foods. Nature.

OurWorldInData.org/fish • CC BY

Freshwater use in capture fisheries relates almost entirely to post harvest activities. Direct freshwater use in aquaculture can be large for some pond-based systems but can vary significantly depending on system characteristics, location, and targeted species. Some forms of aquaculture ‘recycle’ water by using it temporarily on the farm and releasing back in a more polluted form (e.g., added nutrients and chemicals), which might subsequently reduce its useability and/or effect nearby ecosystems. Water use for feed crops can also be significant.²¹⁶ Figure 7.3 shows water use footprints for a range of farmed species. As for GHGs and land use, water usage can vary substantially between species, with trout being the fish with the smallest footprint. Bivalves and seaweed do not require freshwater for production, making them particularly favourable at least from a water scarcity perspective.

Figure 7.3: Freshwater use per kilogram of edible farmed fish and seafood ²¹⁷




Source: Gephart et al. (2021). Environmental performance of blue foods. Nature.

OurWorldInData.org/fish • CC BY

Nitrogen (N) and phosphorus (P) emissions from direct nutrient leakage from aquaculture farms, and also from the agricultural production of feed contribute to marine and freshwater eutrophication. Figures 7.4 and 7.5 show nitrogen and phosphorus footprints of a range of species produced from aquaculture. In fed systems, most N (>87%) and P (>94%) emissions occur on-farm. Silver/bighead fish have the lowest N and P footprints, while seaweeds and bivalves result in negative emissions due to being produced in extractive systems that remove more N and P than is emitted. Changing the feed type can change N and P impacts – for example, replacing feed for rainbow trout with Baltic herring could reduce the total eutrophication impact of rainbow trout. The impacts could be further reduced if the wild caught herring replaced the farmed trout.²¹⁸


Figure 7.4: Nitrogen emissions per kilogram of edible farmed fish and seafood ²¹⁹



Source: Gephart et al. (2021). Environmental performance of blue foods. Nature.

OurWorldInData.org/fish • CC BY

Figure 7.5: Phosphorous emissions per kilogram of edible farmed fish and seafood ²²⁰



Source: Gephart et al. (2021). Environmental performance of blue foods. Nature.

OurWorldInData.org/fish • CC BY

Both capture fisheries and aquaculture have a range of adverse impacts on biodiversity. In relation to capture fisheries, over-harvesting, bycatch and destruction of habitats through equipment are the major routes. Land and aquatic space conversion for farms, effluents (such as nitrogen and phosphorus emissions) to the local environment, and direct killing of wildlife (e.g., if they are a predator of fish) are the major routes for aquaculture. Genetic introgression from escaped species has also been reported.²²¹ For some species (e.g., shrimp), aquaculture requires the capture of

larvae from the wild, which also has bycatch impacts. Biodiversity impacts related to feed production (including habitat loss and degradation, use of plant protection chemicals, and nutrient pollution) also apply to fed aquaculture.

Currently, 92% of the total ocean is unprotected (i.e. is not governed by a global treaty), and 82% of national waters are also unprotected.²²² Globally, 34% of fish stocks are overfished, 60% are maximally fished and 6% are underfished (according to the most recent estimate for 2017).²²³ The Mediterranean and Black Sea have the highest proportion (63%) of stocks fished at unsustainable levels, followed by the Southeast Pacific (54%), and Southwest Atlantic (53%).²²⁴ A number of species are estimated to be in a particularly dangerous state. For example, mackerels are estimated to be below optimal levels due to an increase in fishing intensity. Sharks are rapidly declining – and 37% of all sharks and rays are threatened with extinction. Southern Bluefin tuna need more time to recover to optimal levels. However, there are a number of issues that make such estimates of fish stocks unreliable. First, only the catch brought to land is counted in most databases (bycatch/discards is not counted), and data is much more sparse regarding the status of fish stocks across Asia, Latin America and much of Africa, in comparison with Europe and North America. Assessing the status of fish stocks requires data on catch and how quickly fish populations recover. While there are measures in place to overcome some of the data issues (including an assessment every decade by the UN Fisheries Division), such data gaps present a major hinderance to asserting what is sustainable or not.²²⁵

It has been estimated that most discards (93%) result from large, industrial scale fisheries with the largest discards associated with bottom trawling - on average, 21% of catch from bottom trawling is discarded, increasing to over 50% for shrimp trawling. In terms of global fish discards, bottom trawling is estimated to account for around 50% of the total fish discards from all methods.²²⁶ In addition to high discard rates, trawling physically damages ecosystems. The extent of the damage largely depends on the depth of the trawl into the sediment (although sediment type and lifeforms in the area are also important). On average, 6% (from Otter trawl) to 41% (from hydraulic dredging) of faunal biomass per pass are removed, with recovery of the damage taking 1.9–6.4 years post-trawling, depending on fisheries and environmental context.²²⁷ Trawling, for example, in the Barents Sea affects the biomass of all species but especially those easily caught by a trawl.²²⁸ Around a quarter of global annual fish catch is from bottom trawling, and it is the dominant method used in China and India. Purse seine (vertical nets) accounts for 20% of global annual catch and is also more commonly used in industrial fishing practices as, also for trawling, these methods tend to result in a higher yield. Purse seine is, in some cases, also associated with high levels of bycatch. The lowest levels of bycatch are associated with pole-and-line, and longline methods which are more commonly used in lower income countries for subsistence or small-scale capture.²²⁹

There is growing evidence that seabed habitats throughout the world's oceans are being impacted by physical destruction or selective removal of habitat-forming species. Removal of the latter, or drastic lowering of the population, can impact entire food webs and ecosystem functioning (particularly if apex predators are removed/reduced, and predator-prey interactions are impacted).²³⁰ One consequence is a reduction in vegetation where predator populations are too low and allow herbivorous species to flourish – in turn reducing the carbon storage capacity of marine environments.²³¹ In terms of habitat disturbance, demersal fish (also known as groundfish and include cod and haddock), are estimated to have by far the largest impact from capture fisheries, and shrimp have the largest impact from aquaculture. However, disturbance from capture fisheries tends to be much greater in comparison to that from aquaculture even when accounting for disturbance from feed production.²³²

Aquaculture can also impact entire ecosystems, for example, freshwater and brackish-water pond farming has driven large-scale local and regional landscape transformations including large areas (~60,000 km²) of ecologically valuable coastal agricultural land and wetland habitats (mainly along South China, India, Vietnam, Indonesia and Bangladesh), that have been fragmented through land reclamation and conversion. Mass escapes from fish farms resulting in cross breeding and behaviour change in wild salmon is an ongoing issue, as is pathogen spread to wild species, in addition to nutrient waste polluting local environments.²³³

Ocean acidification (OA) resulting from increasing atmospheric concentrations of CO₂ has been identified as a substantial threat to marine ecosystems, and both capture fisheries and aquaculture – and wider food systems - contribute through fossil fuel use (including direct and indirect use). Capture fisheries are expected to be impacted by OA, as the distribution and catchability of over a hundred fish species could decline by up to 30%.²³⁴ Warm water coral reefs and calciferous marine organisms are among the marine biodiversity impacted by OA.


The fishing industry contributes substantially to ocean plastic pollution – for example, over 46% of the mass comprising the great pacific garbage patch is fishing nets.²³⁵ Such waste from the fishing industry impacts biodiversity in a number of ways, including ingestion of plastic and entanglement in discarded nets and equipment.²³⁶

As with other forms of animal farming, animal welfare and antibiotic use are important issues to consider. Antimicrobials are commonly used in aquaculture, contributing to AMR. While these topics are outside of the scope of the current paper, they are important issues to consider in any assessments of sustainable food systems and diets.

Nordic and Baltic context

Figure 7.6 shows the variation in environmental impacts of fish consumption across the Nordic and Baltic countries and how they relate to global limits for food consumption. Nitrogen, phosphorus and cropland tend to account for the majority of impacts.

Figure 7.6: Impacts of fish consumption across the Nordic and Baltic countries in relation to the food portion of global environmental limits (%)



Source: *Global Nutrition Report 2021*²³⁷ The analysis utilizes country-specific food consumption and environmental footprint data and relates them to the food portions of global environmental limits i.e. a global test to assess the impacts if everyone in the world consumed at the given rate. The methods and data are described in Box 1, and presented in appendices 1-3.

A comprehensive assessment of environmental impacts of fish capture is difficult – particularly in relation to the biodiversity impacts. The lack of data availability regarding fish stocks makes an assessment of the environmental sustainability of capture rates impossible (due to the lack of frequent and comprehensive data collection on discards). Similarly, assessing the biodiversity impacts is impossible without reliable information on bycatch, or data on physical damage to ecosystems via trawling methods. The environmental impacts of aquaculture are generally less challenging to assess, with the exception of biodiversity. A lack of data availability needed to assess environmental sustainability in a comprehensive way, combined with an expected increase in global fish production (mostly via aquaculture) and the impacts of environmental shocks and climate change make a precautionary approach to recommendations for fish consumption important i.e., more information is required before any targeted increase in fish consumption in the Nordic and Baltic countries could be justified from an environmental sustainability perspective, and also from a food security/risk perspective. This might include identifying under-utilized freshwater fish stocks, shifting supply from animal feed to human consumption, and assessing the balance of impacts in different environments including the potential to reduce eutrophication in fresh water bodies. Other potential options to explore (particularly to reduce pressure on capture fisheries) are mussel, algae, and seaweed. For example, mussel farming already takes place along the west coast of Sweden and could provide an important case study in terms of environmental impacts and scalability.

Increasing aquaculture production could be an issue in terms of feed production from agriculture and the associated impacts (for example, see chapter 5 for more information on the environmental issues related to soy production). An increased demand for feed necessarily raises the issue of feed vs food in relation to food security, and competition with other land uses such as biofuel production and carbon capture (this issue applies to all animal farming to varying extents, see for example chapter 8 for more information). Due to their inefficient conversion of shorter-chain fatty acid, α -linolenic acid (ALA; 18:3n-3), into EPA and DHA, salmon along with other cold water marine species of fish must obtain n-3 LC-PUFA through their diet (fish oil and fish meal in aquaculture). However, changes to feed composition (including an increased use of oil seeds), in response to rising demand for fish oil and fish meal, have resulted in a reduced nutritional benefit to humans consuming the farmed fish.²³⁸ Hence, there is scope to explore direct sources of n-3 fatty acids for human consumption as an alternative to consuming fish from aquaculture, such as microalgae already in use as a n-3 fatty acid supplement.

Increasing production from capture fisheries is limited in terms of location across the Nordic and Baltic region due to high levels of pollution in the Baltic Sea and also lakes in Sweden for example, which could make it problematic to recommend local production. Hence, for some countries such as Sweden, increasing fish production would likely be from aquaculture. Otherwise an increase in consumption might be met through imports from Norway, Denmark or Iceland for example, which are all large producers and exporters (table 7.1), or through increasing production in lakes and capturing wild freshwater fish throughout Finland. In these scenarios, it would still be necessary to reduce the overall environmental burden of fishing and fish consumption to prevent problem-shifting e.g., reducing one environmental issue in one location while increasing them in another location, and also to contribute to a range of environmental targets. One aspect to consider in relation to the environmental impacts of importing (and exporting) fish is the use of air cargo. A full assessment of the environmental impacts of capture fisheries (including production methods) and aquaculture (including nutrient emissions and the feasibility of implementing technologies to reduce them, and antibiotic use) across the region would be required in order to identify optimal pathways.²³⁹

Norway is the largest producer, importer and exporter across the Nordic and Baltic countries (table 7.1), and the world's second largest seafood exporter.²⁴⁰ In Norway, a discard ban, which comprises

of several measures, has possibly significantly decreased discards²⁴¹, although monitoring is challenging.²⁴² Iceland and the EU also have discard bans (the ban in the EU is less restrictive compared to the ban in Norway). The discard ban in the EU was launched in 2017 and its impacts have not yet been estimated.²⁴³ Fisheries management is international in scope, meaning that making changes to fisheries in Nordic/Baltic seas (e.g., to harvest rate, selection pattern, discard bans) needs to be decided with other countries with which the stocks are jointly managed through international agreements (e.g., the EU, UK, and Russia). For example, about 90% of Norwegian fisheries, both in value and volume, are taken from stocks it shares with other countries.²⁴⁴ However, given that Norway is the tenth largest producer²⁴⁵ and second largest exporter of fish and seafood globally, it could have more potential to influence such decision making compared to the other Nordic and Baltic countries.

Iceland currently has the highest levels of fish and seafood supply across the region - almost double the amount consumed in Norway, which has the second highest supply levels (per person). Estonia has the lowest supply levels, around 6 times lower than Iceland (table 7.1). For comparison, fish and seafood consumption in line with a flexitarian diet that aligns with planetary boundaries (if implemented in combination with medium-ambition technological measures to reduce inputs during food production, and reductions in food loss and waste of 75%), recommends a minimum daily fish and seafood intake of 28g, or 10.2 kg per year (as part of an average daily intake of 2,100 kcal).²⁴⁶ Based on the annual supply data in table 7.1, all countries exceed this amount (on a per person basis) 1.5 - 9 fold. In contrast, using fish consumption data from table 2 indicates that consumption in Denmark, Iceland, Sweden and Norway exceeds the minimum amount consistent with the flexitarian diet – all other countries consume close to or slightly below this amount.

Table 7.1: Fish (seafood) production and supply across the Nordic and Baltic countries in 2019²⁴⁷


Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	942	1,214	1,955	201	27
Estonia	84	50	112	22	15
Finland	205	134	73	266	34
Iceland	1,184	65	1,146	103	91
Latvia	119	88	134	73	25
Lithuania	94	196	197	93	33
Norway	3,677	1,222	3,105	1,794	51
Sweden	247	1,075	962	360	32

One assessment of fish production suggested at the global level a maximum amount of 3.5 kg per person per year would accord with a sustainable load on fish stocks (maximum sustainable yield of ~66 mt from capture fisheries in 2030).²⁴⁸ Assuming around 50% (in line with the global production split between capture and aquaculture) of consumption within the Nordic and Baltic countries is sourced from capture fisheries, an annual consumption of 3.5 kg is around half of the amount consumed in the lowest consuming nation (Estonia) and 13 times lower than the highest consuming nation (Iceland) within the region. However, this estimated sustainable consumption could be questioned because of large uncertainties due to lack of available data, as described above - a

comprehensive assessment of sustainable fish and seafood yields in the Nordic and Baltic countries is needed.²⁴⁹


For all countries, capture fisheries represent by far the largest proportion of fish production – the biggest contribution from aquaculture is in Norway, where it accounts for around a third of total fish production (figures 7.7 and 7.9). Production levels from capture fisheries have generally declined in Norway, Iceland, Denmark and Sweden since 1997 (figure 7.7).

Figure 7.7: Capture (wild) fishery production (tonnes) in the Nordic and Baltic countries: 1960 - 2018



Capture from bottom trawling has also generally declined from 1997 levels, except for Norway and Iceland, where bottom trawling has increased (figure 7.8). Bottom trawling is used for around a third of fish capture in Norway, and around 21% of fish capture in Iceland.

Figure 7.8: Wild fish catch (tonnes) from bottom trawling in the Nordic and Baltic countries: 1950 – 2018



Source: FishStat via Pauly, Zeller, and Palomares from Sea Around Us Concepts, Design and Data.


OurWorldInData.org/fish • CC BY

Norway has by far the largest production from aquaculture, which increased sharply from 1985 to a current production of around 1.3 mt per year (figure 7.9a). Across all countries in the region, the main production is fed finfish, dominated by salmonoids (salmon, trout and char). The variation in environmental conditions between the countries makes some of them more favourably positioned for aquaculture and impacts sustainability considerations. For example, Iceland, Norway, and Denmark benefit from access to long coastlines with strong currents, whereas Denmark also has coastline towards the Baltic sea, as does Finland and Sweden, which is highly polluted, populated and has low oxygen levels, limiting the possibilities for large scale open-water aquaculture.²⁵⁰


The environmental impacts of most concern in relation to aquaculture production in Norway and Iceland relate to sea lice, escapes and disease and gene transfer to wild fish stocks, organic wastes, and land-use change concerns related to soy production (for feed). Nitrogen and phosphorus emissions, and organic wastes are the main concerns in Denmark, Sweden and Finland. Within the region, the high reliance on inputs from agriculture (of which the high-grade ingredients could be used directly for human consumption), are common concerns. In all settings, antibiotic treatments are of concern.²⁵¹

Nordic aquaculture policies currently focus on growth, both in terms of production volumes and economic gains (through high-value, fed-fish including salmon or trout), with strategies to intensify production through technology rather than shift to less environmentally burdensome species and production levels, and to replace soy with alternative proteins as feed. Hence, the priority appears to be an increased production with a reduced environmental impact per unit of production i.e. reducing relative rather than absolute impacts.²⁵²

Figure 7.9: a) Aquaculture production (tonnes) in the Nordic and Baltic countries: 1960 – 2018; b) Aquaculture production (tonnes) in the Nordic and Baltic countries: 1960 – 2018 (excluding Norway to enhance visibility of production amounts from the remaining countries).



Source: Food and Agriculture Organization of the United Nations (via World Bank) OurWorldInData.org/fish-and-overfishing • CC BY



Source: Food and Agriculture Organization of the United Nations (via World Bank) OurWorldInData.org/fish-and-overfishing • CC BY

Combining supply from all production methods, pelagic fish dominate consumption in Denmark, Iceland, and Lithuania. In Estonia and Norway demersal fish dominate; in Finland freshwater fish dominate; in Sweden the consumption of demersal and freshwater fish is equal, and higher than pelagic; and in Latvia 'other marine' fish dominate (table 7.2). Iceland has the highest consumption of crustaceans, followed by Norway. This group includes lobsters and shrimps, which tend to have

among the highest environmental footprints (especially from capture fisheries) (figures 7.2 – 7.4). Demersal fish tend to be mostly carnivorous and hence have different feed requirements and impacts within an ecosystem compared to herbivorous and planktivorous fish. Bluefin tuna (pelagic fish) are apex predators, which have a particularly important role in maintaining ecosystem functioning. Mackerels are also pelagic fish and are estimated to be below optimal levels due to an increase in fishing intensity. The sum of the unilateral quotas for mackerel and the resulting catches in the Northeast Atlantic and adjacent waters have exceeded the scientific advice by 41% on average since 2010.²⁵³ However, it is generally unknown how much mackerel is consumed and where it is sourced from, in relation to consumption in the Nordic and Baltic countries. There is also lack of consensus on how the quota should be shared between the countries fishing this mackerel.²⁵⁴ Molluscs (a bivalve) tend to have among the lowest environmental impacts across the range of metrics (GHGs, land, water, nitrogen, phosphorus and biodiversity) yet are consumed in lower quantities in comparison to other types of fish and seafood (except for cephalopods and other marine fish, which are consumed in lower amounts in most countries) (table 7.2). Consumption data is not available for seaweed, which is the lowest impact seafood.

Table 7.2: Fish (seafood)²⁵⁵ supply across the Nordic and Baltic countries in 2019 (kg/person/year)²⁵⁶ by fish type

	Pelagic	Demersal	Freshwater	Crustaceans	Molluscs	Cephalopods	Marine fish, other
Denmark	10	6	2	7	1.4	0.1	0.0
Estonia	2	7	1	3	0.6	0.2	0.0
Finland	10	3	18	2	0.2	0.1	1.2
Iceland	44	21	10	16	0.2	0.0	0.0
Latvia	8	4	2	1	0.5	0.2	10.0
Lithuania	28	1	2	2	0.4	0.3	0.0
Norway	6	23	11	10	0.8	0.1	0.1
Sweden	5	9	9	7	0.9	0.1	0.7

Considerations for reducing environmental impacts of fish and seafood consumption in Nordic and Baltic countries

Reducing fish consumption among high consumers in higher income countries with a good food supply will likely be important for reducing the overall environmental burden of the food system. In addition to the current issues regarding fish and seafood production, aquaculture is one of the fastest growing forms of food production globally and will continue to increase pressure on environmental resources and contribute to environmental problems. Challenges remain even for the most technologically advanced aquaculture systems. There is no single measure or innovation that will resolve all environmental challenges related to fish and seafood production (both capture and aquaculture), instead a range of measures are required.²⁵⁷ We offer a range of considerations:

Major:

- Due to the potentially large-scale impacts on ecosystems and the largely unknown nature of fish stocks globally, a precautionary approach to this food group should be taken.

- A comprehensive assessment is needed to identify sustainable yields from marine and lake ecosystems within the Nordic and Baltic countries²⁵⁸, including capture fisheries and aquaculture. It could also identify whether there are any sources of fish that on balance have positive environmental impacts. This should factor in potential impacts of climate change, environmental shocks, antibiotic use, animal welfare, waste, disease transfer and ecosystem-level impacts on biodiversity in addition to potential benefits such as removing nutrients from polluted waters. Due to changes in environmental conditions, regular monitoring, modelling and recalculations will likely be needed.
- Underpinned by a precautionary approach, the assessment should also carefully consider if there is a need to increase consumption of fish and seafood, whether the environmental impacts of increasing consumption are adequately known, and the feasibility of shifting as much continued production and consumption as possible to the lowest impact foods (for example, edible seaweeds which are particularly underrepresented in the literature²⁵⁹).
- The environmental impacts from all types of production should be minimized (this mostly applies to animal sourced capture and aquaculture). One important aspect for aquaculture is improving feed conversion ratios across all fed groups, which could reduce land and water use by up to 50%. Optimising capture equipment could reduce GHG emissions by more than 50% for some groups.²⁶⁰ However, it should be noted that minimising environmental impacts does not necessarily equate with environmental sustainability – an assessment will still be required to ensure production levels are consistent with environmental goals. In addition, impacts will increase if relative impacts are reduced but absolute impacts increase (due to an increase in production).
- Feed to food tensions including food security and efficiency aspects should also be considered, including the potential to reallocate resources used to produce feed to human edible foods.

Minor:

- Environmental impacts that are not included in the data presented in this chapter, for example, the energy used for drying seaweed and CO₂ emissions during shell formation of bivalves, and energy used in the transportation of live bivalves, should be factored in.²⁶¹
- For any production of animals from aquaculture, the focus should be on bivalves (considering the previous point), which are a non-fed species with the lowest environmental impacts across a range of metrics. Currently 30% of aquaculture production is “non-fed” species (mostly carps and bivalves).²⁶²
- Environmental impacts of increasing imports and exports (such as those from air cargo), should also be factored into assessments.


NNR Food Group 8: Meat and meat products

Global context

At the global level, chicken meat is produced in the largest quantity (120 mt), followed by pig meat (110 mt), cattle meat (68 mt), sheep meat (10 mt), and goat meat (6 mt) – which together comprise the top 5 in terms of global meat production quantity in 2020.²⁶³

The environmental impacts of meat can vary substantially depending on the meat type and production method (figure 8.1 and table 8.1). Across the entire product life cycle, most of the impacts from meat production occur during production (~80%), with a relatively small proportion resulting from processing, packaging, and transport. For some meat types and impacts, crop production (for animal feed) accounts for most of the burden e.g. eutrophication for beef meat production from dairy herds, pig meat and poultry meat. Land use change accounts for the largest proportion of GHGs from poultry meat (figure 8.1). For the meats from ruminant animals (beef, lamb and mutton), the largest proportion of GHGs arise directly from the animals in the form of CH₄ from enteric fermentation during digestion.²⁶⁴ For beef meat, the majority of all impacts are directly associated with animal farming e.g., eutrophication and acidification from animal waste (figure 8.1).

Figure 8.1: Proportion of environmental impacts from different life cycle stages: Meat and meat products (global)²⁶⁵



Life cycle assessments of meat products have revealed a general trend in relation to impacts. Compared to meat grown using extensive (or organic) techniques, meat grown using intensive (or conventional) production methods tends to have lower impacts per kilogram across their life cycle in terms of land use, eutrophication and acidification – but similar impacts in terms of GHGs and higher energy use.²⁶⁶ However, extensive methods might have lower impacts in terms of biodiversity loss and soil degradation (if production amounts were equal).²⁶⁷ Extensive ruminant systems generally use a higher share of grassland and less arable land, utilize more permanent grasslands and have higher organic matter applications to soil than intensive systems – all of which can increase soil carbon stocks in extensive production compared to intensive production (depending on current soil carbon content). However, soil carbon stocks are generally not included in such assessments due to a lack of data and methodological limitations.²⁶⁸ The differences between the environmental impacts of intensive and extensive (and broadly conventional vs organic) production occur mostly due to a higher allocation of land (or area) and a longer lifetime of the animal under extensive methods due to a reduced productivity from a lower macronutrient density and digestibility of feed.²⁶⁹ Regardless of subsequent production methods, if initial land clearance for production displaces a native or pristine ecosystem such as primary forest, the impacts of such land use change could be substantial at least in terms of GHGs.²⁷⁰

Table 8.1: Average environmental impacts per kg of retail weight: Meat and meat products (global)²⁷¹

	Land use (m ² /kg)	Greenhouse gas emissions (kg CO ₂ eq/kg)	Acidifying emissions (g SO ₂ eq/kg)	Eutrophying emissions (g PO ₄ ³⁻ eq/kg)	Freshwater (L/kg)

Bovine Meat (beef herd)	326.2	99.5	318.8	301.4	1451
Bovine Meat (dairy herd)	43.2	33.3	343.6	365.3	2714
Lamb & Mutton	369.8	39.7	139.0	97.1	1803
Pig Meat	17.4	12.3	142.7	76.4	1796
Poultry Meat	12.2	9.9	102.4	48.7	660

Depending on where and how feed crops are grown, they might also incur environmental impacts related to land use change, such as deforestation, including GHG emissions and biodiversity loss. Similarly, feed crop production might displace native carbon sinks such as peatland, where they are drained and converted to cropland. Relevant issues related to crop production (in this case, for feed production), are covered in chapters 3 and 5. Animals raised on pasture also have feed requirements, such as over winter where grass production may be less viable, and finishing feed to increase animal weight before slaughter. Pasture lands also have input requirements, for example permanent pasture in the UK is the largest nitrogen user across all agricultural production.²⁷²

Another important consideration is the localised, or site specific impact. For example, manure management and/or use of fertilizers and pesticides could have different impacts on the local environment depending on site-specific conditions, including the scale of operations, and due to nitrogen retention as a result of soils and drainage systems.²⁷³ Stocking density of farmed animals also plays an important role in determining the level and extent of environmental impacts.²⁷⁴ The severity of impacts could also vary depending on the types of inputs – for example, pesticides relatively high in toxic substances could have a more detrimental environmental impact compared to less toxic substances – however this is generally not reflected by the most widely used environmental assessments²⁷⁵ and therefore needs to be addressed in more detail in future studies.

While agricultural land used for grass production (including silage) can result in lower CO₂ emissions from degradation of soil carbon stock per ha in comparison to agricultural land used for monocultures of annual crops²⁷⁶, all land used for meat production, whether it is feed cropland (including silage) or pasture, and whether it is cultivated under organic or non-organic methods, has an associated 'carbon opportunity cost' related to the native vegetation cover that would occur if meat production ceased. Globally, this is mostly related to permanent pastureland that has replaced forest.²⁷⁷

The production of meat, dairy and eggs is the biggest single use of land globally – occupying 38% of all habitable land and 78% of agricultural land.²⁷⁸ The production of feed crops uses 43% of global cropland.²⁷⁹ Farming animals is a major driver of biodiversity loss – being the biggest source of nitrogen pollution (leading to oxygen depleted 'dead zones' in rivers and oceans through the process of eutrophication), ammonia deposition in landscapes, and land use and land use change, including deforestation.²⁸⁰ For example, cattle farming is the single largest direct cause of deforestation, and soy production is the third largest (76% of soy production is used for animal feed, mostly to feed pigs and chickens).²⁸¹ Biodiversity loss also occurs through the killing of animals that are considered detrimental to livestock farming, for example through predation or spread of disease.²⁸² Within key biodiversity areas, cattle farming has been identified as the largest single cause of biodiversity loss (accounting for 31%), with animal farming in total causing more than half of the loss of plants and vertebrates. From the perspective of land use, lightly grazed pasture was associated with half of all biodiversity loss - concentrated in middle- and low-income regions with rich biodiversity.²⁸³ Land classified as permanent pasture occupies 24% of global land area, and temporary pastures occupy

1%.²⁸⁴ The largest meta-analysis conducted on the biodiversity impacts of grazing farmed animals shows that only detritivores (e.g., dung beetles) benefit – all other species of animals and plants analysed decline in abundance and diversity in areas of livestock grazing.²⁸⁵

Beef production

The impact of beef varies mostly depending on whether the meat is sourced from animals grown specifically for meat, or whether the meat is sourced from animals grown for both meat and dairy. For example, in terms of GHG emissions, according to a global average using retail weights, beef from a beef herd has around 3 times the impact of beef from a dairy herd (99.5 kg CO₂e/kg compared to 33.3 kg CO₂e/kg²⁸⁶). The difference is 7-fold greater in relation to land use, but is lower in relation to acidifying emissions (0.9), eutrophying emissions (0.8) and freshwater use (0.5)²⁸⁷ (table 8.1).

Production methods also cause variation in environmental sustainability outcomes. Beef produced from grain fed cattle uses less land and causes fewer GHG emissions and less eutrophication per unit of production compared to beef from grass-fed cattle, but has higher energy requirements. Cattle raised in both systems have feed, land, water and energy requirements and produce manure - the environmental impacts of which can also vary substantially depending on management techniques.²⁸⁸

When considering such trade-offs at the aggregate or system level, demand for beef is an important consideration. If demand for beef declines substantially from current levels, there is more opportunity to focus production on extensive organic methods as the overall system-level land requirement for beef would decline. Adopting extensive organic beef production globally, and even nationally in some cases, would only be possible in combination with a substantial reduction in beef consumption from current levels. For example, a recent analysis of the US found that limiting production to exclusively pasture-fed beef would require reducing beef production to 27% of current levels – or a more than 3-fold increase in pasture area to continue current levels.²⁸⁹ Adoption of integrated mixed farming systems (e.g., extensive livestock coupled with diverse rotations) could be beneficial from a broad sustainability perspective, if demand for ruminant meat declined sufficiently to reduce incentives to expand agricultural land use.²⁹⁰ The main trade off (or opportunity cost) in this situation is the large area of land use that has typically displaced a more carbon- and biodiversity-rich habitat such as forest or grassland. If land-based climate change mitigation and biodiversity needs are considered to be more beneficial than livestock production, the land used for extensively raised organic cattle could instead be restored to native vegetation cover to maximize its carbon sink potential (such as reforestation of pastureland) and in turn provide opportunities for biodiversity that would occur in such native habitats.

Beef production has a substantial climate impact – accounting for 6% of total global anthropogenic GHGs.²⁹¹ Farmed ruminant animals contribute the largest source of anthropogenic methane (CH₄), accounting for around 30% of global emissions.²⁹² While the environmental impacts of beef production could be to some extent reduced through various technological approaches (see section ‘general points across the food groups’), and through utilizing waste streams from other systems for some portions of feed, these improvements alone are not enough. To align with planetary boundaries, reducing overall demand for beef at the global system-level remains essential even when optimistic improvements in production methods (to reduce environmental impacts per unit of production) are considered.²⁹³

Pork production

Several of the issues and trade-offs related to beef production also apply to pig production in terms of large land use, biodiversity impacts, GHGs, feed requirements, and manure waste, albeit in different relative proportions and scales. Pig meat tends to have higher environmental impacts across a range of metrics in comparison to chicken meat, but much lower impacts in comparison to beef meat (including dairy herd) and sheep meat (table 8.1). The largest proportions of overall environmental impacts from pig meat production tend to be a result of feed production and manure management (figure 8.1).²⁹⁴ In terms of GHGs, feed has impacts via land use change (e.g., deforestation to create cropland for soybeans), and also directly in the form of nitrous oxide emissions from nitrogen fertilizer use and manure, and carbon dioxide from fuel used in field machinery and CH₄ emissions from rice production. Hence, the total impacts of feed depend on whether they are driving land use change and the extent to which they generate impacts during production.²⁹⁵ Relevant issues related to crop production (in this case, for feed production), are covered in more detail in chapters 3 and 5.

Chicken meat production

Many of the same issues apply to the production of chicken, also a monogastric animal, as for pig meat production. Several of the issues and trade-offs related to land use, biodiversity, GHGs, feed requirements, water use, and manure waste, are the same albeit in different relative amounts when assessed on a per weight unit basis for example. As for pigs, the largest proportions of environmental impacts from chicken meat production tend to be a result of feed production and manure management (figure 8.1).²⁹⁶ Again, the total impacts of feed largely depend on whether they are driving land use change and the extent to which they generate impacts during production (such as eutrophication of waterways) – see chapters 3 and 5 for more details on the impacts of crop production.


Chicken meat production has risen by 10% globally in the past 5 years alone (and is the fastest growing sector in terms of livestock production).²⁹⁷ Another important consideration is the prospect of shifting from other types of meat with higher environmental impacts, to chicken which, in comparison to the top 5 most highly produced meats, tends to have the lowest environmental impacts across a range of metrics on a per kg basis (table 8.1). However, to be able to assess the impact of shifting from other meat types to chicken, the full range of environmental impacts at the absolute level, and their relation to planetary boundaries, and other environmental risks (e.g., from increased use of antibiotics, spread of zoonotic disease, increased number of farmed animals) must be considered. The prospect of increasing deforestation for feed production is also relevant – for example, around 28% of soy produced globally is used to feed farmed chickens (figure 5.2).

Nordic and Baltic context

It is necessary to reduce meat consumption to some degree across the Nordic and Baltic region. The current levels of red meat intake are higher than the maximum levels recommended in health based dietary guidelines (table 2). Figure 8.2 shows the variation in terms of impacts of meat consumption across the Nordic and Baltic countries and how they relate to the food portion of global environmental limits. For example, if everyone ate the same amount of chicken/pork as Lithuania it would use over 65% of the global cropland limit for food consumption. If everyone ate the same

amount of beef/lamb as Iceland the threshold for GHGs from food consumption would be exceeded by 4.3 times (figure 8.2). Hence, reducing meat consumption would have considerable beneficial environmental impacts.

Figure 8.2: Impacts of meat consumption across the Nordic and Baltic countries in relation to the food portion of global environmental limits (%)



Source: *Global Nutrition Report 2021*²⁹⁸ The analysis utilizes country-specific food consumption and environmental footprint data and relates them to the food portions of global environmental limits i.e. a global test to assess the impacts if everyone in the world consumed at the given rate. The methods and data are described in Box 1, and presented in appendices 1-3.

While the data in table 8.1 suggest a shift from beef to pork or chicken would reduce GHGs, it is important to explore the realities and trade-offs of such an option. For example, shifting some beef consumption to pork and/or chicken could reduce GHGs, but increase cropland, water and nitrogen use to produce feed crops (figure 8.2), and possibly cause further land use change, and biodiversity loss. However, the overall area of pastureland would decline due to less cattle – some of which could be restored to its native habitat cover, in turn sequestering carbon and enhancing biodiversity. Hence, depending on the priorities, such changes could shift rather than solve environmental problems. The current intake of beef, chicken and pig meat in the region causes large impacts on land use, GHGE, and nitrogen and phosphorus application (Figure 8.2), which suggests that a substantially reduced intake of all meats would be optimal in terms of meeting planetary health goals.

The environmental sustainability issues outlined in relation to the global context above also apply to meat production and consumption in the Nordic and Baltic region. However, there are also a number of important location specific issues to consider. Generally, freshwater use is not currently a major problem in Nordic production because of low irrigation rates and abundant water resources. The main issues with water use relate to imported foods produced outside of the region.²⁹⁹ Nitrogen pollution is a major issue and is largely due to pollution from animal agriculture (manure and feed crop production). The Baltic Sea is one of most polluted seas in the world³⁰⁰ and the Oslo fjord is being increasingly polluted by nitrogen from agricultural run-off.³⁰¹ This is a substantial constraint to livestock production and limits other uses of the water bodies. Limits on pollution are required³⁰², although strict regulations of fertilizer use and manure management have resulted in large

reductions in nitrogen emissions in Denmark.³⁰³ A potential precedent has been recently established in the Netherlands, after the European Court of Justice ruled that the state had not taken sufficient action on reducing nitrogen pollution, and resulted in targets being identified to halve livestock numbers in order to tackle the problem.³⁰⁴

Although some environmental impacts of meat from farmed animals might be lower if produced in the Nordic countries compared to some countries that export to the Nordic countries, consumption of locally produced meat, assuming current consumption is maintained, does not sufficiently reduce environmental impacts, as indicated in Nordic studies on dietary patterns.³⁰⁵ Local production does have some practical benefits compared to meat imports in that the storage and transport issues are reduced which can be important for such highly perishable products.

Although in the Nordic region livestock production on semi-natural grasslands does not always directly compete with food production for human consumption, local production of substantial amounts of crops for animal feed has a major environmental impact. This is a problem across the region and highlights the competition for feed versus food. Reconfiguring feed crops for direct human consumption (where possible) in combination with a reduction in livestock production and consumption could therefore be a method for reducing the environmental impacts of food production within the region.^{306,307,308}

Despite biodiversity issues (including those related to feed imports) being particularly important in Sweden, Finland, Norway, Iceland and Denmark, it is not possible to adequately assess them across all countries due to a lack of comparative data. Although land use change e.g., from forest to agriculture is not a major issue in the Nordic and Baltic countries, the negative impacts of livestock production on terrestrial biodiversity are considered a serious threat.³⁰⁹ Across the EU27 (which includes Denmark, Estonia, Finland, Latvia, Lithuania and Sweden), livestock production has been estimated to cause 26% of total biodiversity loss (measured as mean species abundance), mostly through feed production on grasslands and croplands.³¹⁰ There are limited contexts where low density livestock grazing is used to maintain semi-natural grasslands, which are linked to certain types of biodiversity and may contribute to cultural heritage, social cohesion and recreation - particularly in Sweden, Norway and Estonia for example.³¹¹ However, these areas are currently being threatened more by structural changes in agriculture rather than climate or environment related policies. It is important to expand current knowledge in order to assess biodiversity impacts alongside other important environmental metrics. Even with adequate data, it will still be impossible to 'weigh' the impacts e.g., how many units of biodiversity at the local level are equal to how many units of GHGs? Intensive animal production, in general, seems to have little or no benefit for species richness,³¹² or to carbon sequestration³¹³ – but generally has lower environmental footprints in terms of land use and GHGs in comparison to extensive production. Changes in production would therefore likely be assessed using value-based judgements in relation to impacts on biodiversity, and in dealing with trade-offs against other environmental impacts.

There are also major issues related to imported feed. For example, in Norway and Sweden, the majority of imported soy, and in Finland approximately half, was certified as deforestation-free³¹⁴ in 2020, yet in the other Nordic and Baltic countries, the proportion was smaller or negligible³¹⁵ (although there are some potential issues with transparency and shifting demand due to rising costs in relation to certification schemes³¹⁶). In terms of reducing the impact of imported feed, the availability of lower impact feed alternatives and/or whether they can be produced in-country (rather than being imported from another country) could be important areas of inquiry if not already in progress. In Denmark for example, farmers have begun partly replacing soy imports used for

farmed animal feed with vicine-free fava beans.³¹⁷ Similarly, in Finland the Leg4Life initiative is operational.³¹⁸

Iceland dedicates by far the largest proportion (99%) of its agricultural land to pasture (mostly permanent), followed by Norway (67%, mostly temporary), Sweden (54%, mostly temporary), Estonia (48%, mostly permanent), Latvia (47%, mostly permanent), Finland (40%, mostly temporary), Lithuania (34%, mostly permanent), and Denmark (29%, most temporary) (table 1). Large areas of land are used for sheep production in Iceland which causes biodiversity loss and damage to ecosystems via sheep grazing on emerging vegetation and is a particular problem in sensitive areas.³¹⁹

In Sweden, domestic production is dominated by beef from the dairy sector (64%)³²⁰ i.e. from culled cows and offspring raised for meat. There has been a large reduction in dairy cows due to increasing milk yield and these have been replaced by an increasing number of suckler cows. Most bulls are raised indoors to a slaughtering age of between 15 and 24 months on a diet of approximately 50% forage and 50% grains.³²¹ The use of soy for feed is low in beef production and 2.5 % in dairy production (which a substantial part of beef comes from).³²² 15% of slaughtered beef animals are raised under organic production methods. Suckler cows (and their calves), dairy cows and heifers graze in summer (required by Swedish animal welfare legislation). Grass-clover ley grown in rotation with other crops or in monocultures is the main feed source.³²³

In Denmark the largest proportion of all beef originates from dairy systems (83%), estimated from data on the production of bone-free meat from different types of cattle in Denmark. Intensive beef accounts for 15% and extensive beef for 2%, which is mostly under organic production. Within the dairy systems, dairy cows supply the largest share of minced meat and dice/strips (55% and 63%, respectively), whereas calves supply the largest share of steaks (59%). Beef produced from dairy cows in the Nordic countries appears to result in lower environmental impacts compared to beef produced from beef cattle. Mogensen et al., 2015³²⁴ estimated that GHGs from beef produced in Denmark and Sweden from dairy bull calves slaughtered between 9 and 19 months old had the lowest GHGs, ranging from 9 to 12 kg CO₂/kg of carcass weight. Comparatively, GHGs of beef from beef breed systems ranged from 23 to 30 kg CO₂/kg carcass weight. The differences were largely attributed to differences in feed intake over the animal's lifetime. A study of beef production in Finland³²⁵ estimated a range of environmental impacts of (carcass weight) beef produced from dairy cows and beef cattle respectively: GHGs were 25 and 32 kg CO₂eq/kg; aquatic eutrophication was 22 and 33 g PO₄³⁻eq/kg; and acidification potential was 58 and 64 gAE-eq/kg. In these examples, the environmental impacts were higher for beef produced from beef cattle compared to beef produced from dairy cows, which is also apparent for GHGs and land use based on a global average of beef produced from beef herds and beef produced from dairy herds, but not for acidifying and eutrophying emissions (table 8.1).³²⁶

In Finland, the majority of pastures are intensively farmed and are not adequately supporting biodiversity.³²⁷ It is not necessarily possible to expand agricultural land in Finland as this could result in even more peatland being brought into production (currently 8 - 11% of agricultural land is peatland).³²⁸ Reducing the demand for agricultural land through shifting to more plant-based diets would allow peatland to be taken out of production, rather than replace grass production on peatland with grain production for example, which could increase GHG emissions.³²⁹ A reorganization of crop rotations on the remaining agricultural land would be required to maintain or improve the carbon stock and agricultural condition of farmland.³³⁰

In Norway, a large area of land used for temporary pasture is deemed unsuitable for crop production under current market incentives. Most (~90%) crops produced in Norway are used for farmed animal feed. This includes grass and roughage production, but also around two thirds of the grains produced are feed rather than food grains. Outdoor grazing is limited due to seasonal reasons. When indoors, cows receive a mixture of roughage and concentrates. However, recent analysis suggests that more of the cultivated land could be used for producing plant-based foods.³³¹ An increase in vegetable production (including potatoes) is also possible, and portions of feed could be reallocated to human food (e.g., oats).³³² Structural changes have increased the size of farms, with much fewer small farms now operating. Due to smaller production and long distances, production in more marginal areas is no longer financially profitable.³³³ Improving incomes and maintaining viable districts with livestock production is considered a priority area in Norway, and has resulted in conflicting agricultural and environmental policy goals.³³⁴ This demonstrates some of the socio-political challenges in terms of reducing the environmental impacts of food production and consumption within the region and the importance of addressing such issues in order to progress toward more environmentally sustainable agriculture.

The overarching issue across the Nordic and Baltic region in relation to livestock production is land use. If, where and how to reduce production is likely to be considered from the perspective of value judgements which currently dominate decisions around the relative value of maintaining livestock production versus increasing biodiversity and mitigating climate change. The favouring of the latter against the former is largely dependent on whether the focus is on maintaining the status quo (i.e. current income sources, current land use patterns, current levels and types of biodiversity, current response to meeting the Paris Agreement goals), or taking a more ambitious approach to tackling climate and biodiversity issues, and aligning income sources with those goals.³³⁵

A broader point is that current agricultural land use across the countries could usefully be examined to identify 'best' uses for maximizing environmental sustainability outcomes. For example, fully repurposing some areas currently under permanent pasture would allow native vegetation and ecosystems (e.g., the Boreal forest or wetlands in relevant countries), to regenerate, and wildlife to be reinstated (and predator- prey relationships), providing additional benefits in terms of climate change mitigation, biodiversity, and more functional ecosystems. Any such changes to land use might require identification of alternative or replacement income sources.

Beef production and trade

Of the 8 countries, Sweden is the largest beef producer, but none of the countries contribute substantially to global beef production. All countries except Latvia rely on beef imports to some extent to supply national consumption (table 8.2). In countries with a greater reliance on imports, there is less opportunity to influence the environmental impacts of current consumption through changing production methods. However, all countries could influence environmental impacts by reducing national beef consumption, and focusing on production from systems (home and abroad) with lower environmental impacts. Supply levels vary quite substantially across the 8 countries – with levels in Denmark more than four times those in Lithuania (table 8.2). Hence, for some countries there is scope for reducing national beef consumption more significantly.

For comparison, beef consumption in line with a flexitarian diet that aligns with planetary boundaries up to 2050 (if implemented in combination with medium-ambition technological measures to reduce inputs during food production, and reductions in food loss and waste of 75%), recommends a daily maximum beef intake of 7g, or 2.6 kg per year (as part of an average daily

intake of 2,100 kcal).³³⁶ Based on the annual supply data in table 8.2, all countries exceed this amount (on a per person basis) 2 – 9 fold. Similarly, using red meat consumption data from table 2 indicates that consumption in all countries far exceeds the amount consistent with the flexitarian diet.

Table 8.2: Beef production and supply across the Nordic and Baltic countries in 2018³³⁷

Country	Production (1000 tonnes) (% of global total)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	129 (0.2)	124	107	146	23.52
Finland	87 (0.1)	27	4	110	19.26
Iceland	5 (<0.0)	1	0	6	14.89
Norway	89 (0.1)	13	0	102	17.66
Sweden	137 (0.2)	113	12	238	22.58
Estonia	13 (<0.0)	8	4	17	8.64
Latvia	17 (<0.0)	7	13	11	5.73
Lithuania	42 (0.1)	8	33	17	5.36

Pork production and trade

Denmark accounts for 1.3% of global pig meat production and is by far the largest producer from the 8 countries. All countries except Denmark rely on pig meat imports to some extent to supply national consumption (table 8.3). Hence, there could be greater potential to reduce the environmental impacts of production in Denmark compared to other countries that rely more on imports. However, all countries could influence environmental impacts by reducing national consumption, and focusing on production from systems (home and abroad) with the most transparent environmental impact data. Supply levels vary quite substantially across the 8 countries – with levels in Lithuania almost twice those in Denmark. Hence, for some countries there is scope for reducing national consumption more significantly. For comparison, pork consumption in line with a flexitarian diet that aligns with planetary boundaries up to 2050 (if implemented in combination with medium-ambition technological measures to reduce inputs during food production, and reductions in food loss and waste of 75%), recommends a daily maximum pork intake of 7g, or 2.6 kg per year (as part of an average daily intake of 2,100 kcal).³³⁸ Based on the annual supply data in table 8.3, all countries exceed this amount (on a per person basis) 8 – 19 fold. Similarly, using pig meat consumption data from table 2 indicates that consumption in all countries far exceeds the amount consistent with the flexitarian diet, except in Iceland (although data is missing for 4 countries in table 2).

Table 8.3: Pig meat production and supply across the Nordic and Baltic countries in 2018³³⁹

Country	Production (1000 tonnes) (% of global total)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	1,583 (1.3)	148	1,445	286	26.7
Finland	169 (0.1)	42	23	188	37.7
Iceland	7 (<0.0)	1	0	8	20.9
Norway	137 (0.1)	6	5	138	22.9

Sweden	249 (0.2)	111	29	331	30.9
Estonia	42 (<0.0)	40	28	54	37.8
Latvia	39 (<0.0)	58	11	86	43.6
Lithuania	72 (<0.0)	89	19	142	48.4

Chicken meat production and trade

Of the 8 countries, Sweden is the largest chicken meat producer, but none of the countries contribute substantially to global chicken meat production. All countries except Denmark and Lithuania rely on chicken meat imports to some extent to supply national consumption (table 8.4). Hence, there is significant potential to reduce the impacts of both production and consumption in Denmark and Lithuania. In countries with a greater reliance on imports, there is less opportunity to influence the environmental impacts of chicken production. Latvia and Estonia who import more than they produce, have least ability to influence the environmental impacts of production given that it is mostly served via imports. However, all countries could influence environmental impacts by reducing national consumption, and focusing on production from systems (home and abroad) with the most transparent environmental impact data. Supply levels vary quite substantially across the 8 countries – with levels in Lithuania almost twice those in Sweden. Hence, for some countries there is scope for reducing national consumption more significantly and replacing where needed with more environmentally sustainable alternatives such as legumes and pulses.

For comparison, poultry meat consumption in line with a flexitarian diet that aligns with planetary boundaries up to 2050 (if implemented in combination with medium-ambition technological measures to reduce inputs during food production, and reductions in food loss and waste of 75%), recommends a daily maximum poultry meat intake of 29g, or 10.6 kg per year (as part of an average daily intake of 2,100 kcal).³⁴⁰ Based on the annual supply data in table 8.4, all countries exceed this amount (on a per person basis) by around 2 – 3 fold. In contrast, using chicken meat consumption data from table 2 indicates that consumption in Finland, Iceland, Lithuania and Latvia exceeds the amount consistent with the flexitarian diet, whereas other countries consume below or close to this amount.

Table 8.4: Poultry meat production and supply across the Nordic and Baltic countries in 2018³⁴¹

Country	Production (1000 tonnes) (% of global total)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	156 (0.12)	141	142	155	26.9
Finland	135 (0.11)	17	11	141	19.4
Iceland	9 (0.01)	1	0	10	30.9
Norway	98 (0.08)	2	0	100	20.6
Sweden	161 (0.13)	81	26	216	16.9
Estonia	19 (0.02)	30	12	37	22.2
Latvia	34 (0.03)	47	21	60	21.0
Lithuania	134 (0.11)	45	66	113	29.0

Considerations for reducing environmental impacts of meat consumption in Nordic and Baltic countries

- A reduction in meat consumption is a first order priority – with greatest scope in countries consuming above levels compliant with planetary boundaries (mostly beef and pork across all countries, but also chicken for some countries).
- If meat consumption needs to be replaced and not just reduced (for example to meet recommended nutrition intakes), replacement with plant-based foods such as legumes, pulses and whole grains would reduce the impacts of meat consumption as much as possible and avoid trade-offs to the greatest extent (in comparison to replacing beef with chicken or pork for example). The environmental impacts of replacing some meat consumption with the lowest impact seafoods could be assessed, with a requirement to avoid problem shifting.
- Reducing meat production and consumption would reduce the demand for feed crops (see related considerations in sections 3 and 5 for cereals and pulses/legumes), hence replacing meat with plant-based foods would not require an increase in cropland (rather a reconfiguration of crop production to best suit human needs).
- The increased use of locally grown legumes as food could give incentives for farmers to include legumes in the crop rotations, which in turn would reduce the requirements for synthetic nitrogen fertilisers and improve soil quality. Other plant-based meat replacement products might also be suitable alternatives – however this would need to be assessed depending on type and availability. Potential food shifts will be discussed further in the next paper of this series.
- Changing elements of production such as replacing feed with legally permitted and safe by-products and shifting to more extensive methods might offer benefits in some contexts. However, such options do not negate the primary need to reduce meat consumption substantially. For example, shifting all beef production to extensive methods would require a reduction in overall beef production levels to avoid increasing GHG emissions and land use.³⁴²
- Given the range of factors involved and the complexity of trade-offs and location specific impacts and contexts, a national strategy would be important to usefully direct the reduction and replacement of meat production and consumption. An important part of this would be a comprehensive review to identify and explore the relevant range of issues including human health and social impacts, animal welfare, and current and emerging threats e.g., zoonotic-diseases (and the risks of emerging zoonoses with potential pandemic consequences), and environmental change such as temperature extremes.
- Overarching, and in parallel, a national land use assessment could inform optimal land uses for meeting a range of environmental goals, also accounting for the environmental impacts of food imports in producer countries. One important inclusion would be an assessment of different types of pasture lands in terms of their current value and necessity for food production versus alternatives uses, such as restoring native habitats to help meet other social goals (i.e. meeting climate and biodiversity goals). For example, on native grasslands, improved grazing management could partially restore vegetation and related carbon stores but would come with a trade-off of non-CO₂ GHGs from the grazing ruminants. Fully restoring pasture land to native forest cover (where relevant) removes this trade off, and

could have large scale potential to sequester more carbon and provide habitats for a wider range of wildlife across the region.³⁴³

- Ensuring a trajectory within planetary boundaries requires a longer-term perspective on the role of livestock within the food system rather than focussing only on the realities of today’s market and other immediate socio-political determinants. Situations can change rapidly and business as usual cannot be assumed under the expected impacts of climate change alone.


NNR Food Group 9: Milk and dairy products

Global context

At the global level, 876 mt milk (excluding butter) were produced in 2019 of which 3% were used for farmed animal feed. The majority was produced from cows (81%), followed by buffalo (15%), goat (2%), sheep (1%) and camel (<1%). India produces by far the largest quantity of milk (187 mt in 2019), followed by the US (99 mt), Pakistan (56 mt), China (36 mt), and Brazil (35 mt) – making up the top 5 producers globally.^{344,345}

Collectively, dairy production has a large environmental footprint at the global level, accounting for 4% of total anthropogenic GHGs for example.³⁴⁶ This is largely a result of the use of ruminant animals for the majority of milk production (cows, buffalo, goat, and sheep). The environmental impacts of dairy products tend to be concentrated during the production stages of the life cycle, which account for around 80% of impacts. For example, in terms of GHGs, the animal farming component accounts for the largest share of emissions as a result of methane and nitrous oxide released from the ruminant animals and from their manure. Whereas acidification and eutrophication impacts are largely attributed to feed crop production (figure 9.1).

Figure 9.1: Proportion of environmental impacts from different life cycle stages: Milk and dairy products (excl. butter) (global)³⁴⁷



Cheese has a much higher impact compared to dairy milk as it is a concentrated form of milk, requiring around 10 litres of milk to produce 1 kg of cheese (table 9.1). The impacts of butter are not described here, but animal fat is included in section 11 on Fats and oils.

Table 9.1: Average environmental impacts per kg of retail weight: Milk and dairy products (global)³⁴⁸

	Land use (m ² /kg)	Greenhouse gas emissions (kg CO ₂ eq/kg)	Acidifying emissions (g SO ₂ eq/kg)	Eutrophying emissions (g PO ₄ ³⁻ eq/kg)	Freshwater (L/kg)
Milk	9.0	3.2	20.0	10.7	628
Cheese	87.8	23.9	165.5	98.4	5605

Many of the same issues discussed in chapter 8 related to meat production (e.g., feed production, deforestation, large land use, manure and urea, nutrient pollution of surrounding ecosystem, resource use on farm including water and energy for housing, culling of local wildlife), apply to dairy production. Additionally for dairy products, animal welfare and antibiotic use are also important issues to consider,³⁴⁹ but are outside the scope of the current paper.


As for the impacts of meat production, the level of impacts per unit of milk production vary depending on a number of factors including milk yield (which is not uniform across the global herd of farmed cows), and feed type, production methods, and production location. For example, locally grown feed produced by nitrogen fixing legumes will potentially have lower environmental impacts compared to imported grains. CH₄ emissions from ruminant animals can also vary depending on feed type and its digestibility (higher cellulose feeds result in higher CH₄ release from the animals), and emissions from manure can also vary for example, depending on temperature, amounts, oxygen availability and storage type.³⁵⁰ How environmental impacts are reported per unit of milk are also dependent on the allocation method used in the life cycle assessment, for example, if economic allocations are applied, a greater proportion of environmental impacts over a cow's life will be attributed to meat production rather than milk production. However, on a weight basis, most of the environmental impact would be allocated to milk production.³⁵¹

Nordic and Baltic context

The Nordic and Baltic countries have some of the highest levels of dairy milk consumption in the world – for example, Finland currently has the second highest global consumption rates, with Estonia having the 6th highest, Denmark the 9th highest, and Iceland the 18th highest.³⁵² Environmental impacts of dairy consumption are also high across the countries, making up a substantial proportion of total impacts from all food consumption – particularly in relation to GHGs and land use, but phosphorus and nitrogen are also significant. Reducing dairy consumption is the most effective way to reduce environmental impacts from this food group and will be necessary to align with food-related global environmental limits.

Figure 9.2 shows the variation in environmental impacts of dairy milk consumption across the Nordic and Baltic countries and how they relate to the food portion of global limits. For example, if everyone in the world consumed the same amount of dairy as each person in Finland, the global GHG limit for food consumption would be exceeded by almost 1.4 times and would use 60% of the cropland limit for food consumption.

Figure 9.2: Impacts of milk consumption across the Nordic and Baltic countries in relation to the food portion of global environmental limits (%)



Source: *Global Nutrition Report 2021*³⁵³ The analysis utilizes country-specific food consumption and environmental footprint data and relates them to the food portions of global environmental limits i.e. a global test to assess the impacts if everyone in the world consumed at the given rate. The methods and data are described in Box 1, and presented in appendices 1-3.

Many of the issues covered in the global context regarding the environmental impacts of dairy are also relevant to the Nordic and Baltic countries. There are some regional and local variations, for example, some regions are more dependent on feed imports (e.g., rapeseed cake in Finland) due to shorter grazing seasons and less local production of feed crops. Dairy cows are generally fed a mixture of roughage (silage, hay, pasture, maize) and various types of concentrates derived from crops such as cereals, rapeseed, soya bean, beet pulp and molasses. The impacts of feed production vary depending on which crops are used, and where they are sourced from – for example, regulations on plant crop protection varies across the world and therefore soybeans imported from Brazil could have different impacts compared to locally produced pulses. In Sweden soybean use for dairy cows is approximately 2.5% of the feed.³⁵⁴ The amount of soy that Norway imports for animal feed would cover more than 50% of the Norwegian population's basic protein needs.³⁵⁵ Production methods are also important, for example intensively grown animal feed can result in soil erosion within the region (see chapters 3 and 5).³⁵⁶

Intensively produced dairy has a range of environmental issues across the region. Generally, dairy farms have become fewer in number and larger in size, which concentrates certain impacts in the local environment, such as nutrient pollution. Farm location also impacts energy use – for example, areas with shorter grazing seasons will need to use relatively more energy for lighting and ventilation for animal housing.³⁵⁷

Some variation exists in terms of environmental impacts in comparison to the global average of 3.2 kg CO₂e/kg retail weight in table 9.1. For example, a Swedish study³⁵⁸ estimated the impact for energy corrected milk (ECM), including surplus calves and culled cows, to be 1.16 kg CO₂e/kg ECM in Sweden, similar to an estimate of milk in Denmark (1.06 kg CO₂e/kg ECM).³⁵⁹ However, the amount used in figure 9.2 to estimate the GHG impact of milk consumption across the Nordic and Baltic countries in relation to planetary boundaries is 1.39 kg CO₂e/kg (see appendix 2). The main route for reducing the environmental impacts of dairy production is a reduction in production amounts, to

reduce absolute impacts. The main route for reducing GHGs per unit of production (relative impacts) for both organic and non-organic is increasing milk yield.³⁶⁰ However, this is already very high and will not reduce the absolute impact of dairy production if production levels increase with increasing demand. The impact of plant protection products can vary the impacts of organic relative to non-organic production. For example, some chemical products can be used in conventional milk production via feed crops, but are not permitted in organic crop production (although small amounts of non-organic feed might be permitted for organic dairy production in some countries such as Sweden).³⁶¹ Using a larger proportion of locally and regionally produced organic feeds such as grains, oil seeds and pulses, some feed from pastures under extensive production,³⁶² and potentially low-opportunity-biomass (i.e. biomass not in competition with human food)³⁶³ could help to reduce the impacts of dairy per unit of production, however this does not fully address a number of important issues including the competition for agricultural land (e.g., feed versus food), and the need for overall dairy consumption levels to decline across the region.

Denmark is by far the largest producer of dairy milk, followed by Sweden and Finland. Denmark is also the largest net exporter of dairy milk, followed by Estonia and Latvia. All other countries, except Iceland, are net importers of dairy milk. Finland is the largest dairy milk consumer in the region on a per person basis, followed by Estonia and Denmark. Lithuania has the lowest consumption levels (table 9.2). Milk from cows accounted for by far the majority of production and consumption in all countries.

For comparison, milk consumption in line with a flexitarian diet that aligns with planetary boundaries up to 2050 (if implemented in combination with medium-ambition technological measures to reduce inputs during food production, and reductions in food loss and waste of 75%), recommends a daily maximum milk intake of 250g, or 91.3 kg per year (as part of an average daily intake of 2,100 kcal).³⁶⁴ Based on the annual supply data in table 9.2, all countries exceed this amount (on a per person basis), with Lithuania being closest with an average daily consumption of 296g. Average supply in Finland exceeds the amount by 3.7 times, followed by Estonia (3 times), Denmark (2.6 times), Iceland (2.2 times), Sweden (2 times), Norway (1.8 times), and Latvia (1.8 times). In contrast, using milk and cheese consumption data from food diaries in table 2, consumption in all countries exceeds the amount consistent with the flexitarian diet, except for Latvia and Lithuania.

Table 9.2: Milk (excluding butter) production and supply across the Nordic and Baltic countries in 2019³⁶⁵

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	5,615	934	3,148	3,401	238
Estonia	822	81	376	527	278
Finland	2,374	759	209	2,924	336
Iceland	156	1	7	150	203
Latvia	981	260	554	687	165
Lithuania	1,551	657	629	1,579	108
Norway	1,573	131	90	1,614	166
Sweden	2,704	998	340	3,362	182

Considerations for reducing environmental impacts of milk consumption in Nordic and Baltic countries

- Reducing dairy milk consumption, especially cheese, would provide the greatest environmental benefits across a range of metrics (e.g., agricultural land use, GHGs, nutrient pollution, biodiversity loss). Hence, a reduction in consumption should be considered a first order priority – with greatest scope in countries consuming above levels compliant with planetary boundaries. Currently, dairy cows are used to supply milk during their lives and beef meat at the end of their lives. Therefore, for maximum effect, a reduction in dairy consumption should be paired with a reduction in beef consumption – otherwise beef cattle herds (which have higher environmental impacts per unit of beef compared to beef produced from dairy herds) might be increased to compensate for a reduced beef output from the dairy industry.
- If a replacement of nutrition is required (rather than just a reduction), this should also be considered in terms of opportunities to minimise environmental impacts through replacement with plant-based alternatives where possible.
- Reducing dairy production and consumption would reduce the demand for feed crops (see related considerations in chapters 3 and 5 for cereals and pulses/legumes), hence replacing dairy milk with plant foods may not require an increase in cropland.
- Similarly for meat production and consumption, given the range of factors involved and the complexity of trade-offs and location specific impacts and contexts, a national strategy should usefully direct the change in milk production and consumption. An important part of this would be a comprehensive review to identify and explore the range of issues including biodiversity, human health and social impacts, animal welfare, and current and emerging threats e.g., antibiotics, where necessary, and zoonotic-driven pandemics.
- The approach to reducing dairy production and consumption must be done in conjunction across all livestock production and consumption given the important connections, interdependencies and similarities between meat and dairy, and must take a longer-term perspective rather than being limited to the realities of today's market and other socio-political determinants. Situations can change rapidly and business as usual cannot be assumed under the expected impacts of climate change alone.

NNR Food Group 10: Eggs


Global context

At the global level, 90 mt eggs were produced in 2019 of which 93% were from chickens. China produced 37% of the global total (37 mt in 2019), followed by the US (7 mt), Indonesia (5 mt), India (5 mt), and Brazil (3 mt) – making up the top 5 producers globally.³⁶⁶

The environmental impacts of eggs tend to be largely concentrated during the production stages of the life cycle, which account for around 90% of impacts. For example, GHGs are largely attributed to feed crop production, whereas the animal farming component accounts for the largest share of acidification impacts, and eutrophication impacts are more equally spread between feed crop

production and animal farming (figure 10.1). Table 10.1 shows a range of environmental impacts per kg of egg production, as a global average.

Figure 10.1: Proportion of environmental impacts from different life cycle stages: Eggs (global)³⁶⁷



The types of environmental issues related to egg production are largely the same as those related to meat and dairy production (e.g., land use, manure and urea, nutrient pollution of surrounding ecosystem, resource use on farm including water and energy for housing) (see chapters 8 and 9), which includes a significant proportion of impacts from feed production (see chapters 3 and 5 for more information on the environmental impacts of cereals and legumes). Antibiotic use, pathogen spread, and animal welfare are also relevant to egg production, but are beyond the scope of the current paper.


Table 10.1: Average environmental impacts per kg of retail weight: Eggs (global)³⁶⁸

	Land use (m ² /kg)	Greenhouse gas emissions (kg CO ₂ eq/kg)	Acidifying emissions (g SO ₂ eq/kg)	Eutrophying emissions (g PO ₄ ³⁻ eq/kg)	Freshwater (L/kg)
Eggs	6.3	4.7	53.7	21.8	578

Nordic and Baltic context

Figure 10.2 shows the variation in environmental impacts of egg consumption across the Nordic and Baltic countries and how they relate to global limits. For example, if everyone in the world had the same per person egg consumption as Lithuania, it would use 15% of the global cropland limit for food consumption and 14% of the nitrogen limit for food consumption.

Figure 10.2: Impacts of egg consumption across the Nordic and Baltic countries in relation to the food portion of global environmental limits (%)



Source: *Global Nutrition Report 2021*³⁶⁹ The analysis utilizes country-specific food consumption and environmental footprint data and relates them to the food portions of global environmental limits i.e. a global test to assess the impacts if everyone in the world consumed at the given rate. The methods and data are described in Box 1, and presented in appendices 1-3.

While the environmental impacts of egg consumption might appear insignificant or relatively small compared to other animal products such as meat and dairy, the environmental impacts of egg consumption exceed the impacts of other important food groups in a number of ways within the region. For example, egg consumption in Denmark has a greater environmental impact in terms of GHGs, cropland, water, nitrogen and phosphorus than the consumption of root vegetables. In Estonia, egg consumption requires more cropland than legumes and roots combined. Egg consumption in Sweden requires more nitrogen than fruits and vegetables, and more land than the consumption of legumes and nuts combined. In Lithuania, egg consumption uses more cropland and emits more GHGs than those related to fruit and vegetable consumption. More freshwater is used for egg consumption in Iceland compared to legumes and nuts combined (figures 5.3 and 6.2).

Sweden is the biggest producer of eggs, followed by Denmark and Finland. Denmark, Sweden and Latvia have the largest supply of eggs, followed by Estonia. Finland and Latvia are net exporters of eggs, while Denmark is the largest net importer followed by Estonia (table 10.2). For comparison, egg consumption in line with a flexitarian diet that aligns with planetary boundaries up to 2050 (if implemented in combination with medium-ambition technological measures to reduce inputs during food production, and reductions in food loss and waste of 75%), recommends a daily maximum egg intake of 13g, or 4.7 kg per year (as part of an average daily intake of 2,100 kcal).³⁷⁰ Based on the annual supply data in table 10.2, all countries exceed this amount (on a per person basis), by 2 – 3 times. In contrast, using egg consumption data from food diaries in table 2, Sweden is the only slightly above the flexitarian diet recommendations – consumption in all other countries exceeds this amount.

Table 10.2: Egg production, trade and supply across the Nordic and Baltic countries in 2019³⁷¹

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	100	10	5	85	12
Estonia	150	5	10	135	15
Finland	120	15	20	85	10
Iceland	80	2	1	77	8
Lithuania	110	10	5	95	11
Latvia	130	15	20	95	13
Norway	90	5	10	75	9
Sweden	180	10	15	155	18

Denmark	78	40	23	95	14
Estonia	9	11	2	18	13
Finland	76	3	12	67	11
Iceland	5	0	0	5	12
Latvia	46	16	31	31	14
Lithuania	48	18	21	45	12
Norway	71	1	0	72	12
Sweden	143	18	20	141	14

Considerations for reducing environmental impacts of egg consumption in Nordic and Baltic countries

Egg consumption in all countries exceeds amounts consistent with planetary boundaries. While the environmental impacts of egg consumption might appear insignificant or relatively small compared to other animal products such as meat and dairy, they should be considered within the context of the amount of nutrients provided to diets via egg consumption, and the potential to instead at least partly achieve such nutrition from other less environmentally burdensome sources as a contribution to reducing the overall environmental impacts of nutritionally adequate diets and the food system in line with key environmental targets.

SECTION 3: ADDITIONAL FOOD GROUPS TO CONSIDER

NNR Food Group 1: Breastfeeding

While breastfeeding is not a major area of concern regarding the environmental impacts of food consumption across the Nordic and Baltic countries, there is an emerging literature that demonstrates some considerations in terms of reducing environmental impacts. The majority of breast milk substitutes are based on bovine milk (from cows, which are ruminant animals) and environmental impacts occur largely from this ingredient. For example, breast milk substitutes based on bovine milk assessed in 4 different countries (UK, China, Brazil and Vietnam) all revealed the majority (68% - 82%) of the greenhouse gas (GHGs) impact of production arise from raw milk. The remaining emissions arise from other ingredients including lactose or glucose syrup, vitamins and minerals, and a blend of vegetable oils; and transport, production and in-home sterilisation of bottles, and preparation of breast milk substitutes. In comparison, in all four countries the GHG impact of breastfeeding was 40%, 53%, 43% and 46% lower than breast milk substitutes.³⁷² Similarly, a recent study of Norwegian consumption³⁷³ found that 4 months exclusive feeding with infant formula had an environmental impact of 35–72% higher than 4 months exclusive breastfeeding, depending on the impact category which included global warming potential, terrestrial acidification, marine and freshwater eutrophication, and land use. Bovine milk was the main source of environmental impact across all categories, for the infant formula. The environmental impact of breastfeeding was based on the additional food intake required for breastmilk production in lactating mothers. Additionally, a recent analysis of UK consumption estimated that breastfeeding for six months could save 95-153 kg CO₂ equivalents per baby compared with breast milk substitutes

based on bovine milk – equivalent to GHG emissions from 50,000 and 77,500 cars each year in the UK.³⁷⁴ While it is possible to reduce the environmental impacts of breast milk substitutes, it has been demonstrated that for both Ireland and China, the GHG emissions saved by achieving the minimum 50% Exclusive Breast Feeding target are greater than decarbonising current consumption (including breast milk substitutes) with renewable energy alone.³⁷⁵

Dairy milk production has a substantial environmental burden in terms of GHGs, land use, nitrogen use, phosphorus impacts, freshwater use, and biodiversity loss, and is explored in detail in ‘NRR Food Group 9: Milk and dairy products’ (see chapter 9).

NRR Food Group 2: Drinks

The drinks group includes coffee, tea, sugar sweetened and artificially sweetened drinks. However, sugar is covered in ‘NRR Food Group 12: Sweets and confectioneries’ (see chapter 12), therefore this chapter will focus on tea and coffee, and plant-based drinks.

Global context

Globally, in 2019 10 million tonnes (mt) of coffee and 7 mt of tea were produced. Brazil produced the largest quantity of coffee in 2019 (3 mt), followed by Vietnam (1.6 mt), and Columbia (0.9 mt). tea production is concentrated in east and south Asia, with China producing the largest amount in 2019 (2.8 mt), followed by India (1.4 mt).³⁷⁶

The creation of tea plantations typically displaces native forest and results in biodiversity loss.³⁷⁷

Globally, coffee production was estimated to be the sixth largest driver of deforestation – and thus biodiversity loss - from 2001-2015, contributing a slightly smaller share than cocoa (see chapter 12), and around 24 times less deforestation in comparison to cattle.³⁷⁸

The land use change impacts are reflected in the proportion of environmental impacts across different stages of the life cycle, with more than 10% of the GHGs related to coffee resulting from land use change. Crop production, i.e. on farm, is generally the most important life cycle stage in terms of environmental impacts, however coffee has relatively large impacts associated with losses (in comparison to other food products which typically have less than 20% of life cycle impacts from losses e.g., cereals). The impacts of soy milk tend to be more spread across the stages, with retail having the largest proportion of GHGs and acidification impacts, and crop production causing the largest share of eutrophication impacts (figure 2.1).

Figure 2.1: Proportion of environmental impacts from different life cycle stages: Coffee and soy milk (global)³⁷⁹




Table 2.1 shows a range of impacts related to coffee and soy milk production. Note that the amounts given are per kg of retail weight, and a cup of coffee typically requires 15g of coffee, hence the amounts shown represent coffee for around 67 cups. In that case, 1 cup of coffee would for example result in 0.4 kg CO₂eq, in addition to the resources used for water (e.g., energy for boiling, milk, sweetener, packaging – such as disposable cups or aluminum foil for individual portions for coffee machines). Per cup of tea, the GHG impact has been estimated to be up to 0.2 kg CO₂ depending on how it is grown, processed, shipped, packed, brewed and discarded. Loose leaf tea has a smaller impact (around 10 times less) due a reduced need for packaging. Boiling water is one of the highest energy consuming aspects of tea beverages.³⁸⁰


Table 2.1: Average environmental impacts per kg of retail weight: Coffee and soy milk (global)³⁸¹

	Land use (m ² /kg)	Greenhouse gas emissions (kg CO ₂ eq/kg)	Acidifying emissions (g SO ₂ eq/kg)	Eutrophying emissions (g PO ₄ ³⁻ eq/kg)	Freshwater (L/kg)
Coffee	21.6	28.5	83.1	110.5	26
Soy milk	0.7	1.0	2.6	1.1	28

Figure 2.2 shows a range of plant-based drinks in comparison to soy, with dairy milk included as such drinks tend to be used in the same way, and plant based milk or plant based dairy alternatives are mentioned in many food based dietary guidelines.³⁸² An increased demand for such alternatives is expected over the next decade³⁸³ and lactose malabsorption is highly prevalent in the global population (including 28% of the population in western, southern and northern Europe), making it difficult to digest cow's milk.³⁸⁴ In terms of relative environmental impacts, all plant-based drinks have lower impacts than dairy milk across the range of metrics (figure 2.2). However, the nutritional values can differ significantly between dairy and plant-based in terms of saturated fat and cholesterol, and also calcium and protein depending on whether they are supplemented with nutrients. An analysis of 399 products (semi-skimmed cow's milk and soy-, oat-, almond-, coconut- and rice drink) across 6 European countries (including Sweden) found that 50% of the regular plant-based drinks were fortified with calcium, whereas the organic plant-based drinks were mostly unfortified. Soy drink had the best protein quality to carbon footprint ratio, followed by cow's milk.³⁸⁵ However, it is important to note that fortification of dairy cow feed is common practice, in amounts similar to that of plant-based dairy alternatives (~140 mg of minerals and 1 mg vitamin per 100 g milk).³⁸⁶ The main difference between the fortified plant-based dairy alternatives and dairy

milk is in protein content³⁸⁷, which is unlikely to be a concern in most developed/high income country settings where protein is consumed well above the recommended level.³⁸⁸

Figure 2.2: Environmental impacts of plant-based drinks, per litre.³⁸⁹



Source: Poore, J., & Nemecek, T. (2018). Reducing food's environmental impacts through producers and consumers. Science. OurWorldInData.org/environmental-impacts-of-food • CC BY

Many of the same issues related to soy milk, and tea and coffee production (e.g., land use, land use change, freshwater use, nutrient pollution of surrounding ecosystems, monocultures, intensive farming methods, soil erosion, use of crop protection products, and biodiversity loss), apply to cereal, fruit and legume production and are therefore covered in detail in the relative chapters (see chapters 3, 4 and 5 for more details). An additional issue related to coffee production is a partial proportion (up to 40%) of the yield dependency on pollinators.³⁹⁰

Nordic and Baltic context

All countries are entirely dependent on tea and coffee imports from tropical areas for national consumption (tables 2.2 and 2.3). In all countries, coffee is consumed in much higher quantities in comparison to tea – in some cases substantially more, for example 59 and 41 times more coffee is consumed compared to tea in Finland and Norway respectively. Globally, Finland, Sweden, Iceland, Norway and Denmark have the 3rd, 4th, 5th, 6th and 8th highest coffee consumption respectively.

Table 2.2: Coffee production, trade and supply across the Nordic and Baltic countries in 2019³⁹¹

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Supply (Kg/person/Year)

Denmark	0	50	8	7.3
Estonia	0	8	1	4.7
Finland	0	91	10	11.8
Iceland	0	3	0	8.9
Latvia	0	12	6	2.8
Lithuania	0	24	11	4.9
Norway	0	45	1	8.2
Sweden	0	135	34	10.1

Table 2.3: Tea production, trade and supply across the Nordic and Baltic countries in 2019³⁹²

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Supply (Kg/person/ Year)
Denmark	0	6	1	-	0.7
Estonia	0	0	0	-	0.3
Finland	0	1	0	-	0.2
Iceland	0	0	0	-	0.6
Latvia	0	1	0	-	0.3
Lithuania	0	1	0	-	0.3
Norway	0	1	0	-	0.2
Sweden	0	5	1	-	0.4

In terms of plant-based drink products on the Danish market, recent testing revealed the protein content is generally low, except for soy drinks, and the vitamin and mineral content such as calcium may also be low if not fortified.³⁹³ This difference might be important for consumers with poor dietary diversity, or a reliance on dairy milk for protein intake for example (although this is unlikely in a developed/high income country setting where protein is generally consumed above recommended levels³⁹⁴).

Considerations for reducing environmental impacts of drinks consumption in Nordic and Baltic countries

While tea and coffee consumption is not a major area in terms of potential to reduce environmental impacts from food consumption within the Nordic and Baltic countries, there are some aspects to consider. For example, coffee consumption could be reduced without a need for replacement, to amounts that correlate positively with health. In addition, herbal teas sourced from plants grown within the region (and possibly considered weeds, such as nettles) could be alternatives to imported tea. However, herbal teas do not contain caffeine and therefore are not a functional replacement for caffeinated drinks. Reducing the environmental impacts of other ingredients added to coffee and tea, such as dairy milk, could also be explored – for example, the replacement of dairy milk with fortified plant-based drinks (see chapter 9 for considerations on reducing the impacts of dairy consumption). There are a number of measures to take regarding energy consumption of boiling water, for example, using renewable energy sources and more energy efficient equipment.

NNR Food Group 6: Nuts

Global context

At the global level, 131 mt nuts were produced in 2019 of which 48% were coconuts, and 38% were groundnuts (peanuts). Around 1% of total groundnut production was used for farmed animal feed. Cashews and almonds each accounted for ~3% of global production, and walnuts and chestnuts each accounted for ~2%. Combined, Indonesia, Philippines, and India produced 67% of all coconuts, and China produced the largest quantity of groundnuts (37% of the global total). The US produced the most almonds and India produced the most cashews. China was the main producer of walnuts and chestnuts.³⁹⁵

The environmental impacts of tree nuts and groundnuts tend to be largely concentrated during the production stages of the life cycle, particularly in terms of eutrophication for which crop production accounts for around 80% of impacts. Tree nuts have a large carbon sequestration effect through the process of photosynthesis, resulting in a net negative land use change impact in terms of GHGs, whereas groundnuts have net positive GHG emissions related to land use change for crop production (figure 6.1).

Figure 6.1: Proportion of environmental impacts from different life cycle stages: Nuts (global)³⁹⁶




Table 6.1 shows a range of environmental impacts per kg of groundnut and tree nut production, as a global average. Groundnuts generally have lower impacts compared to tree nuts, except for GHGs. Groundnuts are a leguminous crop and fix nitrogen during production, hence the need for nitrogen fertilizer is reduced. This again highlights the importance of trade-offs regarding environmental impacts – for example, selecting to increase the production of groundnuts rather than tree nuts on the basis of lower land use, acidifying emissions, eutrophying impacts and freshwater use would increase GHGs and result in further land use change for crop production, in turn adversely impacting biodiversity (which is not indicated in table 6.1). In addition to being important for carbon storage, maintaining and increasing biodiversity (depending on pesticide regimes and scale of monoculture), trees also help to regulate weather across large areas and can therefore be important from a micro-climate perspective, e.g., rainfall and floods. Hence, there are important factors to consider beyond the utility of LCA metrics in determining the relative costs and benefits of increasing tree nut and groundnut production.

Table 6.1: Average environmental impacts per kg of retail weight: Nuts (global)³⁹⁷

	Land use (m ² /kg)	Greenhouse gas emissions	Acidifying emissions	Eutrophying emissions	Freshwater (L/kg)
--	----------------------------------	-----------------------------	-------------------------	--------------------------	----------------------


		(kg CO ₂ eq/kg)	(g SO ₂ eq/kg)	(g PO ₄ ³⁻ eq/kg)	
Nuts	13.0	0.4	45.2	19.2	4134
Groundnuts	9.1	3.2	22.6	14.1	1852

Many of the same issues related to nut production (e.g., land use change, nutrient pollution of surrounding ecosystems, monocultures, use of crop protection products, biodiversity loss), apply to cereal and legume production (see chapters 3 and 5 for more details). Additionally for groundnuts, aflatoxins (produced by fungus *Aspergillus flavus*) could be an increasing problem due to more frequent weather extremes (flood and drought).

Nordic and Baltic context

Figure 6.2 shows the variation in environmental impacts of nut and seed consumption across the Nordic and Baltic countries and how they relate to the food portion of global limits (as nuts consumed are imported, the majority of impacts occur outside of the region). GHGs from nut and seed consumption tend to account for the least impact on planetary boundaries, with cropland being the biggest impact, followed by freshwater and nitrogen use (biodiversity impacts are not included in the LCA data presented here). As nut consumption is recommended to increase across the Nordic and Baltic countries, minimizing such impacts per unit of production would be beneficial in order to reduce environmental burdens as consumption rises. Precision agriculture, a method used to apply inputs such as water and fertilizers in a direct rather than diffuse way in appropriate amounts at the correct times, could be important for minimizing nitrogen and water use, and agroforestry could be important in terms of land use as the impacts could be spread across different crops and potential leisure activities. While table 6.1 shows environmental impacts are generally lower for groundnuts in comparison to tree nuts, there is potential to grow tree nuts across the region (and possibly groundnuts in sunnier areas with some protection). Large areas of the region were covered in native Boreal forest, hence there is potential for tree growth in conjunction with other food group changes. For example, reconfiguring portions of permanent or temporary pastureland that displaced forest, with tree nut production – and reconfiguring portions of animal feed crop land with groundnuts. Increasing tree nut production could also make important contributions to climate (in terms of carbon sequestration) and biodiversity goals. Hence, increasing the consumption of nuts should be considered in conjunction with reduced environmental impacts from shifts in other food groups, rather than being considered as an increase in the overall environmental burden of diets.

Figure 6.2: Impacts of nut and seed consumption across the Nordic and Baltic countries in relation to the food portion of global environmental limits (%)



Source: *Global Nutrition Report 2021*³⁹⁸ The analysis utilizes country-specific food consumption and environmental footprint data and relates them to the food portions of global environmental limits i.e. a global test to assess the impacts if everyone in the world consumed at the given rate. The methods and data are described in Box 1, and presented in appendices 1-3.

All countries are currently fully reliant on imports for nut consumption. In all countries, nut consumption is dominated by tree nuts, except for Finland where consumption is equally shared with groundnuts. The lowest proportion of groundnuts in relation to total nut intake is in Estonia, where it accounts for 13%. In Latvia, groundnuts account for 30% of total nut consumption, 32% in Denmark, 33% in Iceland, 38% in both Lithuania and Sweden, and 43% in Norway.³⁹⁹ For comparison, nut and seed consumption in line with a flexitarian diet that aligns with planetary boundaries up to 2050 (if implemented in combination with medium-ambition technological measures to reduce inputs during food production, and reductions in food loss and waste of 75%), recommends a daily minimum nut and seed intake of 50g, or 18.3 kg per year (as part of an average daily intake of 2,100 kcal).⁴⁰⁰ Average supply of tree nuts and groundnuts shown in table 6.2 falls far short in all 8 countries. Based on consumption amounts for nuts in table 2, consumption is below the amounts by at least 6-fold in all countries (there is no data on nut consumption for Lithuania in table 2).

Table 6.2: Tree nut and groundnut production and supply across the Nordic and Baltic countries in 2019⁴⁰¹

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	0	34	6	28	4.2
Estonia	0	5	0	5	3.0
Finland	0	14	0	14	2.3
Iceland	0	1	0	1	2.8
Latvia	0	6	2	4	2.2
Lithuania	0	15	7	8	2.7
Norway	0	21	1	20	3.3

Sweden	0	40	5	35	3.2
--------	---	----	---	----	-----

Considerations for reducing environmental impacts of nut consumption in Nordic and Baltic countries

As a substantial increase in nut consumption would be required across the region to meet current dietary guidelines (table 2), or to align with recommended intakes in the Planetary Health Diet, it is important to consider the environmental impacts in an absolute and relative sense. The main goal is to reduce the relative environmental impacts of nuts per unit of production while also reducing the absolute impacts of diets in terms of total food consumption, by reducing environmental impacts through shifts in other food groups that have higher environmental impacts such as meat and dairy. We identify the following considerations in relation to the environmental impacts of increasing nut consumption:

- In a relative sense, the environmental impacts of nut production per unit of output could be reduced through the use of precision agriculture, polycultures, and organic production methods (or similar), for example to reduce the impacts of crop protection products and provide more opportunities for biodiversity. However, the overall, or absolute impacts of nuts in the diet might still increase due to an increase in consumption.
- Reducing the disproportionately sized environmental impacts of animal products (particularly meat and dairy) will be important in reducing the total, or absolute, environmental impacts of diets, and therefore in creating 'space' for increased nut consumption. The potential to shift some animal product consumption to nuts could also be explored.
- The potential to increase nut production (such as hazelnuts and acorns) within the region could be included in a land use assessment, and also considered in relation to major environmental goals (such as climate - (particularly in relation to carbon sequestration benefits of tree nut production,- and biodiversity) and also in reducing nutrient pollution to ecosystems such as the Baltic sea. Hazelnuts have a legacy in the region and could therefore provide a useful starting point. A land use assessment would bring a comprehensive perspective, which is important for avoiding increasing total environmental loads through increased overall agricultural production, in comparison to reducing overall environmental loads through land use reconfiguration.
- Increasing nut production within the region could help to reduce import dependency, and in turn the potential insecurity of imports due to environmental issues such as increased aflatoxin in production areas more prone to weather extremes (e.g., Africa), and water stress (e.g., China).
- Other options might be useful to explore, for example rather than increasing nut consumption to the full extent in line with dietary guidelines, seeds (e.g., sesame, linseed or hemp), might provide nutritional qualities and have lower environmental impacts.


NNR Food Group 11: Fats and oils

Global context

At the global level, 223 mt of fats and oils were produced in 2019 of which 93% were vegetable oils. Palm oil was produced in the largest quantity (75 mt) - accounting for 33% of total production, followed by soybean oil (60 mt), rapeseed oil (24 mt), sunflower oil (20 mt) and pig fat (11 mt).⁴⁰² Globally, more than 300 million hectares of land is used for oil crop production.⁴⁰³ The majority (68%) of palm oil is used for human food, and 27% is used for industrial applications including soaps. While palm oil is consumed around the world, Indonesia accounts for 57% of annual production, and Malaysia produces 27%. The amount of land used globally for palm oil production has increased from 4 million hectares in 1980 to 19 million hectares in 2018, and is concentrated across the tropics. Most of this land use is in Malaysia and Indonesia (63%). In Indonesia, palm oil plantations are estimated to account for 23% of deforestation over a 15-year period from 2001 - 2016. Globally, from 2001-2015, oil palm was responsible for around 25% more deforestation in comparison to soy, and around 4 times less deforestation in comparison to cattle.⁴⁰⁴

The land use change impacts are reflected in the proportion of environmental impacts across different stages of the life cycle (figure 11.1). For example, palm oil and soybean oil have large GHG emissions associated with land use change. Conversely, olive oil has net negative GHG emissions related to land use change, due to olive trees sequestering carbon during photosynthesis. For olive, rapeseed and sunflower oil, the majority of environmental impacts occur during crop production.

Figure 11.1: Proportion of environmental impacts from different life cycle stages: Oils (global)⁴⁰⁵



None of the oils listed in table 11.1 have the lowest relative impacts across the board. For example, palm oil has the lowest impacts in comparison to the other oils listed, except for acidifying emissions, where soybean oil has a lower impact. This variation across the oils in terms of relative environmental impacts demonstrates the trade-offs involved in assessing environmental sustainability of food and diets. It also omits information that is important when assessing potential shifts, such as growing conditions. For example, palm is a tropical fruit and hence its production location is limited to the tropics.


Table 11.1: Average environmental impacts per kg of retail weight: Oils (global)⁴⁰⁶

	Land use (m ² /kg)	Greenhouse gas emissions	Acidifying emissions	Eutrophying emissions	Freshwater (L/kg)
--	----------------------------------	-----------------------------	-------------------------	--------------------------	----------------------

		(kg CO ₂ eq/kg)	(g SO ₂ eq/kg)	(g PO ₄ ³⁻ eq/kg)	
Soybean Oil	10.5	6.3	15.7	11.7	415
Palm Oil	2.4	7.3	17.5	10.7	6
Sunflower Oil	17.7	3.6	28.0	50.7	1008
Rapeseed Oil	10.6	3.8	28.5	19.2	238
Olive Oil	26.3	5.4	37.6	37.3	2142

Focusing on single metrics, such as land use, when assessing potential to reduce environmental impacts can also be problematic. For example, as demonstrated in figure 11.2, shifting all oil production to palm oil would reduce the overall land requirements for oil production, but as indicated by table 11.1, this shift would in turn increase GHGs. It also fails to account for location specific impacts relating to crop production, for example growing sunflower seeds in a temperature climate in comparison to clearance of carbon and biodiversity rich forest in the tropics to grow palm oil (the area needed for palm oil equates to a 4-fold increase from the current crop area of 19 million ha). Hence, in some cases, a shift to other oils could potentially have a lower overall impact, for example replacing some palm oil in Europe with rapeseed or sunflower oil. However, the substitutability of palm oil depends on its intended use – it is a largely unique oil in terms of its use in food products. A more effective strategy would be to reduce the use of palm oil in diets/food manufacture, and also to consider shifting palm oil used for biofuels, as a way to reduce overall demand more substantially.⁴⁰⁷ Nutritional properties should also be considered, for example the requirement for poly unsaturated fatty acid rather than saturated fat.

Figure 11.2: Area of land needed if each oil type alone was to meet global vegetable oil demand, 2018.⁴⁰⁸



Source: Calculated by Our World in Data based on data from the UN Food and Agriculture Organization (FAO)
OurWorldInData.org/palm-oil • CC BY

Nordic and Baltic context

All countries, with the exception of Estonia which is a net exporter, are dependent on vegetable oil imports to meet national supply. Norway and Sweden have the highest dependency on imports

(table 11.2). A flexitarian diet that aligns with planetary boundaries up to 2050 (if implemented in combination with medium-ambition technological measures to reduce inputs during food production, and reductions in food loss and waste of 75%), recommends a daily maximum vegetable oil intake of 80g, or 29.2 kg per year, in addition to a maximum intake of 2.5 kg of palm oil per year (as part of an average daily intake of 2,100 kcal).⁴⁰⁹ Based on the annual supply data in table 11.2, all countries consume below this amount, from 2 to around 15-fold (on a per person basis). Vegetable oil consumption data from national food diaries (i.e., table 2) is not available for comparison.

Table 11.2: Vegetable oil⁴¹⁰ production, trade and consumption across the Nordic and Baltic countries in 2019⁴¹¹

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	247	724	368	603	2
Estonia	53	34	71	16	13
Finland	70	178	43	205	3
Iceland	2	10	2	10	11
Latvia	68	103	26	145	16
Lithuania	128	239	97	270	10
Norway	96	571	100	567	5
Sweden	139	880	175	844	8

Finland, Norway and Sweden are net importers of animal fats, whereas Denmark, Iceland, and Lithuania are net exporters (table 11.3). Denmark, Finland, Iceland, Latvia, Norway and Sweden all consume a larger amount of animal fats per person in comparison to vegetable oils. Only Estonia has a lower consumption of animal fats in comparison to vegetable oils, on a per person basis (tables 11.2 and 11.3).

Table 11.3: Animal fat⁴¹² production, trade and supply across the Nordic and Baltic countries in 2019⁴¹³

Country	Production (1000 tonnes)	Imports (1000 tonnes)	Exports (1000 tonnes)	Balance (1000 tonnes)	Food supply (Kg/person/year)
Denmark	269	184	301	152	19
Estonia	9	4	5	8	5
Finland	134	69	45	158	15
Iceland	63	8	49	22	18
Latvia	44	10	10	44	22
Lithuania	87	27	76	38	10
Norway	117	194	91	220	11
Sweden	172	45	25	192	17

Considerations for reducing environmental impacts of oil and fat consumption in Nordic and Baltic countries

The main considerations are:

- Consider the overall consumption needed in healthy plant rich diets.
- Shift from animal to (mostly unsaturated) plant-based fats for human consumption.
- Focus consumption on local (e.g., rapeseed, sunflower) over imported palm or soy oil if the environmental impacts are lower and health impacts are equal or improved.

NNR Food Group 12: Sweets and confectioneries


Global context

Due to a lack of environmental impact assessment data on processed food products (which are generically covered in chapter 15), this chapter focuses on some of the key commodity crops that are major constituents of sweets and confectioneries – sugar and cocoa. In addition, a number of important foods relevant to sweets and confectioneries are covered in other chapters, including cereals (chapter 3), dairy (chapter 9), eggs (chapter 10), and oils and fats (chapter 11).

Sugar cane is produced in higher quantities than any other crop. At the global level, 2 billion tonnes of sugar cane, 280 million tonnes of sugar beet and 6 million tonnes of cocoa beans. were produced in 2019. Around 3.5% of sugar cane and 6% of sugar beet was used for farmed animal feed. Sugar cane is generally grown in tropical and subtropical regions. Brazil produced the most sugar cane in 2019 (752 mt), followed by India (405 mt) and Thailand (131 mt). Sugar beet production is concentrated in temperate regions of the northern hemisphere. Russia produced the most sugar beet in 2019 (54 mt), followed by France (38 mt) and Germany (30 mt). Cocoa beans are generally grown in tropical and subtropical regions. Cote d'Ivoire produced the largest quantity of cocoa beans (2 mt), followed by Ghana (0.8 mt) and Indonesia (0.8 mt).⁴¹⁴ Globally, cocoa production was estimated to be the fourth largest driver of deforestation – and thus biodiversity loss - from 2001-2015, contributing a slightly larger share than coffee, and around 20 times less deforestation in comparison to cattle.⁴¹⁵

The land use change impacts are reflected in the proportion of environmental impacts across different stages of the life cycle, with more than half of the GHGs related to dark chocolate resulting from land use change – despite cocoa being a seed from tree fruit. Crop production is most important in terms of eutrophication impacts, however other stages (mainly transport) are also important for GHGs and acidification impacts of beet and cane sugars (figure 12.1).

Figure 12.1: Proportion of environmental impacts from different life cycle stages: Sweets and confectioneries (global)⁴¹⁶



On a per unit basis, beet sugar has lower environmental impacts than cane sugar across the spectrum of metrics shown in table 12.1. Sugar cane is produced from a grass (sugar beet is a root) and generally requires more processing in comparison to sugar beet, although this does not account for the majority of the difference in impacts.

Table 12.1: Average environmental impacts per kg of retail weight: Sweets and confectioneries (global)⁴¹⁷


	Land use (m ² /kg)	Greenhouse gas emissions (kg CO ₂ eq/kg)	Acidifying emissions (g SO ₂ eq/kg)	Eutrophying emissions (g PO ₄ ³⁻ eq/kg)	Freshwater (L/kg)
Cane sugar	2.0	3.2	18.0	16.9	620
Beet sugar	1.8	1.8	12.6	5.4	218
Dark chocolate	69.0	46.7	46.3	87.1	541

Many of the same issues related to sugar and cocoa production (e.g., land use, land use change, freshwater use, nutrient pollution of surrounding ecosystems, monocultures, intensive farming methods, soil erosion, use of crop protection products, and biodiversity loss), apply to cereal and legume production (see chapters 3 and 5 for more details). An additional issue related to cocoa bean production is a very high (>90%) yield dependency on pollinators.⁴¹⁸

Considerations for reducing environmental impacts of sweets and confectioneries consumption in Nordic and Baltic countries

Figure 12.2 shows the variation in terms of the impacts of sugar consumption across the Nordic and Baltic countries and how they relate to the food portion of global environmental limits. For example, if everyone ate the same amount of sugar as Lithuania it would use over 70% of the global freshwater limit for food consumption. The impacts are highest in terms of freshwater use, but nitrogen and phosphorus use are also substantial, and cropland to a lesser extent.

Figure 12.2: Impacts of sugar consumption across the Nordic and Baltic countries in relation to the food portion of global environmental limits (%)



Source: *Global Nutrition Report 2021*⁴¹⁹ The analysis utilizes country-specific food consumption and environmental footprint data and relates them to the food portions of global environmental limits i.e. a global test to assess the impacts if everyone in the world consumed at the given rate. The methods and data are described in Box 1, and presented in appendices 1-3.

Annual supply of sugar per person varies among the countries, being 42 kg in Denmark, 20 kg in Estonia, 33 kg in Finland, 30 kg in Iceland, 26 kg in Latvia, 30 kg in Lithuania, 27 kg in Norway and 33 kg in Sweden in 2019.⁴²⁰ In comparison, a flexitarian diet that aligns with planetary boundaries up to 2050 (if implemented in combination with medium-ambition technological measures to reduce inputs during food production, and reductions in food loss and waste of 75%), recommends a daily maximum sugar intake of 31g, or 11.3 kg per year (as part of an average daily intake of 2,100 kcal).⁴²¹ Based on the annual supply of sugar, all countries consume above this amount, by around 2 to 4-fold (on a per person basis). Sugar consumption data from national food diaries (i.e., table 2) is not available for comparison.

The biggest challenge in relation to sweets and confectioneries is reducing their consumption in line with dietary recommendations. However, there are a number of considerations for reducing environmental impacts:

- Replacing sweets and confectioneries as much as possible with a diversity of nutritious foods (such as nuts, seeds, pulses/legumes, grains and fruits/vegetables) would likely reduce environmental impacts of diets overall, and free up agricultural land and resources that in turn could enable an increased production of such foods while also providing a nutritional and health benefit. Any continued consumption of sweets and confectioneries (i.e., that meet health criteria) should be plant-based (e.g., made without dairy and animal fats where possible to minimise negative environmental impacts) and should diversify ingredients to help promote a diversity of crop production from farming techniques that favour biodiversity and minimise adverse environmental impacts. For example, carob (a nitrogen fixing leguminous crop) could be explored as an alternative to cocoa.
- More understanding of complex food products and supply chains relative to the Nordic and Baltic countries (and their impacts, such as deforestation) would help to shape

recommendations on sweets and confectioneries from an environmental sustainability perspective. Sugar taxes that help reduce imports could provide a useful tool if carefully planned.

NNR Food Group 15: Ultra processed foods


Global context

The risks and benefits of ultra-processed foods (UPF) have, until recently, largely been understudied. The definition of UPF itself can be contested.⁴²² The NOVA classification system has been endorsed by the WHO/FAO and is a widely accepted definition in the scientific literature.⁴²³ NOVA classifies all foods and food products into one of four categories due to the nature and extent of industrial processing to which they have been submitted (unprocessed and minimally processed; processed culinary ingredients; processed foods; and ultra-processed foods).⁴²⁴ In this classification, UPFs are 'ready-to-consume and ready-to-heat formulations, made by combining substances derived from foods with cosmetic additives, typically through a series of industrial processes' and often contain little or no whole foods. Some examples of UPFs are biscuits, confectionery, reconstituted meats, margarine, and many ready-made meals.⁴²⁵

The definition of UPF is still contested in some areas, with a resistance to classifying foods as ultra-processed because they contain certain additives, irrespective of their full nutritional profile or an argument that these foods are discretionary and can still play a role in a balanced diet. Nevertheless, there is growing concern that the average global diet has quickly transitioned to a highly processed form. Since the 1950's, UPFs have been increasingly consumed in high income countries and now contribute to a substantial proportion of energy intake (for example, more than 50% in the USA and UK; and more than 33% in Australia and France).⁴²⁶ From around 1980 onwards, following the globalisation of food systems, consumption of UPFs began to increase in middle-income countries and now accounts for 30%, 29% and 22% of energy intake in Mexico, Chile and Brazil, respectively.⁴²⁷ UPF consumption is also rising in lower income countries within Asia, Africa and Latin America.⁴²⁸

A general observation is that as country income rises, greater volumes and a wider variety of UPFs, are sold. At first high-income groups account for the majority of UPF consumption within a country. Then as a country continues to prosper, and similar to trends in obesity prevalence, low-income groups begin to account for the majority of UPF consumption.⁴²⁹ There are however, large variations within these trends at regional and country levels. Sales volumes are highest in Australasia, North America, Europe and Latin America, yet growing most rapidly in Asia, the Middle East and Africa (figure 15.1). UPF transitions are closely linked with the industrialisation of food systems (the mass production of primary agricultural commodities), technological change (conversion of these commodities into a diverse range of cheaper ingredients available for use in food manufacturing), and globalisation (opening up market opportunities) and diet-related ill-health associated with excessive energy (kcal) consumption or risks associated with non-sugar sweeteners.⁴³⁰

Figure 15.1: Ultra-processed food sales (kg) per capita by region 2006-2019 with projections to 2024.⁴³¹



Currently, there is a paucity of evidence in terms of environmental impacts of UPFs and their role in environmentally sustainable diets.⁴³² One analysis of food consumed in Australia estimated that ‘discretionary foods’, which includes UPFs, accounted for 29.4% of diet related GHGs.⁴³³ An assessment of UK consumption found that UPFs had lower GHGs and were cheaper to buy compared to minimally processed foods, yet there were of a lower nutritional quality.⁴³⁴ An important concern with UPFs is not necessarily processing per se, but their high palatability and ability to encourage overconsumption of energy (kcal) with minimal nutritional benefit. This has direct consequences for the environment. For example, many UPFs contain palm and soy oils, which have substantial negative environmental effects on land use change, such as deforestation (see chapter 11: fats and oils).⁴³⁵

The impacts of UPFs on agrobiodiversity is an emerging area of research. One concern is that UPFs displace the cultivation and consumption of fresh and minimally processed foods, and in turn further the reduction of crop diversity as their ingredients tend to be sourced from a few high-yielding plant species grown as global commodities in monocultures (such as sugar, maize and wheat).⁴³⁶ The homogeneity of agricultural landscapes reduces opportunities for wildlife by limiting the types of habitats and available resources.⁴³⁷ UPFs containing animal products such as meat further increase their environmental impacts due to the feed requirements for farmed animals (particularly if the feed is sourced from large scale monocultures and limited to a few crop types), in addition to the large environmental impacts of animal farming (see chapter 8 for more information on the inefficiency of and environmental impacts of animal feed and farmed animals).⁴³⁸

Nordic and Baltic context

Taking the global context regarding UPFs, the same concerns would apply in the Nordic context. For example, absolute (system level) vs relative environmental impacts of UPFs, and contribution to land

use change and biodiversity loss via imports as consumption rises. Some countries already have a high consumption of UPFs (e.g., Iceland, where UPFs account for 40% of adult energy intake).⁴³⁹

To assess UPFs in relation to the Nordic and Baltic countries it is considered essential to define them adequately and consistently. For example, using the NOVA definition might lead to the assumption that processing is generally a negative aspect of food production, and that all ready meals are nutritionally inadequate (which is not the case). It is also unclear how plant-based meat replacement products relate to UPFs, since some are highly processed, and others are less processed. In some cases, such as in Finland, meat replacements made from local grains and pulses, such as fava beans, peas, and oats have high-quality nutrition profiles in terms of low contents of salt and saturated fat and also have a high-quality amino acid profile. In some cases, processing can also make plant-based ingredients more digestible and thus make their nutrients more bioavailable. Therefore, nutritional adequacy (or inadequacy) may sometimes be problematic to associate with food processing.

Increasing consumption of discretionary/ultra processed foods is mostly considered a health issue in the region, rather than an environmental issue, due to lack of data on environmental impact of these products. Some of the increase in consumption is attributed to a lack of time and knowledge needed for home cooking, while intense marketing and high palatability might also play a role, along with price and availability. Hence, interventions to reduce discretionary/ultra processed food consumption could usefully consider such aspects.⁴⁴⁰

Considerations for reducing environmental impacts of Ultra Processed Foods in Nordic and Baltic countries

The biggest challenge in relation to reducing the environmental impacts of UPFs is reducing their consumption across populations that are increasingly accustomed to their lower price, taste and convenience. However, there are several important considerations for reducing environmental impacts:

- Replacing UPFs, i.e. foods that are discretionary in terms of nutritional requirements, as needed with a diversity of nutritious foods (such as nuts, seeds, pulses/legumes, grains and fruits/vegetables) would likely reduce environmental impacts of diets overall, and a reduced agricultural production of ingredients used in UPFs could allow more cropland to be used for the production of such nutritious foods. Potential barriers to implementation, such as lack of knowledge of ingredients and cooking/preparation methods, should be considered.
- Any consumption of UPFs should not exceed an amount that would result in higher energy, salt, sucrose or saturated fatty acid contents than recommended, should be plant-based and should diversify ingredients to help promote a diversity of crop production from farming techniques that favour biodiversity and minimise adverse environmental impacts.⁴⁴¹
- More understanding of complex food products and supporting supply chains relative to the Nordic and Baltic countries would help to shape recommendations on UPFs from an environmental sustainability perspective.

SECTION 4: Key environmental sustainability considerations for food consumption in the Nordic and Baltic region

This paper's principal conclusion is that, even though environmental impacts differ between different production systems and region-specific circumstances, the primary determinant of the environmental impact of food consumption is not the production system or place, but what is consumed, and its quantity. In short, substantially reducing the consumption of animal sourced foods (particularly meat and dairy) is the key approach to reducing adverse environmental impacts at the system, aggregate or absolute level. Radical changes to current consumption patterns are urgently required to provide the best chance of avoiding the worst impacts of climate change and ecosystem destabilisation. We therefore focus our advice for reducing the environmental impacts of food consumption in the Nordic and Baltic countries on measures that could be implemented in the immediate to short term. We recognise that other options might become relevant in the medium to long term, such as effective methods to reduce methane emissions from farmed animals, and novel food production technologies, such as precision fermentation, cell-cultured foods, controlled-environment farming and algae production that could improve the environmental performance of food systems. However, the required change is large and immediate enough to necessitate change in both food production and consumption. Delaying remediation of today's burgeoning environmental impacts of the food system in the hope they will be resolved by technologies in the future is a very high-risk strategy that should be avoided.

The overarching advice for all countries, in line with the current body of scientific literature, is to shift to more plant-based dietary patterns – and avoid food waste. From this, we can deduce that there is high potential and necessity to shift food consumption across the Nordic and Baltic countries to minimise its environmental impacts. The extent to which this is necessary depends on current consumption patterns. More specifically, we suggest the following as priority interventions for increasing the environmental sustainability of food consumption:

- Reduce meat and dairy consumption substantially and increase the consumption of legumes/pulses, whole grains, vegetables, fruits, nuts and seeds – and explore the potential to increase consumption of wild berries (while potential impacts on wildlife should also be considered) and cultivation of legumes/pulses, vegetables and grains within the region.
- Explore potential seafood shifts from species with higher impacts to species with lower impacts (e.g., seaweed, bivalves), and explore options for direct human consumption of omega 3 fatty acids e.g., microalgae as a replacement for salmon. Due to the potentially large-scale impacts on ecosystems and the largely unknown nature of fish stocks globally, a precautionary approach to the fish and seafood food group is essential – particularly in relation to pursuing an increase in consumption i.e., more information is required before any targeted increase in fish consumption in the Nordic and Baltic countries could be justified from an environmental sustainability perspective, and also from a food security/risk perspective.
- Explore options that support a reduction in consumption of animal products and have potential to increase provision of plant-based foods through feed-to-food shifts. This is relevant to cereals and pulses, as well as nuts, vegetables and fruits.

In addition to the specific suggestions above, we identify a number of overarching, broader points for consideration:

- Identifying suitable options for reducing the environmental impacts of dietary consumption should be in conjunction with a nutritional assessment to ensure that food shifts align with nutritional adequacy and positive health impacts at the dietary level. This is explored in a subsequent paper in this series.⁴⁴²
- Changes to demand *and* supply are needed to align food systems with environmental thresholds. In this context where consumption of fruits and vegetables must increase, shifting production methods could help to further reduce environmental impacts (particularly water and fertilizer use). In addition, fruits and vegetables that require less resources to produce could be prioritised if in alignment with the requirements of a healthy diet.
- The literature suggests that cultivation methods such as those used in organic production (or similar) result in greater biodiversity benefits. At the global level it is only possible to convert agricultural production to such methods in conjunction with substantial shifts in demand to mostly plant-based diets.⁴⁴³
- The total impacts of food consumption must be considered, including imports, and the impacts must include biodiversity. Impacts must be considered in terms of environmental thresholds at the local, regional and global level.
- National strategies that facilitate changes to food consumption and production (including exports) should consider the complexity of trade-offs and location specific impacts and contexts, and implications for trade. An important part of such strategies would be a comprehensive review to identify and explore the range of issues including human health and social impacts, animal welfare, and current and emerging threats e.g., antimicrobial resistance and zoonotic-driven pandemics.
- A national land use assessment could inform optimal land uses for meeting a range of environmental goals, also accounting for the environmental impacts of food imports in producer countries. One important inclusion would be an assessment of pasture land in terms of its current value and necessity for food production and the potential for alternative food production such as agroforestry of tree nuts, or other uses to help meet social and environmental goals (i.e. mitigating and adapting to climate change and restoring biodiversity), such as restoring portions of native ecosystems (such as Boreal forest in countries with some territory in the Boreal forest zone including Norway, Sweden, Finland, Estonia, Lithuania and Latvia).
- The national strategies and land use assessments could usefully apply the 5 key considerations identified in the first paper in this series as consistent framings: 1. Consider the thresholds, 2. Consider the system, 3. Consider the variables, 4. Consider the context, and 5. Consider the spill over.⁴⁴⁴
- The overarching approach to reducing the environmental impacts of food consumption must not be limited to the realities of today's market and other socio-political determinants. The potential impacts and risks of environmental destabilization on food production and supply (including those related to climate change and biodiversity loss, such as temperature extremes, droughts and floods) should also be considered.

- While urgent and fundamental changes to food production and consumption are required to help meet climate change and biodiversity goals⁴⁴⁵, tackling such issues does not remove the need for urgent reform in other sectors, including energy. Instead, transformation of food systems must be incorporated as one part of a comprehensive ‘green transition’ plan that includes all systems.

Acknowledgements

List of experts engaged in the research workshops

We are very grateful to the experts who participated in our workshops during the formation of this paper and to those who also reviewed earlier drafts of the paper. The following experts who participated in the workshops/reviews have consented to being listed here:

Eirin Marie Skjøndal Bar, PhD, Norwegian University of Science and Technology.

Maija Ceruka, Ministry of Health of the Republic of Latvia.

Hanna Eneroth, Swedish Food Agency.

Hanna Karlsson Potter, Department of Energy and Technology; Agricultural technology and systems, Swedish University of Agricultural Sciences.

Helle Margrete Meltzer, former Research Director, Norwegian Institute of Public Health.

Sirli Pehme, sustainability team lead, Civitta Eesti AS.

Ursula Schwab, PhD, Professor (nutrition therapy), Vice head, Institute of Public Health and Clinical Nutrition, University of Eastern Finland.

Inga Thorsdottir, professor, School of Health Sciences/Faculty of Food Science and Nutrition, and Health Science Institute/Unit for Nutrition Research, University of Iceland and Landspítali University Hospital, Reykjavik, Iceland.

The NNR2023 process and role of background papers

NNR2023 was tasked by the Nordic Council of Ministers to integrate sustainability into the food-based dietary guidelines and thereby contribute to their vision of making the Nordic Region the most sustainable region in the world by 2030.

The NNR2023 Committee collaborated with Chatham House in London, UK and a number of other sustainability experts in the development of several background papers on sustainability. These external experts (including lead-authors and co-authors) will not be involved in the final integration of sustainability and the formulation of the food-based dietary guidelines. All external experts are acknowledged in the background papers.

When developing sustainable food-based dietary guidelines, several major reports covering the main dimensions of sustainability, including the environmental, social and economic dimension will be considered, in addition to the background papers developed in the NNR2023 project.

Appendix 1 Overview of global health and environmental targets and their derivation

Global targets	Comment	Implementation
Paris Climate Agreement	The Paris Agreement's long-term goal is to keep the increase in global average temperature to well below 2°C above pre-industrial levels; and to limit the increase to 1.5°C, since this would substantially reduce the risks and effects of climate change. The goal is reflected in SDG 13 and in the planetary boundary for climate change.	The target for agricultural emissions in line with the 2°C target was derived as 4.7 (4.3–5.3) GtCO ₂ -eq (Wollenberg et al, 2016; Springmann et al, 2018). We adjusted this value for the proportion of emissions related specifically to food consumption (92% of emissions of the whole food system, according to Springmann et al, 2018).
Aichi Biodiversity Targets	Target 5: By 2020, the rate of loss of all natural habitats, including forests, is at least halved and where feasible brought close to zero, and degradation and fragmentation is significantly reduced. The target is related to SDG 15 and the planetary boundary for land- system change.	Contribute to target by not increasing pressure to convert natural land into cropland (or pastures), in line with the food-related planetary boundary for land-systems change (Steffen et al, 2015; Springmann et al, 2018). The planetary boundary value was set to the extent of current cropland (+/- 16%). We internally recalculated the value for consistency with the baseline parameters and our focus on food available for consumption (9.9Mkm ² , 8.3–11.5).
SDG target on water withdrawals	SDG 6.4: By 2030, substantially increase water-use efficiency across all sectors and ensure sustainable withdrawals and supply of freshwater to address water scarcity and substantially reduce the number of people suffering from water scarcity. The goal is in line with the planetary boundary for fresh-water use.	Adopt the food-related planetary-boundary target of maintaining environmental flow requirements by limiting agricultural fresh-water use to below 2,000km ³ , with a range of 800–3,350 km ³ (Springmann et al, 2018). We adjusted the value for the proportion of the food system attributed to diets (1,600km ³ , 640–2,600).
SDG target on nutrient pollution	SDG 14.1: By 2025, prevent and significantly reduce marine pollution of all kinds, in particular from land-based activities, including marine debris and nutrient pollution. The goal is in line with the planetary boundary for biogeochemical flows of nitrogen and phosphorus.	Adopt the food-related planetary-boundary target for nitrogen and phosphorus application in line with limiting eutrophication risk (de Vries et al, 2013; Springmann et al, 2018). We recalculated the value for our focus on consumption-related impacts by applying the original risk fractions to estimates of baseline use, which yielded target values of 51TgN (38–83) and 11TgP (5.6–12.9).

Source: SI Table 4. <https://globalnutritionreport.org/reports/2021-global-nutrition-report/appendix-chapter-2-methodology-and-data-sources/>

Appendix 2: Environmental footprints of food commodities (per kg of product), 2010 and 2050.

Food groups	Greenhouse gas emissions (kgCO ₂ eq/kg)	Greenhouse gas emissions (kgCO ₂ eq/kg)	Cropland use (m ² /kg)	Cropland use (m ² /kg)	Fresh-water use (m ³ /kg)	Fresh-water use (m ³ /kg)	Nitrogen use (kgN/t)	Nitrogen use (kgN/t)	Phosphorus use (kgP/t)	Phosphorus use (kgP/t)
	2010	2050	2010	2050	2010	2050	2010	2050	2010	2050
Wheat	0.23	0.21	3.36	2.46	0.49	0.37	28.73	19.78	4.39	2.01
Rice	1.18	0.90	3.51	2.78	1.07	0.89	36.64	25.07	5.20	2.28
Maize	0.19	0.17	1.98	1.40	0.15	0.12	22.77	14.36	3.57	1.55
Other grains	0.29	0.22	6.17	4.43	0.17	0.14	16.39	9.82	2.72	0.97
Roots	0.07	0.06	0.69	0.52	0.04	0.04	3.60	2.07	0.71	0.30
Legumes	0.23	0.19	11.11	6.89	0.94	0.61	0.00	0.00	0.00	0.00
Soybeans	0.12	0.09	3.95	3.14	0.14	0.15	2.75	1.75	5.88	3.17
Nuts & seeds	0.69	0.65	6.39	5.13	0.43	0.33	14.16	10.84	2.10	1.17
Vegetables	0.06	0.07	0.49	0.34	0.09	0.06	9.55	6.32	1.67	0.81
Oil crops	0.70	0.64	3.12	2.37	0.22	0.19	13.33	8.50	2.86	1.32
Fruits (temperate)	0.08	0.08	1.18	0.97	0.33	0.28	12.73	8.57	1.91	0.92
Fruits (tropical)	0.09	0.10	0.94	0.62	0.32	0.23	10.27	6.10	1.58	0.70
Fruits (starchy)	0.11	0.10	0.88	0.59	0.11	0.08	6.15	3.76	1.05	0.48
Sugar	0.19	0.19	1.67	1.35	1.22	0.88	22.34	15.26	3.84	1.86
Palm oil	1.85	2.03	3.10	2.39	0.00	0.00	22.34	16.29	3.57	1.85
Vegetable oil	0.67	0.63	10.31	8.46	0.47	0.45	42.73	28.19	11.47	5.66
Beef	36.78	40.36	4.21	2.78	0.22	0.17	27.29	17.16	5.36	2.29
Lamb	36.73	37.21	6.24	4.48	0.49	0.42	27.52	21.82	4.94	2.47
Pork	3.14	3.25	6.08	4.90	0.35	0.29	51.52	34.19	8.87	4.05
Poultry	1.45	1.39	6.59	5.18	0.40	0.36	50.20	36.00	9.02	4.35
Eggs	1.61	1.48	6.86	5.19	0.44	0.39	51.22	35.09	8.81	4.18
Milk	1.28	1.39	1.34	1.01	0.08	0.08	6.32	4.63	1.58	0.78
Shellfish	0.03	0.04	0.36	0.46	0.03	0.04	2.19	2.39	0.50	0.40
Fish (freshwater)	0.12	0.12	1.51	1.37	0.10	0.10	11.26	8.39	2.37	1.29
Fish (pelagic)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Fish (demersal)	0.01	0.01	0.13	0.20	0.01	0.01	0.75	0.99	0.19	0.18

Notes: Values shown are global averages per kilogram of product. Footprints for animal products represent feed-related impacts, except for greenhouse gas emissions of livestock, which also have a direct component. Footprints for fish and seafood represent feed-related impacts of aquaculture production weighted by total production volumes. The global averages account for expected efficiency improvements, such as improved feed for livestock, and changes in production by 2050, such as increases in extensive beef production in middle-income countries. The analysis is based on country-specific values.

Source: SI Table 3. <https://globalnutritionreport.org/reports/2021-global-nutrition-report/appendix-chapter-2-methodology-and-data-sources/>. Note, future years factor in improvements in technologies and management practices, including reductions in food loss and waste, along a middle-of-the-road socioeconomic development pathway. The environmental footprint data was also used in Springmann, M et al (2020) The healthiness and sustainability of national and global food based dietary guidelines: modelling study. BMJ <https://doi.org/10.1136/bmj.m2322>.

Appendix 3: Food consumption data (g/day) for Nordic and Baltic countries used in environmental analyses

	Beef	Lamb	Poultry	Pork	Eggs	Fish	Milk	Legumes	Fruit	Vegetables	Nuts/seeds	Whole grains	Sugar	Oils
Denmark	55.7	3.6	39.6	36.1	40.8	28.5	702.7	2.9	192.1	210.7	10.5	84.2	85.1	20.0
Estonia	26.0	1.5	35.7	66.4	29.3	18.2	624.8	8.8	134.8	198.5	6.7	27.7	72.9	19.5
Finland	33.7	4.2	32.1	59.5	21.9	43.9	978.8	3.3	161.7	148.2	6.2	80.4	51.3	26.7
Latvia	12.1	1.2	37.5	74.1	35.6	33.2	567.5	0.2	86.0	190.1	7.1	27.5	56.9	33.2
Lithuania	8.4	0.9	42.1	83.0	32.2	53.6	731.5	9.6	84.1	182.4	4.6	25.2	72.9	24.1
Norway	35.0	11.7	30.4	39.9	27.7	66.0	664.4	11.9	232.8	133.9	9.6	85.2	62.0	47.2
Sweden	42.6	5.9	27.9	62.8	30.9	38.0	892.8	5.2	210.5	160.5	14.5	89.3	63.7	44.2

Notes: Baseline food consumption was estimated by adopting estimates of food availability from the FAO's food balance sheets, and adjusting those for the amount of food wasted at the point of consumption. A three-year average of food availability data around the year 2010 (2009, 2010 and 2011) was derived, to be consistent with the data on food waste, and the data on the environmental impacts of food production (appendix 2).

Source: Supplementary data to Springmann, M et al (2020) The healthiness and sustainability of national and global food based dietary guidelines: modelling study. *BMJ* <https://doi.org/10.1136/bmj.m2322>. Benchmark (BMK) mean consumption data for each country and food extracted from tab 'cons_data' in FBDG_data_file.xlsx. Available to download: <https://ora.ox.ac.uk/objects/uuid:bbf756dd-5556-41f6-b969-a9bee18cdb21/files/d9p2909776>

¹ Halpern, BS, Frazier, M, Verstaen, J et al. (14 more authors) (2022) The environmental footprint of global food production. *Nature Sustainability*, 5 (12). pp. 1027-1039. ISSN 2398-9629.

<https://doi.org/10.1038/s41893-022-00965-x>; Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392), pp. 987–92, <https://doi.org/10.1126/science.aag0216>; Benton et al 2021. Food system impacts on biodiversity loss Three levers for food system transformation in support of nature. Chatham House report, February 2021.

² Williams, D. R. et al. (2020), 'Proactive conservation to prevent habitat losses to agricultural expansion', *Nature Sustainability*, 4, pp. 314–22, <https://doi.org/10.1038/s41893-020-00656-5>; Bodirsky, B. L. et al. (2020), 'The ongoing nutrition transition thwarts long-term targets for food security, public health and environmental protection', *Scientific Reports*, 10(1), pp. 1–14, <https://doi.org/10.1038/s41598-020-75213-3>; Clark, M. A. et al. (2020), 'Global food system emissions could preclude achieving the 1.5° and 2°C climate change targets', *Science*, 370(6517), pp. 705–708, <https://doi.org/10.1126/science.aba7357>; M.

Springmann, H. C. J. Godfray, M. Rayner, P. Scarborough, Analysis and valuation of the health and climate change cobenefits of dietary change. *Proc. Natl. Acad. Sci. U.S.A.* 113, 4146–4151 (2016).

doi:10.1073/pnas.1523119113 Medline

³ <https://www.ipcc.ch/report/ar6/syr/resources/spm-headline-statements/>

⁴ Steffen W, Richardson K, Rockström J, Cornell SE, Fetzer I, Bennett EM, Biggs R, Carpenter SR, de Vries W, de Wit CA, Folke C, Gerten D, Heinke J, Mace GM, Persson LM, Ramanathan V, Reyers B, Sörlin S. Sustainability. Planetary boundaries: guiding human development on a changing planet. *Science*. 2015 Feb 13;347(6223):1259855. doi: 10.1126/science.1259855. Epub 2015 Jan 15. PMID: 25592418.

⁵ Steffen, W., K. Richardson, J. Rockström, S.E. Cornell, et al. 2015. [Planetary boundaries: Guiding human development on a changing planet. *Science* 347: 736, 1259855](https://doi.org/10.1038/s41893-022-00287-8); Wang-Erlandsson, L., Tobian, A., van der Ent, R.J. et al. A planetary boundary for green water. *Nat Rev Earth Environ* 3, 380–392 (2022). <https://doi.org/10.1038/s41893-022-00287-8>; Outside the Safe Operating Space of the Planetary Boundary for Novel Entities. Linn Persson, Bethanie M. Carney Almroth, Christopher D. Collins, Sarah Cornell, Cynthia A. de Wit, Miriam L. Diamond, Peter Fantke, Martin Hassellöv, Matthew MacLeod, Morten W. Ryberg, Peter Søgaard Jørgensen, Patricia Villarrubia-Gómez, Zhanyun Wang, and Michael Zwicky Hauschild. *Environmental Science & Technology* 2022 56 (3), 1510-1521. DOI:10.1021/acs.est.1c04158

⁶ Schulte-Uebbing, L.F., Beusen, A.H.W., Bouwman, A.F. et al. From planetary to regional boundaries for agricultural nitrogen pollution. *Nature* 610, 507–512 (2022). <https://doi.org/10.1038/s41586-022-05158-2>

-
- ⁷ Benton, T et al (2023) Forthcoming. Assessing the environmental sustainability of diets – an overview of approaches and identification of 5 key considerations for comprehensive assessments.
- ⁸ Springmann, M., Clark, M., Mason-D’Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>
- ⁹ Springmann, M., Mozaffarian, D., Rosenzweig, C. and Micha, R. (2021), ‘Chapter 02: What we eat matters: Health and environmental impacts of diets worldwide’, in 2021 Global Nutrition Report: The state of global nutrition, Bristol, UK: Development Initiatives, <https://globalnutritionreport.org/reports/2021-global-nutrition-report/health-and-environmental-impacts-of-diets-worldwide/>.
- ¹⁰ Springmann, M., Mozaffarian, D., Rosenzweig, C. and Micha, R. (2021), ‘Chapter 02: What we eat matters: Health and environmental impacts of diets worldwide’, in 2021 Global Nutrition Report: The state of global nutrition, Bristol, UK: Development Initiatives, <https://globalnutritionreport.org/reports/2021-global-nutrition-report/health-and-environmental-impacts-of-diets-worldwide/>.
- ¹¹ Sustainable Development Report 2022. <https://dashboards.sdgindex.org/rankings/spillovers>
- ¹² Based on estimates of food demand from the Food and Agriculture Organization of the United Nations and a database of country and food group-specific environmental footprints. The target values for sustainable food production that would be in line with Sustainable Development Goals were specified by and adapted from the EAT-Lancet Commission. Note: In this test, regional diets in 2010 and 2018 are universally adopted and compared to global environmental targets. Source: Springmann, M., Mozaffarian, D., Rosenzweig, C. and Micha, R. (2021), ‘Chapter 02: What we eat matters: Health and environmental impacts of diets worldwide’, in 2021 Global Nutrition Report: The state of global nutrition, Bristol, UK: Development Initiatives, <https://globalnutritionreport.org/reports/2021-global-nutrition-report/health-and-environmental-impacts-of-diets-worldwide/>.
- ¹³ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives. Notes: Planetary boundaries define the threshold related to global environmental processes beyond which humanity should not go. Planetary boundaries align with the targets for sustainable food production as set out by the Sustainable Development Goals. If globalised impacts exceed 100% of the planetary boundary, the dietary pattern can be considered unsustainable in light of global environmental targets, and disproportionate in the context of an equitable distribution of environmental resources and mitigation efforts. The methods and data are described in Box 1, and presented in appendices 1-2.
- ¹⁴ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives.
- ¹⁵ <https://www.ipcc.ch/report/ar6/syr/resources/spm-headline-statements/>
- ¹⁶ <https://www.ipcc.ch/report/ar6/syr/resources/spm-headline-statements/>
- ¹⁷ Webb, P., Benton, T.G., Beddington, J. et al. The urgency of food system transformation is now irrefutable. *Nat Food* 1, 584–585 (2020). <https://doi.org/10.1038/s43016-020-00161-0>
- ¹⁸ Rockström, J., Gupta, J., Qin, D. et al. Safe and just Earth system boundaries. *Nature* (2023). <https://doi.org/10.1038/s41586-023-06083-8>
- ¹⁹ Karlsson J, Rööös E, Sjunnestrand T, Pira K, Larsson M, Hessellund Andersen B, Sørensen J, Veistola T, Rantakokko J, Manninen S, Brubæk S. Future Nordic Diets Exploring ways for sustainably feeding the Nordics. *TemaNord 2017:566*. <https://norden.diva.portal.org/smash/get/diva2:1163192/FULLTEXT01.pdf>
- ²⁰ Mittenzwei K, van Oort B (2022) Hvordan øke selvforsyningsgraden i norsk jordbruk? *Ruralis note 3/22*
- ²¹ Bakken AK, Mittenzwei K (2023) Produksjonspotensial i jordbruket og nasjonal sjølforsyning med mat. *Utredning for Klimautvalget 2050*. NIBIO, Ruralis
- ²² Future Nordic Diets. Exploring ways for sustainably feeding the Nordics. Johan Karlsson, Elin Rööös, Tove Sjunnestrand, Kajsa Pira, Malin Larsson, Bente Hessellund Andersen, Jacob Sørensen, Tapani Veistola, Jaana Rantakokko, Sirkku Manninen and Stein Brubæk.
- ²³ Source: FAO Statistics Division, http://www.fao.org/es/ess/compendium_2004/technotes.asp
- ²⁴ Definitions and standards - Food Balances (2010-). Glossary. Per capita supply. Available at: <https://www.fao.org/faostat/en/#data/FBS>
- ²⁵ Springmann et al. Options for keeping the food system within environmental limits. *Nature* 2018; 562: 519–25; Poore & Nemecek. Reducing food’s environmental impacts through producers and consumers. *Science* 2018; 360: 987–92; Springmann, M, Spajic, L, Clark, MA, Poore, J, Hertforth, A, Webb, P, Rayner, M, Scarborough, P. The healthiness and sustainability of national and global food-based dietary guidelines: modelling study. *The BMJ*, 2020. <http://dx.doi.org/10.1136/bmj.m2322>; Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives.
- ²⁶ Available to download from: https://globalnutritionreport.org/documents/917/Country-Nutrition-Profiles-data_February_20,23.xlsx (the Global Nutrition Report 2021: Country Nutrition Profiles dataset).
- ²⁷ Available at: https://www.bmj.com/highwire/filestream/1031038/field_highwire_adjunct_files/2/sprm054404.ww3.xlsx
- ²⁸ Poore, J. and Nemecek, T. (2018), ‘Reducing food’s environmental impacts through producers and consumers’, *Science*, 360(6392): pp. 987–92, doi: 10.1126/science.aqa0216.
- ²⁹ CML, “CML2 Baseline Method 2000” (CML, 2001).
- ³⁰ Available to download from: <https://www.science.org/doi/10.1126/science.aqa0216>
- ³¹ Calculated by dividing agricultural land area by total land area for each country, multiplied by 100 (2019 data on land use: <https://www.fao.org/faostat/en/#data/RL>). Land area is defined in the FAOSTAT as: ‘Country area excluding area under inland waters and coastal waters.’
- ³² Land Use 2019. **FAOSTAT**. <https://www.fao.org/faostat/en/#data/RL>. For an overview of agricultural land use types in the FAOSTAT, see figure 1: [Land use statistics and indicators \(fao.org\)](#)

³³ Land used for cultivation of crops. The total of areas under "Arable land" and "Permanent crops". Arable land is the total of areas under temporary crops, temporary meadows and pastures, and land with temporary fallow. Arable land does not include land that is potentially cultivable but is not normally cultivated.

³⁴ Land used permanently (five years or more) to grow herbaceous forage crops through cultivation or naturally (wild prairie or grazing land). Permanent meadows and pastures on which trees and shrubs are grown should be recorded under this heading only if the growing of forage crops is the most important use of the area. Measures may be taken to keep or increase productivity of the land (i.e., use of fertilizers, mowing or systematic grazing by domestic animals.) This class includes: • Grazing in wooded areas (agroforestry areas, for example) • Grazing in shrubby zones (heath, maquis, garigue) • Grassland in the plain or low mountain areas used for grazing: land crossed during transhumance where the animals spend a part of the year (approximately 100 days) without returning to the holding in the evening: mountain and subalpine meadows and similar; and steppes and dry meadows used for pasture.

³⁵ Land cultivated with long-term crops which do not have to be replanted for several years (such as cocoa and coffee), land under trees and shrubs producing flowers (such as roses and jasmine), and nurseries (except those for forest trees, which should be classified under "Forestry"). Permanent meadows and pastures are excluded from land under permanent crops.

³⁶ Land used for crops with a less-than-one-year growing cycle, which must be newly sown or planted for further production after the harvest. Some crops that remain in the field for more than one year may also be considered as temporary crops e.g., asparagus, strawberries, pineapples, bananas and sugar cane. Multiple-cropped areas are counted only once.

³⁷ Land temporarily cultivated with herbaceous forage crops for mowing or pasture. A period of less than five years is used to differentiate between temporary and permanent meadows and pastures.

³⁸ Of which 626,000 ha are cultivated and 102,000 are natural growing.

³⁹ Of which 20,000 ha are cultivated and 159,100 are natural growing.

⁴⁰ [FiBL Statistics - Area data. Organic area \(share of total farmland \[%\]\) in 2019.](#)

⁴¹ [From farm to fork: Our food, our health, our plan \(europa.eu\)](#)

⁴² Herzon, I., K. J. Raatikainen, S. Wehn, S. Rūsiņa, A. Helm, S. A. O. Cousins, and V. Rašomavičius. 2021. Semi-natural habitats in boreal Europe: a rise of a social-ecological research agenda. *Ecology and Society* 26(2):13. <https://doi.org/10.5751/ES-12313-260213>

⁴³ Herzon, I., K. J. Raatikainen, S. Wehn, S. Rūsiņa, A. Helm, S. A. O. Cousins, and V. Rašomavičius. 2021. Semi-natural habitats in boreal Europe: a rise of a social-ecological research agenda. *Ecology and Society* 26(2):13. <https://doi.org/10.5751/ES-12313-260213>

⁴⁴ Warensjö Lemming E., & Pitsi T. (2022). The Nordic Nutrition Recommendations 2022 – food consumption and nutrient intake in the adult population of the Nordic and Baltic countries. *Food & Nutrition Research*, 66. <https://doi.org/10.29219/fnr.v66.8572>

⁴⁵ Warensjö Lemming E., & Pitsi T. (2022). The Nordic Nutrition Recommendations 2022 – food consumption and nutrient intake in the adult population of the Nordic and Baltic countries. *Food & Nutrition Research*, 66. <https://doi.org/10.29219/fnr.v66.8572>

⁴⁶ Nordic Nutrition Recommendations 2012. Integrating nutrition and physical activity. ISBN 978-92-893-2670-4. <http://dx.doi.org/10.6027/Nord2014-002>

⁴⁷ Warensjö Lemming E., & Pitsi T. (2022). The Nordic Nutrition Recommendations 2022 – food consumption and nutrient intake in the adult population of the Nordic and Baltic countries. *Food & Nutrition Research*, 66. <https://doi.org/10.29219/fnr.v66.8572>

⁴⁸ Andersen JLM et al. Intake of whole grain and associations with lifestyle and demographics: a cross-sectional study based on the Danish Diet, Cancer and Health—Next Generations cohort. *European Journal of Nutrition* (2021) 60:883–895 <https://doi.org/10.1007/s00394-020-02289-y>

⁴⁹ Where consumption data is unavailable, the space in the table has been left empty.

References for Table 2:

Denmark: Danskernes kostvaner 2011-2013, 18-75y, n=3016, http://www.food.dtu.dk/-/media/Institutter/Foedevareinstituttet/Publikationer/Pub_2015/Rapport_Danskernes-Kostvaner-2011-2013.ashx?la=da

Estonia: Estonian National Dietary Survey 2014, 18-74Y, n=2713, <https://efsa.onlinelibrary.wiley.com/doi/abs/10.2903/sp.efsa.2017.EN-1198>

Finland: FinDiet 2017, 25-74 y, N=3099, https://www.julkari.fi/bitstream/handle/10024/137433/Raportti_12_2018_nettti%20uusi%202.4.pdf?sequence=1&isAllowed=y

Iceland: Hvatd bordar Íslendingar 2022, 18-80y, N=1312, https://www.landlaeknir.is/servlet/file/store93/item49171/Hvatdbordaislendingar_vefur_endanlegt.pdf

Latvia: Latvian National Dietary Survey (2007-09), N=1377

Latvia: Study of salt and iodine consumption in the adult population in Latvia (2018-2020), 19-64y, N=1011, <https://esparveselibu.lv/petijums/sals-un-joda-paterins-latvijas-pieauguso-iedzivotaju-populacija>

Lithuania: Study of the actual nutrition and nutrition habits of the Lithuanian adult population, (2013-2014), N=2513,

[https://hi.lt/uploads/pdf/visuomenes%20sveikata/2016.01.72/VS%202016%201\(72\)%20ORIG%20Mitybos%20i%20prociai.pdf](https://hi.lt/uploads/pdf/visuomenes%20sveikata/2016.01.72/VS%202016%201(72)%20ORIG%20Mitybos%20i%20prociai.pdf)

Norway: Norkost 3, 2011, 18-70y, N=1787, https://www.helsedirektoratet.no/rapporter/norkost-3-en-landsomfattende-kostholdsundersokelse-blant-menn-og-kvinner-i-norge-i-alderen-18-70-ar-2010-11/Norkost%203%20en%20landsomfattende%20kostholdsundersokelse%20blant%20menn%20og%20kvinner%20i%20Norge%20i%20alderen-18-70%20%C3%A5r%202010-11.pdf/_attachment/inline/b7bafaab-

Sweden: Riksmaten vuxna 2010–2011, 18–80y, N=1797, https://www.livsmedelsverket.se/globalassets/publikationsdatabas/rapporter/2011/riksmaten_2010_2011.pdf?AspxAutoDetectCookieSupport=1

⁵⁰ Nordic Nutrition Recommendations 2012. Integrating nutrition and physical activity. ISBN 978-92-893-2670-4. <http://dx.doi.org/10.6027/Nord2014-002>

⁵¹ <https://globalnutritionreport.org/resources/nutrition-profiles/methodology/> The target values for sustainable food production are in line with the Sustainable Development Goals specified by and adapted from the EAT-Lancet Commission (Willett et al. *The Lancet* 2019; 393: 447–92; Springmann et al. *The British Medical Journal* 2020; 370: 2322).

⁵² National Dietary Guideline year and country used in Figure 3: Denmark, 2015; Estonia, 2015; Latvia, 2008; Lithuania, 2018; Finland, 2014; Sweden, 2015; Iceland, 2014; Norway, 2014. Source data: Springmann, M, Spajic, L, Clark, MA, Poore, J, Hertforth, A, Webb, P, Rayner, M, Scarborough, P. The healthiness and sustainability of national and global food-based dietary guidelines: modelling study. *The BMJ*, 2020. <http://dx.doi.org/10.1136/bmj.m2322>. The methods and data are described in Box 1, and presented in appendices 1–3.

⁵³ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392), pp. 987–92, <https://doi.org/10.1126/science.aag0216>; Springmann, M., Clark, M., Mason-D'Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>; Clark et al 2020; Springmann, M, Spajic, L, Clark, MA, Poore, J, Hertforth, A, Webb, P, Rayner, M, Scarborough, P. The healthiness and sustainability of national and global food-based dietary guidelines: modelling study. *The BMJ*, 2020.

<http://dx.doi.org/10.1136/bmj.m2322>; Hayek, M.N., Harwatt, H., Ripple, W.J., and Mueller, N (2020). The carbon opportunity cost of animal-sourced food production on land. *Nature Sustainability*.

<https://doi.org/10.1038/s41893-020-00603-4>; Erb, KH., Lauk, C., Kastner, T. et al. Exploring the biophysical option space for feeding the world without deforestation. *Nat Commun* 7, 11382 (2016).

<https://doi.org/10.1038/ncomms11382>; Gerald Kalt, Andreas Mayer, Helmut Haberl, Lisa Kaufmann, Christian Lauk, Sarah Matej, Elin Rööös, Michaela C. Theurl, Karl-Heinz Erb, Exploring the option space for land system futures at regional to global scales: The diagnostic agro-food, land use and greenhouse gas emission model BioBaM-GHG 2.0, *Ecological Modelling*, Volume 459, 2021, 109729,

<https://doi.org/10.1016/j.ecolmodel.2021.109729>; David Bryngelsson, Stefan Wirsenius, Fredrik Hedenus, Ulf Sonesson, How can the EU climate targets be met? A combined analysis of technological and demand-side changes in food and agriculture, *Food Policy*, Volume 59, 2016, Pages 152-164, <https://doi.org/10.1016/j.foodpol.2015.12.012>; Moberg et al (2020) Benchmarking the Swedish Diet Relative to Global and National Environmental Targets—Identification of Indicator Limitations and Data Gaps. *Sustainability* 2020, 12, 1407; doi:10.3390/su12041407; Bajzelj, B. et al. Importance of food-demand management for climate mitigation. *Nat. Clim. Change* 4, 924–929 (2014); Tilman, D. & Clark, M. Global diets link environmental sustainability and human health. *Nature* 515, 518–522 (2014); S. Clune, E. Crossin, K. Verghese, Systematic review of greenhouse gas emissions for different fresh food categories. *J. Clean. Prod.* 140, 766–783 (2017); Clark, M et al (2022) Estimating the environmental impacts of 57,000 food products. *PNAS* 2022 Vol. 119 No. 33, <https://doi.org/10.1073/pnas.2120584119>; Intergovernmental Panel on Climate Change (2019), 'Climate Change and Land: An IPCC Special Report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems', <https://www.ipcc.ch/srccl>.

⁵⁴ For example: E. Hallström, A. Carlsson-Kanyama, P. Börjesson, Environmental impact of dietary change: a systematic review, *Journal of Cleaner Production*, Volume 91, 2015, Pages 1-11,

<https://doi.org/10.1016/j.jclepro.2014.12.008>; and Aleksandrowicz L, Green R, Joy EJM, Smith P, Haines A (2016) The Impacts of Dietary Change on Greenhouse Gas Emissions, Land Use, Water Use, and Health: A Systematic Review. *PLoS ONE* 11(11): e0165797. doi:10.1371/journal.pone.0165797; and Jarmul S, Dangour AD, Green R, Liew Z, Haines A, Scheelbeek PF. Climate change mitigation through dietary change: a systematic review of empirical and modelling studies on the environmental footprints and health effects of 'sustainable diets'. *Environ Res Lett.* 2020;15:123014. Published 2020 Dec 22. doi:10.1088/1748-9326/abc2f7

⁵⁵ For example: Wallén, A., Brandt, N., & Wennersten, R. (2004). Does the Swedish consumer's choice of food influence greenhouse gas emissions? *Environmental Science & Policy*, 7, 525-535; Martin M, Brandão M. Evaluating the Environmental Consequences of Swedish Food Consumption and MDietary Choices. *Sustainability* 2017;9:2227. <https://doi.org/10.3390/su9122227>; Saari M, Kaljonen M, Niemi J, Antikainen R, Hakala K, Hartikainen H, Heikkinen J, Joensuu K, Lehtonen H, Mattila T, Nisonen S, Ketoja E, Knuutila M, Regina K, Rikkonen P, Seppälä J, Varho V. Effects of dietary change and policy mix supporting the change, End report of the FoodMin project. (in Finnish with an English abstract). Publications of the Government's analysis, assessment, and research activities 2019:47; Rööös E, Karlsson H, Witthöft C, Sundberg C. Evaluating the sustainability of diets—combining environmental and nutritional aspects. *Environmental Science & Policy* 2015;47:157-166.

⁵⁶ Wood A, Gordon LJ, Rööös E, Karlsson JO, Häyhä T, Bignet V, Rydenstam T, Hård af Segerstad L, Bruckner M. Nordic Food Systems for Improved Health and Sustainability; Stockholm Resilience Centre, University of Stockholm: Stockholm, Sweden, 2019.

⁵⁷ Benton, T et al (2023) *Forthcoming*. Assessing the environmental sustainability of diets – an overview of approaches and identification of 5 key considerations for comprehensive assessments.

⁵⁸ Trolle et al 2023 Integrating environmental sustainability into Food-Based Dietary Guidelines in the Nordic countries (in press).

⁵⁹ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 *Environ. Res. Lett.* 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>

⁶⁰ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392), pp. 987–92, <https://doi.org/10.1126/science.aag0216>.

⁶¹ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 *Environ. Res. Lett.* 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>

-
- ⁶² Halpern, B.S., Frazier, M., Verstaen, J. et al. The environmental footprint of global food production. *Nat Sustain* 5, 1027–1039 (2022). <https://doi.org/10.1038/s41893-022-00965-x>
- ⁶³ Järviö, N., Maljanen, N.-L., Kobayashi, Y., Ryyänen, T. & Tuomisto, H.L. 2021. An attributional life cycle assessment of microbial protein production: a case study on using hydrogen-oxidizing bacteria. *Science of the Total Environment*. 776, 12 p. doi: 10.1016/j.scitotenv.2021.145764
- ⁶⁴ Behm, K., Nappa, M., Aro, N., Welman, A., Ledgard, S., Suomalainen, M., Hill, J., 2022. Comparison of carbon footprint and water scarcity footprint of milk protein produced by cellular agriculture and the dairy industry. *The International Journal of Life Cycle Assessment*, 1-18.
- ⁶⁵ Järviö, N., Parviainen, T., Maljanen, N.-L., Kobayashi, Y., Kujanpää, L., Ercili-Cura, D., Landowski, C.P., Ryyänen, T., Nordlund, E., Tuomisto, H.L., 2021. Ovalbumin production using *Trichoderma reesei* culture and low-carbon energy could mitigate the environmental impacts of chicken-egg-derived ovalbumin. *Nature Food* 2, 1005-1013.
- ⁶⁶ Amiri-Jami, M., Abdelhamid, A.G., Hazaa, M., Kakuda, Y., Griffiths, M.W., 2015. Recombinant production of omega-3 fatty acids by probiotic *Escherichia coli* Nissle 1917. *FEMS microbiology letters* 362.
- ⁶⁷ Springmann, M., Clark, M., Mason-D’Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>; Clark, M. A. et al. (2020), ‘Global food system emissions could preclude achieving the 1.5° and 2°C climate change targets’, *Science*, 370(6517), pp. 705–708, <https://doi.org/10.1126/science.aba7357>.
- ⁶⁸ Benton, T. G. and Harwatt, H. (2022), *Sustainable agriculture and food systems: Comparing contrasting and contested versions*, Research Paper, London: Royal Institute of International Affairs, <https://doi.org/10.55317/9781784135263>
- ⁶⁹ Francaviglia, Rosa, Jorge Álvaro-Fuentes, Claudia Di Bene, Lingtong Gai, Kristiina Regina, and Eila Turtola. 2020. Diversification and Management Practices in Selected European Regions. A Data Analysis of Arable Crops Production. Vol. 10. doi:10.3390/agronomy10020297.
- ⁷⁰ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 *Environ. Res. Lett.* 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ⁷¹ Clark, M. A. et al. (2020), ‘Global food system emissions could preclude achieving the 1.5° and 2°C climate change targets’, *Science*, 370(6517), pp. 705–708, <https://doi.org/10.1126/science.aba7357>.
- ⁷² Francaviglia, Rosa, Jorge Álvaro-Fuentes, Claudia Di Bene, Lingtong Gai, Kristiina Regina, and Eila Turtola. 2020. *Diversification and Management Practices in Selected European Regions. A Data Analysis of Arable Crops Production*. Vol. 10. doi:10.3390/agronomy10020297; Ferreira Helena, Pinto Elisabete, Vasconcelos Marta W. Legumes as a Cornerstone of the Transition Toward More Sustainable Agri-Food Systems and Diets in Europe. *Frontiers in Sustainable Food Systems*. VOLUME 5. 2021. DOI=10.3389/fsufs.2021.694121
- ⁷³ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 *Environ. Res. Lett.* 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ⁷⁴ Tuck, S.L., C. Winqvist, F. Mota, J. Ahnström, L.A. Turnbull & J. Bengtsson. 2014. Land-use intensity and the effects of organic farming on biodiversity: a hierarchical meta-analysis. *Journal of Applied Ecology* 51: 746-755; Muller et al. (2017), ‘Strategies for feeding the world more sustainably with organic agriculture’; Seufert et al. (2018), ‘Current and potential contributions of organic agriculture to diversification of the food production system’; H.L. Tuomisto, I.D. Hodge, P. Riordan, D.W. Macdonald, Does organic farming reduce environmental impacts? – A meta-analysis of European research, *Journal of Environmental Management*, Volume 112, 2012, Pages 309-320, <https://doi.org/10.1016/j.jenvman.2012.08.018>.
- ⁷⁵ Benton, T. G., Vickery, J. A., & Wilson, J. D. (2003). Farmland biodiversity: is habitat heterogeneity the key?. *Trends in ecology & evolution*, 18(4), 182-188.
- ⁷⁶ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 *Environ. Res. Lett.* 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ⁷⁷ Seufert, V., Ramankutty, N. & Foley, J. Comparing the yields of organic and conventional agriculture. *Nature* 485, 229–232 (2012). <https://doi.org/10.1038/nature11069>
- ⁷⁸ Muller, A., Schader, C., El-Hage Scialabba, N., Brüggemann, J., Isensee, A., Erb, K.-H., Smith, P., Klocke, P., Liebe, F., Stolze, M. and Niggli, U. (2017), ‘Strategies for feeding the world more sustainably with organic agriculture’, *Nature Communications*, 8(1290), doi: 10.1038/s41467-017-01410-w
- ⁷⁹ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 *Environ. Res. Lett.* 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ⁸⁰ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 *Environ. Res. Lett.* 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ⁸¹ [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](https://www.livsmedelsverket.se)

-
- ⁸² Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 Environ. Res. Lett. 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ⁸³ [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ⁸⁴ [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ⁸⁵ Varis, A.-L. (1991). Effect of intercropping carrots and onions on damage caused by the carrot fly, *Psila rosae* (F.) (Dipt., Psilidae). *Agricultural and Food Science*, 63(5), 411–414. <https://doi.org/10.23986/afsci.72420>; Lizarazo, Clara I., Antti Tuulos, Venla Jokela, and Pirjo S. A. Mäkelä. 2020. "Sustainable Mixed Cropping Systems for the Boreal-Nemoral Region." *Frontiers in Sustainable Food Systems* 4. doi:10.3389/fsufs.2020.00103.
- ⁸⁶ Peltoniemi, Krista, Sannakajsa Velmala, Hannu Fritze, Riitta Lemola, and Taina Pennanen. 2021. "Long-Term Impacts of Organic and Conventional Farming on the Soil Microbiome in Boreal Arable Soil." *European Journal of Soil Biology* 104: 103314. doi:<https://doi.org/10.1016/j.ejsobi.2021.103314>.
- ⁸⁷ Muller, A., Schader, C., El-Hage Scialabba, N., Brüggemann, J., Isensee, A., Erb, K.-H., Smith, P., Klocke, P., Liebe, F., Stolze, M. and Niggli, U. (2017), 'Strategies for feeding the world more sustainably with organic agriculture', *Nature Communications*, 8(1290), doi:10.1038/s41467-017-01410-w; Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 Environ. Res. Lett. 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ⁸⁸ Benton, T. G. and Harwatt, H. (2022), Sustainable agriculture and food systems: Comparing contrasting and contested versions, Research Paper, London: Royal Institute of International Affairs, <https://doi.org/10.55317/9781784135263>; Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 Environ. Res. Lett. 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ⁸⁹ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392), pp. 987–92, <https://doi.org/10.1126/science.aag0216>.
- ⁹⁰ Halpern, B.S., Frazier, M., Verstaen, J. et al. The environmental footprint of global food production. *Nat Sustain* 5, 1027–1039 (2022). <https://doi.org/10.1038/s41893-022-00965-x>
- ⁹¹ Halpern, B.S., Frazier, M., Verstaen, J. et al. The environmental footprint of global food production. *Nat Sustain* 5, 1027–1039 (2022). <https://doi.org/10.1038/s41893-022-00965-x>
- ⁹² Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 Environ. Res. Lett. 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ⁹³ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 Environ. Res. Lett. 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ⁹⁴ Gabriel, D., Sait, S. M., Hodgson, J. A., Schmutz, U., Kunin, W. E., & Benton, T. G. (2010). Scale matters: the impact of organic farming on biodiversity at different spatial scales. *Ecology letters*, 13(7), 858-869.
- ⁹⁵ Benton, T. G., Bieg, C., Harwatt, H., Pudasaini, R., & Wellesley, L. (2021). Food system impacts on biodiversity loss. *Three levers for food system transformation in support of nature*; and Benton, T. G., and Harwatt, H. "Sustainable agriculture and food systems." (2022). *Chatham House, London*.
- ⁹⁶ Benton, T. G., Bieg, C., Harwatt, H., Pudasaini, R., & Wellesley, L. (2021). Food system impacts on biodiversity loss. *Three levers for food system transformation in support of nature*; and Benton, T. G., and Harwatt, H. "Sustainable agriculture and food systems." (2022). *Chatham House, London*.
- ⁹⁷ Benton, Tim G., Juliet A. Vickery and Jeremy D. Wilson. "Farmland biodiversity: is habitat heterogeneity the key?." *Trends in ecology & evolution* 18.4 (2003): 182-188.
- ⁹⁸ [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ⁹⁹ Stanislas Rigal et al, Farmland practices are driving bird population decline across Europe, Proceedings of the National Academy of Sciences (2023). DOI: 10.1073/pnas.2216573120
- ¹⁰⁰ Benton, T. G., Bieg, C., Harwatt, H., Pudasaini, R., & Wellesley, L. (2021). Food system impacts on biodiversity loss. *Three levers for food system transformation in support of nature*; and Benton, Tim G., and Helen Harwatt. "Sustainable agriculture and food systems." (2022). *Chatham House, London*.
- ¹⁰¹ Outhwaite, Charlotte L., Peter McCann, and Tim Newbold. "Agriculture and climate change are reshaping insect biodiversity worldwide." *Nature* 605.7908 (2022): 97-102.
- ¹⁰² Benton, T. G., Bieg, C., Harwatt, H., Pudasaini, R., & Wellesley, L. (2021). Food system impacts on biodiversity loss. *Three levers for food system transformation in support of nature*; and Benton, Tim G., and Helen Harwatt. "Sustainable agriculture and food systems." (2022). *Chatham House, London*; Harwatt, H., Wetterberg, K., Giritharan, A. and Benton, T. G. (2022), Aligning food systems with climate and biodiversity targets: Assessing the suitability of policy action over the next decade, Research Paper, London: Royal Institute of International Affairs, <https://doi.org/10.55317/9781784135416>; Williams, D. R. et al. (2020), 'Proactive conservation to prevent habitat losses to agricultural expansion', *Nature Sustainability*, 4, pp. 314–22, <https://doi.org/10.1038/s41893-020-00656-5>.
- ¹⁰³ The Kunming-Montreal Global Biodiversity Framework: <https://www.cbd.int/doc/decisions/cop-15/cop-15-dec-04-en.pdf>
- ¹⁰⁴ Production quantity data from FAOSTAT (2020) <https://www.fao.org/faostat/en/#data/QCL>

-
- ¹⁰⁵ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216; FAOSTAT food balance 2019 data: FAOSTAT
- ¹⁰⁶ Source: FAOSTAT food balance 2019 data: [FAOSTAT](#)
- ¹⁰⁷ Grains are an average of wheat, maize, oats, and flooded rice. From figure caption for S13 in Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ¹⁰⁸ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ¹⁰⁹ Jagdish K. Ladha, Mark B. Peoples, Pallavolu M. Reddy, et al. Biological nitrogen fixation and prospects for ecological intensification in cereal-based cropping systems. *Field Crops Research*, V 283, 2022, <https://doi.org/10.1016/j.fcr.2022.108541>.
- ¹¹⁰ Lassaletta, L., Billen, G., Grizzetti, B., Anglade, J., & Garnier, J. (2014). 50 year trends in nitrogen use efficiency of world cropping systems: the relationship between yield and nitrogen input to cropland. *Environmental Research Letters*, 9(10), 105011.
- ¹¹¹ West, P. C., Gerber, J. S., Engstrom, P. M., Mueller, N. D., et al. (2014). Leverage points for improving global food security and the environment. *Science*, 345(6194), 325-328.
- ¹¹² [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#); Breiling, M. Hashimoto, S. Sato, Y. & Ahamer, G. 2005. Rice-related greenhouse gases in Japan, variations in scale and time and significance for the Kyoto Protocol. *Paddy and Water Environment* 3(1), 39-46.
- ¹¹³ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ¹¹⁴ [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ¹¹⁵ Francaviglia, Rosa, Jorge Alvaro-Fuentes, Claudia Di Bene, Lingtong Gai, Kristiina Regina, and Eila Turtola. 2020. *Diversification and Management Practices in Selected European Regions. A Data Analysis of Arable Crops Production*. Vol. 10. doi:10.3390/agronomy10020297; Ferreira Helena, Pinto Elisabete, Vasconcelos Marta W. Legumes as a Cornerstone of the Transition Toward More Sustainable Agri-Food Systems and Diets in Europe. *Frontiers in Sustainable Food Systems*. VOLUME 5. 2021. DOI=10.3389/fsufs.2021.694121
- ¹¹⁶ Jagdish K. Ladha, Mark B. Peoples, Pallavolu M. Reddy, et al. Biological nitrogen fixation and prospects for ecological intensification in cereal-based cropping systems. *Field Crops Research*, V 283, 2022, <https://doi.org/10.1016/j.fcr.2022.108541>.
- ¹¹⁷ Eshel, G., Shepon, A., Makov, T., & Milo, R. (2014). Land, irrigation water, greenhouse gas, and reactive nitrogen burdens of meat, eggs, and dairy production in the United States. *Proceedings of the National Academy of Sciences*, 111, 11996–12001.
- ¹¹⁸ Alexander, P. et al. (2016), 'Human appropriation of land for food: The role of diet', *Global Environmental Change*, 41, pp. 88–98, <http://www.sciencedirect.com/science/article/pii/S0959378016302370?via%3Dihub#bib0330>
- ¹¹⁹ Cassidy, E., West, P., Gerber, J., & Foley, J. (2013). Redefining agricultural yields: From tonnes to people nourished per hectare. *Environmental Research Letters*, 8, 034015.
- ¹²⁰ Harwatt, H (2018) Including animal to plant protein shifts in climate change mitigation policy: a proposed three-step strategy. *Climate Policy* 19(5):1-9 DOI:10.1080/14693062.2018.1528965
- ¹²¹ West, Paul C., et al. "Leverage points for improving global food security and the environment." *Science* 345.6194 (2014): 325-328; Cassidy, E., West, P., Gerber, J., & Foley, J. (2013). Redefining agricultural yields: From tonnes to people nourished per hectare. *Environmental Research Letters*, 8, 034015.
- ¹²² Shepon, A., Eshel, G., Noor, E., & Milo, R. (2018). The opportunity cost of animal based diets exceeds all food losses. *Proceedings of the National Academy of Sciences of the United States of America*, 115, 3804–3809.
- ¹²³ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives.
- ¹²⁴ Production quantity data from FAOSTAT (2020) <https://www.fao.org/faostat/en/#data/QCL>
- ¹²⁵ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ¹²⁶ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ¹²⁷ Percentage of total national supply as shown in table 3.2 under 'balance'.
- ¹²⁸ Calculated from total feed amount in 2019 divided by total food amount in 2019, using FBS data: <https://www.fao.org/faostat/en/#data/FBS>.
- ¹²⁹ Food supply quantity data for 2019 from FBS: <https://www.fao.org/faostat/en/#data/FBS>
- ¹³⁰ [Ellesmere farm produces UK's first large-scale quinoa crop - BBC News](#)

-
- ¹³¹ NNR2012: <https://www.norden.org/en/publication/nordic-nutrition-recommendations-2012>; Aune D, Keum N, Giovannucci E, et al. Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: systematic review and dose-response meta-analysis of prospective studies. *BMJ*. 2016;353:i2716; Nordic Council of Ministers Nordic Nutrition Recommendations 2012: integrating nutrition and physical activity. Nordic Nutrition Recommendations. Report No: 5. Copenhagen: Nordic Council of Ministers; 2014, p. 627. doi: 10.6027/Nord2014-002
- ¹³² Karlsson J, Rööös E, Sjunnestrand T, Pira K, Larsson M, Hesselund Andersen B, Sørensen J, Veistola T, Rantakokko J, Manninen S, Brubæk S. Future Nordic Diets Exploring ways for sustainably feeding the Nordics. *TemaNord* 2017:566.
- ¹³³ Source: FAOSTAT production data, world total 2019 [FAOSTAT](#)
- ¹³⁴ Source: FAOSTAT feed data, world total 2019 [FAOSTAT](#)
- ¹³⁵ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216. Note – tomatoes are fruits but are grouped here with vegetables as they are most commonly consumed as a vegetable.
- ¹³⁶ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ¹³⁷ [Towards Environmentally Sound Dietary Guidelines – Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ¹³⁸ [Towards Environmentally Sound Dietary Guidelines – Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#). Section 3.4: A Non-Toxic Environment.
- ¹³⁹ [Towards Environmentally Sound Dietary Guidelines – Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ¹⁴⁰ [Towards Environmentally Sound Dietary Guidelines – Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ¹⁴¹ [Towards Environmentally Sound Dietary Guidelines – Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ¹⁴² Davis J, Wallman M, Sund V, Emanuelsson A, Cederberg C, Sonesson U. Emissions of Greenhouse Gases from Production of Horticultural Products Analysis of 17 products cultivated in Sweden. SIK 2011:SR 828. <https://www.diva-portal.org/smash/get/diva2:943913/FULLTEXT01.pdf>
- ¹⁴³ van Lenteren, Joop & Alomar, Oscar & Ravensberg, Willem & Urbaneja, Alberto. (2020). Biological Control Agents for Control of Pests in Greenhouses. 10.1007/978-3-030-22304-5_14
- ¹⁴⁴ Carlsson-Kanyama, A. & Boström-Carlsson, K. 2001. Energy use for cooking and other stages in the life cycle of food. A study of wheat, spaghetti, pasta, barley, rice, potatoes, couscous and mashed potatoes. Stockholms universitet och FOI. Forskningsgruppen för miljöstrategiska studier. Fms-rapport 160.
- ¹⁴⁵ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives. Chart does not include root vegetables.
- ¹⁴⁶ <https://www.ri.se/en/barometer-2021-what-happens-to-the-wild-berries>
- ¹⁴⁷ Mittenzwei K, Milford AB, Grønlund A (2017) Status og potensial for økt produksjon og forbruk av vegetabiliske matvarer i Norge
- ¹⁴⁸ van Lenteren, Joop & Alomar, Oscar & Ravensberg, Willem & Urbaneja, Alberto. (2020). Biological Control Agents for Control of Pests in Greenhouses. 10.1007/978-3-030-22304-5_14; <https://www.hortidaily.com/article/9498631/use-of-biological-control-still-increasing-in-dutch-greenhouse/>
- ¹⁴⁹ Naseer M, Persson T, Hjelkrem A-GR, Ruoff P, Verheul MJ (2022a) Life cycle assessment of tomato production for different production strategies in Norway. *Journal of Cleaner Production* 372:133659. <https://doi.org/10.1016/j.jclepro.2022.133659>; Naseer M, Persson T, Righini I, Stanghellini C, Maessen H, Ruoff P, Verheul MJ (2022b) Bioeconomic evaluation of extended season and year-round tomato production in Norway using supplemental light. *Agricultural Systems* 198:103391. <https://doi.org/10.1016/j.agsy.2022.103391>; Naseer M, Persson T, Righini I, Stanghellini C, Maessen H, Verheul MJ (2021) Bio-economic evaluation of greenhouse designs for seasonal tomato production in Norway. *Biosystems Engineering* 212:413–430. <https://doi.org/10.1016/j.biosystemseng.2021.11.005>; Wacker J-D, Verheul MJ, Righini I, Maessen H, Stanghellini C (2022) Optimisation of supplemental light systems in Norwegian tomato greenhouses - A simulation study. *Biosystems Engineering* 215:129–142. <https://doi.org/10.1016/j.biosystemseng.2021.12.020>; SMARTGREEN pioneers zero emission greenhouse | Interreg North Sea; Muhammad Naseer, Tomas Persson, Isabela Righini, Cecilia Stanghellini, Henk Maessen, Peter Ruoff, Michel J. Verheul. Bioeconomic evaluation of extended season and year-round tomato production in Norway using supplemental light. *Agricultural Systems*, Volume 198, 2022. <https://doi.org/10.1016/j.agsy.2022.103391>.
- ¹⁵⁰ <https://sciendo.com/article/10.2478/euco-2018-0039>

-
- ¹⁵¹ Björklund, J., Eksvärd, K. & Schaffer, C. Exploring the potential of edible forest gardens: experiences from a participatory action research project in Sweden. *Agroforest Syst* 93, 1107–1118 (2019). <https://doi.org/10.1007/s10457-018-0208-8>
- ¹⁵² Farstad FM, Hermansen EAT, Grasbekk BS, Brudevoll K, van Oort B (2022) Explaining radical policy change: Norwegian climate policy and the ban on cultivating peatlands. *Global Environmental Change* 74:102517. <https://doi.org/10.1016/j.gloenvcha.2022.102517>
- ¹⁵³ Please note: This information was provided by regional experts during a series of workshops and has not been substantiated by documented evidence.
- ¹⁵⁴ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ⁹⁵ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ¹⁵⁶ Source: FAOSTAT food balances available at [FAOSTAT](#). This figure does not include starchy root vegetables.
- ¹⁵⁷ Calculated from total feed amount divided by total food amount. NA accounts for cases where consumption equalled zero.
- ¹⁵⁸ https://skogsstatistik.slu.se/pxweb/sv/OffStat/OffStat_Skogsmark_StandortVegetation/SM_Barproduktion_tab.px/; <https://www.skogsaktuellt.se/artikel/31849/svenska-bar-kan-bli-guld-varda.html>;
<https://www.skogssverige.se/hur-mycket-atliga-bar-finns-det-i-sverige-idag>
- ¹⁵⁹ <https://pulses.org/nap/what-are-pulses/>
- ¹⁶⁰ Crop products (production quantity, world totals in 2019). <https://www.fao.org/faostat/en/#data/QCL>
- ¹⁶¹ [Soy - Our World in Data](#)
- ¹⁶² [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ¹⁶³ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ¹⁶⁴ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ¹⁶⁵ Nemecek, T & Baumgartner, D. 2006. Environmental impacts of introducing grain legumes into European crop rotations and pig feed formulas. Concerted action GL-Pro WP4: Environmental analysis. Final report. Deliverable 4.3.
- ¹⁶⁶ Smil, V. 2002. Nitrogen and Food Production: Proteins for human Diets. *Ambio* 31(2), 126-131.
- ¹⁶⁷ Andersson, A. Jansson, A & Hellström, A. 2006. The Swedish monitoring of pesticide residues in food of plant origin: 2005. EC and National Report. Livsmedelsverket. Rapport 13 - 2006.
- ¹⁶⁸ Jungbluth, N. & Frischknecht. 2007. LCA of Imported Agricultural Products – Impacts due to Deforestation and Burning of residues. Konferens. LCA in Foods, Gothenburg, Sweden, 25 - 26 April, 2007, Organised by SIK, The Swedish Institute for Food and Biotechnology.
- ¹⁶⁹ [Soy - Our World in Data](#)
- ¹⁷⁰ [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ¹⁷¹ [Soy - Our World in Data](#)
- ¹⁷² Goldman, E., M.J. Weisse, N. Harris, and M. Schneider. 2020. "Estimating the Role of Seven Commodities in Agriculture-Linked Deforestation: Oil Palm, Soy, Cattle, Wood Fiber, Cocoa, Coffee, and Rubber." Technical Note. Washington, DC: World Resources Institute. Available online at: wri.org/publication/estimating-the-role-of-sevencommodities-in-agriculture-linked-deforestation
- ¹⁷³ [Soy - Our World in Data](#)
- ¹⁷⁴ Kuepper, B. and M. Stravens (January 2022), 'Mapping the European Soy Supply Chain – Embedded Soy in Animal Products Consumed in the EU27+UK'. Profundo, Amsterdam, The Netherlands. Table 11 Soy consumption in the EU27+UK (2020).
- ¹⁷⁵ [Soy - Our World in Data](#)
- ¹⁷⁶ Source: FAOSTAT production data for pulses, world total 2019. [FAOSTAT. 85,456,000 tonnes of pulses were produced in 2019, of which 19,761,000 tonnes \(23%\) were allocated to feed.](#)
- ¹⁷⁷ Harwatt, H., Sabaté, J., Eshel, G., Soret, S. and Ripple, W. (2017), 'Substituting beans for beef as a contribution toward US climate change targets', *Climatic Change*, 143(1–2), doi: 10.1007/s10584-017-1969-1
- ¹⁷⁸ Alexander, P. et al. (2016), 'Human appropriation of land for food: The role of diet', *Global Environmental Change*, 41, pp. 88–98, <http://www.sciencedirect.com/science/article/pii/S0959378016302370?via%3Dihub#bib0330>
- ¹⁷⁹ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives.

-
- ¹⁸⁰ WWF-Denmark-Risky-Business-Summary-Report-design_final_23-January.pdf (3keel.com).
- ¹⁸¹ Please note: This information was provided by regional experts during a series of workshops and has not been verified.
- ¹⁸² [UK's first commercial crop of chickpeas harvested in Norfolk | Food & drink industry | The Guardian](#)
- ¹⁸³ Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines (livsmedelsverket.se); M. Vogelsang-O'Dwyer. Comparison of Faba Bean Protein Ingredients Produced Using Dry Fractionation and Isoelectric Precipitation: Techno-Functional, Nutritional and Environmental Performance, 2020; Gonera, A. og Milford, A.B.: The plant protein trend in Norway. Market overview and future perspectives". Rapport fra Nofima og NIBIO (2018); Abrahamsen U. m.fl: Muligheter for økt proteinproduksjon på kornarealene. Rapport fra NIBIO (2019).
- ¹⁸⁴ Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines (livsmedelsverket.se)
- ¹⁸⁵ <https://projects.luke.fi/scenoprot/en/>
- ¹⁸⁶ Project Info - Leg4Life
- ¹⁸⁷ Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines (livsmedelsverket.se)
- ¹⁸⁸ Seehusen T, Uhlen A.K. Analyses of Yield Gaps for the production of wheat and barley in Norway - Potential to increase yields on existing farmland. NIBIO Rapport;5(166) 2019 <https://core.ac.uk/download/pdf/287378885.pdf>
- ¹⁸⁹ Rööös E, Carlsson G, Ferawati F, Hefni M, Stephan A, Tidåker P, Witthöft C. Less meat, more legumes: Prospects and challenges in the transition toward sustainable diets in Sweden. *Renewable Agriculture and Food Systems* 2020;35:192-205. doi:10.1017/S1742170518000443
- ¹⁹⁰ [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ¹⁹¹ [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ¹⁹² Rööös E, Carlsson G, Ferawati F, Hefni M, Stephan A, Tidåker P, Witthöft C. Less meat, more legumes: Prospects and challenges in the transition toward sustainable diets in Sweden. *Renewable Agriculture and Food Systems* 2020;35:192-205. doi:10.1017/S1742170518000443
- ¹⁹³ Springmann, M., Clark, M., Mason-D'Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>
- ¹⁹⁴ Includes beans, peas, pulses (other), and soybeans. Pulses (other) is a group which includes broad beans, chickpeas and lentils.
- ¹⁹⁵ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ¹⁹⁶ NA accounts for cases where consumption equalled zero. Source: FAOSTAT food balances available at [FAOSTAT](#)
- ¹⁹⁷ Benton et al 2021. Food system impacts on biodiversity loss Three levers for food system transformation in support of nature. Chatham House report, February 2021.
- ¹⁹⁸ World production of fish and seafood, 2019. <https://www.fao.org/faostat/en/#data/FBS>
- ¹⁹⁹ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.
- ²⁰⁰ [Fish and Overfishing - Our World in Data](#)
- ²⁰¹ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.
- ²⁰² [Fish and Overfishing - Our World in Data](#)
- ²⁰³ Halpern, B.S., Frazier, M., Verstaen, J. et al. The environmental footprint of global food production. *Nat Sustain* 5, 1027–1039 (2022). <https://doi.org/10.1038/s41893-022-00965-x>
- ²⁰⁴ Winther U, Skontorp Hognes E, Jafarzadeh S, Ziegler F (2020) Greenhouse gas emissions of Norwegian seafood products in 2017. Sintef
- ²⁰⁵ Winther U, Skontorp Hognes E, Jafarzadeh S, Ziegler F (2020) Greenhouse gas emissions of Norwegian seafood products in 2017. Sintef
- ²⁰⁶ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.
- ²⁰⁷ Henry C. Holm et al, Global ocean lipidomes show a universal relationship between temperature and lipid unsaturation, *Science* (2022). DOI: 10.1126/science.abn7455
- ²⁰⁸ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.
- ²⁰⁹ Henry C. Holm et al, Global ocean lipidomes show a universal relationship between temperature and lipid unsaturation, *Science* (2022). DOI: 10.1126/science.abn7455
- ²¹⁰ Gephart, J.A., Henriksson, P.J.G., Parker, R.W.R. et al. Environmental performance of blue foods. *Nature* 597, 360–365 (2021). <https://doi.org/10.1038/s41586-021-03889-2>
- ²¹¹ Sala, E., Mayorga, J., Bradley, D. et al. Protecting the global ocean for biodiversity, food and climate. *Nature* 592, 397–402 (2021). <https://doi.org/10.1038/s41586-021-03371-z>
- ²¹² Ziegler and Hilborn (2023) Fished or farmed: Life cycle impacts of salmon consumer decisions and opportunities for reducing impacts. *Science of the Total Environment* 854 (2023) 158591. <https://doi.org/10.1016/j.scitotenv.2022.158591>
- ²¹³ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.

-
- ²¹⁴ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.
- ²¹⁵ Based on data from 1690 fish farms. Source data: Gephart, J.A., Henriksson, P.J.G., Parker, R.W.R. *et al.* Environmental performance of blue foods. *Nature* **597**, 360–365 (2021). <https://doi.org/10.1038/s41586-021-03889-2>
- ²¹⁶ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.
- ²¹⁷ Based on data from 1690 fish farms. Source data: Gephart, J.A., Henriksson, P.J.G., Parker, R.W.R. *et al.* Environmental performance of blue foods. *Nature* **597**, 360–365 (2021). <https://doi.org/10.1038/s41586-021-03889-2>
- ²¹⁸ Silvenius, F., J. Grönroos, M. Kankainen, S. Kurppa, T. Mäkinen, and J. Vielma. 2017. "Impact of Feed Raw Material to Climate and Eutrophication Impacts of Finnish Rainbow Trout Farming and Comparisons on Climate Impact and Eutrophication between Farmed and Wild Fish." *Journal of Cleaner Production* **164**:1467-1473. doi: <https://doi.org/10.1016/j.jclepro.2017.07.069>.
- ²¹⁹ Based on data from 1690 fish farms. Source data: Gephart, J.A., Henriksson, P.J.G., Parker, R.W.R. *et al.* Environmental performance of blue foods. *Nature* **597**, 360–365 (2021). <https://doi.org/10.1038/s41586-021-03889-2>
- ²²⁰ Based on data from 1690 fish farms. Source data: Gephart, J.A., Henriksson, P.J.G., Parker, R.W.R. *et al.* Environmental performance of blue foods. *Nature* **597**, 360–365 (2021). <https://doi.org/10.1038/s41586-021-03889-2>
- ²²¹ Bolstad, Geir H., Sten Karlsson, Ingerid J. Hagen, Peder Fiske, Kurt Urdal, Harald Sægrov, Bjørn Florø-Larsen et al. "Introgression from farmed escapees affects the full life cycle of wild Atlantic salmon." *Science Advances* **7**, no. 52 (2021): eabj3397.
- ²²² [Fish and Overfishing - Our World in Data](#)
- ²²³ FAO. 2020. The State of World Fisheries and Aquaculture 2020. Sustainability in action. Rome. <https://doi.org/10.4060/ca9229en>
- ²²⁴ FAO. 2020. The State of World Fisheries and Aquaculture 2020. Sustainability in action. Rome. Figure 20: percentages of stocks fished at biologically sustainable and unsustainable levels, by FAO statistical area, 2017. <https://doi.org/10.4060/ca9229en>
- ²²⁵ [Fish and Overfishing - Our World in Data](#)
- ²²⁶ [Fish and Overfishing - Our World in Data](#)
- ²²⁷ Hiddink, J. G., Jennings, S., Sciberras, M., Szostek, C. L., Hughes, K. M., Ellis, N., ... & Kaiser, M. J. (2017). Global analysis of depletion and recovery of seabed biota after bottom trawling disturbance. *Proceedings of the National Academy of Sciences*, **114**(31), 8301-8306.
- ²²⁸ Lis Lindal Jørgensen, Benjamin Planque, Trude Hauge Thangstad, Grégoire Certain, Vulnerability of megabenthic species to trawling in the Barents Sea, *ICES Journal of Marine Science*, Volume 73, Issue suppl_1, January 2016, Pages i84–i97.
- ²²⁹ [Fish and Overfishing - Our World in Data](#)
- ²³⁰ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.
- ²³¹ Atwood, T., Connolly, R., Ritchie, E. et al. Predators help protect carbon stocks in blue carbon ecosystems. *Nature Clim Change* **5**, 1038–1045 (2015). <https://doi.org/10.1038/nclimate2763>
- ²³² Halpern, B.S., Frazier, M., Verstaen, J. et al. The environmental footprint of global food production. *Nat Sustain* **5**, 1027–1039 (2022). <https://doi.org/10.1038/s41893-022-00965-x>
- ²³³ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.
- ²³⁴ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.
- ²³⁵ Lebreton, L., Slat, B., Ferrari, F. et al. Evidence that the Great Pacific Garbage Patch is rapidly accumulating plastic. *Sci Rep* **8**, 4666 (2018). <https://doi.org/10.1038/s41598-018-22939-w>
- ²³⁶ Wilcox, C. et al. Understanding the sources and effects of abandoned, lost, and discarded fishing gear on marine turtles in northern Australia. *Conserv. Biol.* **29**, 198–206 (2015).
- ²³⁷ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives.
- ²³⁸ Sprague, M., Dick, J. & Tocher, D. Impact of sustainable feeds on omega-3 long-chain fatty acid levels in farmed Atlantic salmon, 2006–2015. *Sci Rep* **6**, 21892 (2016). <https://doi.org/10.1038/srep21892>
- ²³⁹ Future Nordic Diets. Exploring ways for sustainably feeding the Nordics. Johan Karlsson, Elin Röö, Tove Sjunnestrånd, Kajsa Pira, Malin Larsson, Bente Hesselund Andersen, Jacob Sørensen, Tapani Veistola, Jaana Rantakokko, Sirkku Manninen and Stein Brubæk.
- ²⁴⁰ Ziegler, F, Jafarzadeh, S, Hognes, ES, & Winther, U. Greenhouse gas emissions of Norwegian seafoods: From comprehensive to simplified assessment. *J Ind Ecol.* **2022**; **26**: 1908– 1919. <https://doi.org/10.1111/jiec.13150>
- ²⁴¹ Gullestad, P. & Blom, G. & Bakke, G. & Bogstad, B., 2015. "The "Discard Ban Package": Experiences in efforts to improve the exploitation patterns in Norwegian fisheries," *Marine Policy*, Elsevier, vol. 54(C), pages 1-9. <https://doi.org/10.1016/j.marpol.2014.09.025>

²⁴² Lisa Borges, Luis Cocas, Kåre Nolde Nielsen, Discard ban and balanced harvest: a contradiction? *ICES Journal of Marine Science*, Volume 73, Issue 6, May/June 2016, Pages 1632–1639, <https://doi.org/10.1093/icesjms/fsw065>

²⁴³ Lisa Borges, Luis Cocas, Kåre Nolde Nielsen, Discard ban and balanced harvest: a contradiction? *ICES Journal of Marine Science*, Volume 73, Issue 6, May/June 2016, Pages 1632–1639, <https://doi.org/10.1093/icesjms/fsw065>

²⁴⁴ <https://eurofish.dk/member-countries/norway/#:~:text=Approximately%2090%25%20of%20Norway%E2%80%99s%20catch%20volume%20comes%20from,Iceland%2C%20the%20Faroe%20Islands%2C%20Greenland%2C%20and%20the%20EU.>

²⁴⁵ Fish and seafood production quantity, 2019. <https://www.fao.org/faostat/en/#data/FBS>

²⁴⁶ Springmann, M., Clark, M., Mason-D'Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>

²⁴⁷ Source: FAOSTAT food balances available at [FAOSTAT](https://www.fao.org/faostat/en/#data/FBS). Includes all fish species and major seafood commodities, including crustaceans, cephalopods and other mollusc species. Data is based on per capita food supply at the consumer level, but does not account for food waste at the consumer level.

²⁴⁸ The World Bank, 2013. Fish to 2030: prospects for fisheries and aquaculture (No. 83177). The World Bank.

²⁴⁹ Future Nordic Diets. Exploring ways for sustainably feeding the Nordics. Johan Karlsson, Elin Röö, Tove Sjunnestrand, Kajsa Pira, Malin Larsson, Bente Hesselund Andersen, Jacob Sørensen, Tapani Veistola, Jaana Rantakokko, Sirkku Manninen and Stein Brubæk.

²⁵⁰ Thorsøe, M.H., Andersen, M.S., Brady, M.V. et al. Promise and performance of agricultural nutrient management policy: Lessons from the Baltic Sea. *Ambio* 51, 36–50 (2022).

<https://doi.org/10.1007/s13280-021-01549-3>; Ola Luthman, Malin Jonell, Patrik Rönnbäck, Max Troell, Strong and weak sustainability in Nordic aquaculture policies, *Aquaculture*, Volume 550, 2022, 737841, <https://doi.org/10.1016/j.aquaculture.2021.737841>.

²⁵¹ Ola Luthman, Malin Jonell, Patrik Rönnbäck, Max Troell, Strong and weak sustainability in Nordic aquaculture policies, *Aquaculture*, Volume 550, 2022, 737841, <https://doi.org/10.1016/j.aquaculture.2021.737841>.

²⁵² Ola Luthman, Malin Jonell, Patrik Rönnbäck, Max Troell, Strong and weak sustainability in Nordic aquaculture policies, *Aquaculture*, Volume 550, 2022, 737841, <https://doi.org/10.1016/j.aquaculture.2021.737841>.

²⁵³ [Mackerel \(Scomber scombrus\) in subareas 1–8 and 14, and in Division 9.a \(Northeast Atlantic and adjacent waters\) \(figshare.com\)](https://www.figshare.com/articles/report/Mackerel_Scomber_scombrus_in_subareas_1_8_and_14_and_in_Division_9_a_Northeast_Atlantic_and_adjacent_waters/19772392?backTo=/collections/ICES_Advice_2022/5796935)

²⁵⁴ https://ices-library.figshare.com/articles/report/Mackerel_Scomber_scombrus_in_subareas_1_8_and_14_and_in_Division_9_a_Northeast_Atlantic_and_adjacent_waters/19772392?backTo=/collections/ICES_Advice_2022/5796935

²⁵⁵ Pelagic fish include: herring, mackerel, capelin, blue whiting, bluefin tuna. Demersal fish include: cod, haddock, whiting, hake, pollock and plaice. Freshwater fish include: salmon (technically an anadromous species), trout, bass, carp, bream, eel, and sturgeon. Crustaceans include: crabs, lobsters, crayfish, shrimp, krill, and prawns. Molluscs include: mussels, scallops, oysters, clam and whelk. Cephalopods include: octopuses, squid, cuttlefish and nautilus.

²⁵⁶ Source: FAOSTAT food balances available at [FAOSTAT](https://www.fao.org/faostat/en/#data/FBS)

²⁵⁷ The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.

²⁵⁸ Future Nordic Diets. Exploring ways for sustainably feeding the Nordics. Johan Karlsson, Elin Röö, Tove Sjunnestrand, Kajsa Pira, Malin Larsson, Bente Hesselund Andersen, Jacob Sørensen, Tapani Veistola, Jaana Rantakokko, Sirkku Manninen and Stein Brubæk.

²⁵⁹ Gephart, J.A., Henriksson, P.J.G., Parker, R.W.R. et al. Environmental performance of blue foods. *Nature* 597, 360–365 (2021). <https://doi.org/10.1038/s41586-021-03889-2>

²⁶⁰ Gephart, J.A., Henriksson, P.J.G., Parker, R.W.R. et al. Environmental performance of blue foods. *Nature* 597, 360–365 (2021). <https://doi.org/10.1038/s41586-021-03889-2>

²⁶¹ Gephart, J.A., Henriksson, P.J.G., Parker, R.W.R. et al. Environmental performance of blue foods. *Nature* 597, 360–365 (2021). <https://doi.org/10.1038/s41586-021-03889-2>

²⁶² The role of seafood for sustainable and healthy diets. The EAT-Lancet commission report through a blue lens. Max Troell, Malin Jonell and Beatrice Crona. 2019.

-
- ²⁶³ Source: FAOSTAT production data, world total 2020 [FAOSTAT](#)
- ²⁶⁴ Gerber, P.J., Steinfeld, H., Henderson, B., Mottet, A., Opio, C., Dijkman, J., Faluccci, A. & Tempio, G. 2013. Tackling climate change through livestock – A global assessment of emissions and mitigation opportunities. Food and Agriculture Organization of the United Nations (FAO), Rome.
- ²⁶⁵ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/science.aaq0216.
- ²⁶⁶ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 *Environ. Res. Lett.* 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ²⁶⁷ Benton and Harwatt (2022) Sustainable agriculture and food systems. Comparing contrasting and contested versions. RESEARCH PAPER. 24 MAY 2022. ISBN: 978 1 78413 526 3. DOI: 10.55317/9781784135263
- ²⁶⁸ M. Torres-Miralles, K. Särkelä, K. Koppelmäki, M. Lamminen, H.L. Tuomisto, I. Herzon. Contribution of High Nature Value farming systems to sustainable livestock production: A case from Finland. *Science of The Total Environment*, Volume 839, 2022. 156267. <https://doi.org/10.1016/j.scitotenv.2022.156267>; Bjørn Molt Petersen, Marie Trydeman Knudsen, John Erik Hermansen, Niels Halberg, An approach to include soil carbon changes in life cycle assessments, *Journal of Cleaner Production*, Volume 52, 2013, Pages 217-224, <https://doi.org/10.1016/j.jclepro.2013.03.007>.
- ²⁶⁹ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 *Environ. Res. Lett.* 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>
- ²⁷⁰ Hayek, M. N., Harwatt, H., Ripple, W. J. and Mueller, N. D. (2020), 'The carbon opportunity cost of animal-sourced food production on land', *Nature Sustainability* (2020), doi: 10.1038/s41893-020-00603-4
- ²⁷¹ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/science.aaq0216.
- ²⁷² UK Greenhouse Gas Inventory 1990 to 2019: Annual Report for submission under the Framework Convention on Climate Change. Table A 3.3.9 Areas of UK Crops and quantities of fertiliser applied for 2019. UK BIES. 15 April 2021.
- ²⁷³ Roos et al (2014) Communicating the environmental impact of meat production: challenges in the development of a Swedish meat guide. *Journal of Cleaner Production* 73 (2014) 154e164; Højberg A L, Thodsen H, Børgesen C D, Tornbjerg H, Nordstrøm B O, Troldborg L, Hoffmann C C, Kjeldgaard A, Holm H, Audet J, Ellermann T, Christensen J H, Bach E O and Pedersen B F 2021 National kvælstofmodel – version 2020. Bilag Online: https://www.geus.dk/Media/637576521825668552/NKM2020_Bilag_18maj2021_web.pdf; Saghar K. Motarjemi, Merete E. Styczen, Rasmus J. Petersen, Kasper J.S. Jensen, Finn Plauborg, Effects of different drainage conditions on nitrogen losses of an agricultural sandy loam soil, *Journal of Environmental Management*, Volume 325, Part A, 2023, 116267, <https://doi.org/10.1016/j.jenvman.2022.116267>.
- ²⁷⁴ Roos et al (2014) Communicating the environmental impact of meat production: challenges in the development of a Swedish meat guide. *Journal of Cleaner Production* 73 (2014) 154e164.
- ²⁷⁵ Roos et al (2014) Communicating the environmental impact of meat production: challenges in the development of a Swedish meat guide. *Journal of Cleaner Production* 73 (2014) 154e164.
- ²⁷⁶ Heikkinen, Jaakko, Elise Ketoja, Visa Nuutinen, and Kristiina Regina. 2013. "Declining Trend of Carbon in Finnish Cropland Soils in 1974–2009." *Global Change Biology* 19 (5): 1456-1469. doi:<https://doi.org/10.1111/gcb.12137>.
- ²⁷⁷ Hayek, M. N., Harwatt, H., Ripple, W. J. and Mueller, N. D. (2020), 'The carbon opportunity cost of animal-sourced food production on land', *Nature Sustainability* (2020), doi: 10.1038/s41893-020-00603-4
- ²⁷⁸ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392), pp. 987–92, <https://doi.org/10.1126/science.aaq0216>.
- ²⁷⁹ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392), pp. 987–92, <https://doi.org/10.1126/science.aaq0216>.
- ²⁸⁰ David De Pue, Andreas Bral, Jeroen Buysse, Abatement of ammonia emissions from livestock housing fine-tuned according to impact on protected habitats, *Agricultural Systems*, Volume 176, 2019, 102667, <https://doi.org/10.1016/j.agsy.2019.102667>; Benton, T. G. et al. (2021), Food system impacts on biodiversity loss: Three levers for food system transformation in support of nature, Research Paper, London: Royal Institute of International Affairs, <https://www.chathamhouse.org/2021/02/food-system-impacts-biodiversity-loss>; Leip et al 2015 *Environ. Res. Lett.* 10 115004 DOI 10.1088/1748-9326/10/11/115004.
- ²⁸¹ Ritchie, H. and Roser, M. (2021), 'Soy', <https://ourworldindata.org/soy>; World Resources Institute (2020), Estimating the Role of Seven Commodities in Agriculture-Linked Deforestation: Oil Palm, Soy, Cattle, Wood Fiber, Cocoa, Coffee, and Rubber, Technical Note, Washington, DC: World Resources Institute, <https://www.wri.org/research/estimating-role-seven-commodities-agriculture-linked-deforestation-oil-palm-soy-cattle>.
- ²⁸² Machovina et al (2015). Biodiversity conservation: The key is reducing meat consumption. *Science of The Total Environment*. Volume 536, 1 December 2015, Pages 419-431. <https://doi.org/10.1016/j.scitotenv.2015.07.022>
- ²⁸³ Sun et al (2022) Global Human Consumption Threatens Key Biodiversity Areas. *Environ. Sci. Technol.* 2022, 56, 9003–9014
- ²⁸⁴ Land use: area 2020. [FAOSTAT](#)
- ²⁸⁵ Filazzola et al (2020) The effects of livestock grazing on biodiversity are multi-trophic: a meta-analysis. *Ecology Letters* (2020) 23: 1298 – 1309.
- ²⁸⁶ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/science.aaq0216.
- ²⁸⁷ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/science.aaq0216.

- ²⁸⁸ Clark, M and Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. 2017 *Environ. Res. Lett.* 12 064016. <http://iopscience.iop.org/1748-9326/12/6/064016>; Halpern, B.S., Frazier, M., Verstaen, J. et al. The environmental footprint of global food production. *Nat Sustain* 5, 1027–1039 (2022). <https://doi.org/10.1038/s41893-022-00965-x>; Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/science.aaq0216; Roos et al (2014) Communicating the environmental impact of meat production: challenges in the development of a Swedish meat guide. *Journal of Cleaner Production* 73 (2014) 154e164.
- ²⁸⁹ Hayek, M. and Garrett, R (2018) Nationwide shift to grass-fed beef requires larger cattle population. *Environ. Res. Lett.* 13 (2018) 084005.
- ²⁹⁰ Benton, T. G. and Harwatt, H. (2022), Sustainable agriculture and food systems: Comparing contrasting and contested versions, Research Paper, London: Royal Institute of International Affairs, <https://doi.org/10.55317/9781784135263>
- ²⁹¹ Harwatt, H. (2018), 'Including animal to plant protein shifts in climate change mitigation policy: a proposed three-step strategy', *Climate Policy*, 19(5), pp. 533–541, <https://doi.org/10.1080/14693062.2018.1528965>.
- ²⁹² United Nations Development Programme & Climate and Clean Air Coalition (2020), *Global Methane Assessment: Benefits and Costs of Mitigating Methane Emissions*, Nairobi: United Nations Environment Programme, <https://www.ccacoalition.org/en/resources/global-methane-assessment-full-report>.
- ²⁹³ Clark et al 2020; Springmann et al 2018; Willet et al 2019.
- ²⁹⁴ Poore and Nemecek (2018); Gonzalez-Garcia et al (2015) Life cycle assessment of pig meat production: Portuguese case study and proposal of improvement options. *Journal of Cleaner Production*, V 100, pp 126-139; Noya et al (2017) Life Cycle Assessment of pig production: A case study in Galicia. *Journal of Cleaner Production*. V 142 (4) pp 4327-4338.
- ²⁹⁵ Gerber, P.J., Steinfeld, H., Henderson, B., Mottet, A., Opio, C., Dijkman, J., Falucci, A. & Tempio, G. 2013. Tackling climate change through livestock – A global assessment of emissions and mitigation opportunities. Food and Agriculture Organization of the United Nations (FAO), Rome.
- ²⁹⁶ Poore and Nemecek (2018).
- ²⁹⁷ Poultry meat production, world total 2015 – 2020. <https://www.fao.org/faostat/en/#data/FBS>
- ²⁹⁸ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives.
- ²⁹⁹ Wood A, Gordon LJ, Rööös E, Karlsson JO, Häyhä T, Bignet V, Rydenstam T, Hård af Segerstad L, Bruckner M. Nordic Food Systems for Improved Health and Sustainability; Stockholm Resilience Centre, University of Stockholm: Stockholm, Sweden, 2019.
- ³⁰⁰ Thorsøe, M.H., Andersen, M.S., Brady, M.V. et al. Promise and performance of agricultural nutrient management policy: Lessons from the Baltic Sea. *Ambio* 51, 36–50 (2022). <https://doi.org/10.1007/s13280-021-01549-3>
- ³⁰¹ Staalstrøm A, Walday M, Vogelsang C, Frigstad H, Borgersen G, Albreten J, Naustvoll L-J (2022) Utredning av behovet for å redusere tilførselen av nitrogen til Ytre Oslofjord. NIVA; Styrket innsats for renere Oslofjord - men langt frem | Statsforvalteren i Oslo og Viken.
- ³⁰² Sofie Hellsten, Tommy Dalgaard, Katri Rankinen, Kjetil Tørseth, Airi Kulmala, Eila Turtola, Filip Moldan, Kajsa Pira, Kristoffer Piil, Lars Bakken, Marianne Bechmann and Stina Olofsson. Nordic nitrogen and agriculture Policy, measures and recommendations to reduce environmental impact. *TemaNord* 2017:547; Leip et al 2015 *Environ. Res. Lett.* 10 115004DOI 10.1088/1748-9326/10/11/115004
- ³⁰³ Hellsten, S., et al. (2017), Nordic nitrogen and agriculture: Policy, measures and recommendations to reduce environmental impact, *TemaNord*, Nordic Council of Ministers, Copenhagen. <https://doi.org/10.6027/TN2017-547>; T. Dalgaard, B. Hansen, B. Hasler, O. Hertel, N. J. Hutchings, B. H. Jacobsen, L. S. Jensen, B. Kronvang, J. E. Olesen, J. K. Schjørring, I. S. Kristensen, M. Graversgaard, M. Termansen, H. Vejre, Policies for agricultural nitrogen management—Trends, challenges and prospects for improved efficiency in Denmark. *Environ. Res. Lett.* 6, 115002 (2014) <https://www.reuters.com/world/europe/eu-okays-161-bln-dutch-govt-buy-out-farmers-reduce-nitrogen-2023-05-02/>;
- ³⁰⁴ <https://www.reuters.com/world/europe/eu-okays-161-bln-dutch-govt-buy-out-farmers-reduce-nitrogen-2023-05-02/>; Boezeman, D., De Pue, D., Graversgaard, M., Möckel, S. 2023. Less Livestock in North-western Europe? Discourses and Drivers Behind Livestock Buyout Policies. *Eurochoices*. volume 22, issue 2.
- ³⁰⁵ Hallstrom E, Davis J, Hakansson N, Ahlgren S, Åkesson A, Wolk A, et al. Dietary environmental impacts relative to planetary boundaries for six environmental indicators-A population-based study. *Journal of Cleaner Production* 2022;373; Moberg E, Karlsson Potter H, Wood A, Hansson P-A, Rööös E. Benchmarking the Swedish Diet Relative to Global and National Environmental Targets—Identification of Indicator Limitations and Data Gaps. *Sustainability* 2020;12:1407; Wood A, Gordon LJ, Rööös E, Karlsson JO, Häyhä T, Bignet V, Rydenstam T, Hård af Segerstad L, Bruckner M. Nordic Food Systems for Improved Health and Sustainability; Stockholm Resilience Centre, University of Stockholm: Stockholm, Sweden, 2019.
- ³⁰⁶ Karlsson J, Rööös E, Sjunnestrand T, Pira K, Larsson M, Hessellund Andersen B, Sørensen J, Veistola T, Rantakokko J, Manninen S, Brubæk S. Future Nordic Diets Exploring ways for sustainably feeding the Nordics. *TemaNord* 2017:566. <https://norden.diva.portal.org/smash/get/diva2:1163192/FULLTEXT01.pdf>
- ³⁰⁷ Mittenzwei K, van Oort B (2022) Hvordan øke selvforsyningsgraden i norsk jordbruk? *Ruralis* note 3/22
- ³⁰⁸ Bakken AK, Mittenzwei K (2023) Produksjonspotensial i jordbruket og nasjonal sjølforsyning med mat. Utredning for Klimautvalget 2050. NIBIO, Ruralis

- ³⁰⁹ Tiainen, J., Hyvönen, T., Hagner, M., Huusela-Veistola, E., Louhi, P., Miettinen, A., Nieminen, T M., Palojärvi, A., Seimola, T., Taimisto, P & Virkajärvi, P 2020, ' Biodiversity in intensive and extensive grasslands in Finland : the impacts of spatial and temporal changes of agricultural land use ', *Agricultural and Food Science*, vol. 29, no. 2, pp. 68-97. <https://doi.org/10.23986/afsci.86811>; Leip et al 2015 *Environ. Res. Lett.* 10 115004DOI 10.1088/1748-9326/10/11/115004
- ³¹⁰ Adrian Leip et al 2015 *Environ. Res. Lett.* 10 115004DOI 10.1088/1748-9326/10/11/115004
- ³¹¹ Bengtsson, J., J. M. Bullock, B. Egoh, C. Everson, T. Everson, T. O'Connor, P. J. O'Farrell, H. G. Smith, and R. Lindborg. 2019. Grasslands—more important for ecosystem services than you might think. *Ecosphere* 10(2):e02582. 10.1002/ecs2.2582; <https://www.sverigesmiljomal.se/miljomalen/ett-rikt-odlingslandskap/betesmarker-och-slatteangar/>
- ³¹² Karlsson J. Livestock as resource users and landscape managers. A food system perspective. *Acta Universitatis Agriculturae Seuciae*, Doctoral thesis no 2022:19; Tiainen, J., Hyvönen, T., Hagner, M., Huusela-Veistola, E., Louhi, P., Miettinen, A., Nieminen, T M., Palojärvi, A., Seimola, T., Taimisto, P & Virkajärvi, P 2020, ' Biodiversity in intensive and extensive grasslands in Finland : the impacts of spatial and temporal changes of agricultural land use ', *Agricultural and Food Science*, vol. 29, no. 2, pp. 68-97. <https://doi.org/10.23986/afsci.86811>
- ³¹³ Chang, J., Ciais, P., Gasser, T. et al. Climate warming from managed grasslands cancels the cooling effect of carbon sinks in sparsely grazed and natural grasslands. *Nat Commun* 12, 118 (2021). <https://doi.org/10.1038/s41467-020-20406-7>
- ³¹⁴ IDH (2021) European Soy Monitor; Insights on European uptake of responsible, deforestation and conversion-free soy in 2020. June 2022.
- ³¹⁵ Schilling-Vacaflor A, Lenschow A, Challies E, Cotta B, Newig J. Contextualizing certification and auditing: Soy certification and access of local communities to land and water in Brazil. *World Development* 2021;140;105281. <https://doi.org/10.1016/j.worlddev.2020.105281>.
- ³¹⁶ <https://d5i6is0eze552.cloudfront.net/documents/Publikasjoner/Andre-rapporter/Salmon-on-soy-beans-deforestation-and-land-conflict-in-Brazil.pdf?mtime=20181029093010>
- ³¹⁷ Haskell2_2022_BioVit.pdf (unit.no); Faba bean production under Norwegian growing conditions | (foodprofuture.no); Denmark: Ancient riddle solved: Danish researchers find a way to make faba beans safe for all to eat – University of Copenhagen (ku.dk).
- ³¹⁸ <https://www.leg4life.fi/en/project-info/>
- ³¹⁹ https://www.lbhi.is/images/pdf/utgefird%20efni/fjolrit%20rannsoknastofnunar%20landbunadarins/rit_lbhi_nr_133_ok.pdf
- ³²⁰ Cederberg et al (2009). Greenhouse gas emissions from Swedish production of meat, milk and eggs 1990 and 2005. SIK Report No 793.
- ³²¹ Cederberg et al (2009). Greenhouse gas emissions from Swedish production of meat, milk and eggs 1990 and 2005. SIK Report No 793.
- ³²² <https://www.slu.se/institutioner/husdjurens-utfodring-varld/nyheter-huv/soja-till-husdjur/>
- ³²³ Röös et al. 2015; Moberg et al. 2020
- ³²⁴ Mogensen, L., Kristensen, T., Nielsen, N.I., Spleth, P., Henriksson, M., Swensson, C., Hessel, A., Vestergaard, M., 2015. Greenhouse gas emissions from beef production systems in Denmark and Sweden. *Livest. Sci.* 174 (April 2015), 126–143. <https://doi.org/10.1016/j.livsci.2015.01.021>.
- ³²⁵ Hietala S, Heusala H, Katajajuuri J-M, Järvenranta K, Virkajärvi P, Huuskonen A, et al. Environmental life cycle assessment of Finnish beef – cradle-to-farm gate analysis of dairy and beef breed beef production. *Agricultural Systems* [Internet]. 2021 Dec 1;194.
- ³²⁶ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/science.aaq0216.
- ³²⁷ Tiainen, J., Hyvönen, T., Hagner, M., Huusela-Veistola, E., Louhi, P., Miettinen, A., Nieminen, T M., Palojärvi, A., Seimola, T., Taimisto, P & Virkajärvi, P 2020, ' Biodiversity in intensive and extensive grasslands in Finland : the impacts of spatial and temporal changes of agricultural land use ', *Agricultural and Food Science*, vol. 29, no. 2, pp. 68-97.
- ³²⁸ Please note: This information was provided by regional experts during a series of workshops and has not been verified.
- ³²⁹ Lehtonen, Heikki, Ellen Huan-Niemi, and Jyrki Niemi. 2022. "The Transition of Agriculture to Low Carbon Pathways with Regional Distributive Impacts." *Environmental Innovation and Societal Transitions* 44: 1-13. doi:<https://doi.org/10.1016/j.eist.2022.05.002>.
- ³³⁰ Heikkinen, J. (2016). Carbon storage of Finnish agricultural mineral soils and its long-term change. University of Helsinki <https://helda.helsinki.fi/bitstream/handle/10138/160237/CARBONST.pdf?sequence=1&isAllowed=y>
- ³³¹ <https://www.agrianalyse.no/getfile.php/13898-1513669775/Dokumenter/Dokumenter%202014/R6%20%C3%98kt%20matproduksjon%20p%C3%A5%20norske%20arealer.pdf>
- ³³² Mittenzwei K, Milford AB, Grønland A (2017) Status og potensial for økt produksjon og forbruk av vegetabilske matvarer i Norge. NIBIO note.
- ³³³ AgriAnalyse (2023) Landbruksbarometeret 2023. AgriAnalyse.
- ³³⁴ The agricultural policy environment in Norway | Policies for the Future of Farming and Food in Norway | OECD iLibrary (oecd-ilibrary.org).

- ³³⁵ Benton, T. G. and Harwatt, H. (2022), Sustainable agriculture and food systems: Comparing contrasting and contested versions, Research Paper, London: Royal Institute of International Affairs, <https://doi.org/10.55317/9781784135263>
- ³³⁶ Springmann, M., Clark, M., Mason-D’Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>
- ³³⁷ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ³³⁸ Springmann, M., Clark, M., Mason-D’Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>
- ³³⁹ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ³⁴⁰ Springmann, M., Clark, M., Mason-D’Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>
- ³⁴¹ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ³⁴² Mittenzwei K, van Oort B (2022) Hvordan øke selvforsyningsgraden i norsk jordbruk? *Ruralis note* 3/22
- ³⁴³ Hayek, M. N., Harwatt, H., Ripple, W. J. and Mueller, N. D. (2020), ‘The carbon opportunity cost of animal-sourced food production on land’, *Nature Sustainability* (2020), doi: 10.1038/s41893-020-00603-4; Sun, Z. et al. (2022), ‘Dietary change in high-income nations alone can lead to substantial double climate dividend’, *Nature Food*, 2(12), pp. 29–37, <https://doi.org/10.1038/s43016-021-00431-5>;
- ³⁴⁴ Source: FAOSTAT production data, world total 2019 [FAOSTAT](#)
- ³⁴⁵ Source: FAOSTAT feed data, world total 2019 [FAOSTAT](#)
- ³⁴⁶ Harwatt, H. (2018), ‘Including animal to plant protein shifts in climate change mitigation policy: a proposed three-step strategy’, *Climate Policy*, 19(5), pp. 533–541, <https://doi.org/10.1080/14693062.2018.1528965>.
- ³⁴⁷ Poore, J. and Nemecek, T. (2018), ‘Reducing food’s environmental impacts through producers and consumers’, *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ³⁴⁸ Poore, J. and Nemecek, T. (2018), ‘Reducing food’s environmental impacts through producers and consumers’, *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ³⁴⁹ Nunan, C. 2022. Ending routine farm antibiotic use in Europe. Achieving responsible farm antibiotic use through improving animal health and welfare in pig and poultry production. edited by Written by Cólín Nunan for the European Public Health Alliance (EPHA).
- ³⁵⁰ [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ³⁵¹ Benton, T et al (2023) Forthcoming. Assessing the environmental sustainability of diets – an overview of approaches and identification of 5 key considerations for comprehensive assessments.
- ³⁵² Food supply quantity 2019. Source: [FAOSTAT](#)
- ³⁵³ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives.
- ³⁵⁴ <https://www.slu.se/institutioner/husdjurens-utfodring-varld/nyheter-huv/soja-till-husdjur/>
- ³⁵⁵ Landbruksdirektoratet. 2021. Bruk av norske fôrressurser. Utredning av forbedring av virkemidler med sikte på økt produksjon og bruk av norsk fôr. Norwegian Directorate of Agriculture.
- ³⁵⁶ Future Nordic Diets. Exploring ways for sustainably feeding the Nordics. Johan Karlsson, Elin Rööf, Tove Sjunnestrand, Kajsa Pira, Malin Larsson, Bente Hesselund Andersen, Jacob Sørensen, Tapani Veistola, Jaana Rantakokko, Sirkku Manninen and Stein Brubæk.
- ³⁵⁷ [Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines \(livsmedelsverket.se\)](#)
- ³⁵⁸ Flysjö et al 2011 The impact of various parameters on the carbon footprint of milk production in New Zealand and Sweden. July 2011, Pages 459-469.
- ³⁵⁹ Kristensen, T., Mogensen, L., Trydeman Knudsen, M., & Hermansen, J.E. (2011). Effect of production system and farming strategy on greenhouse gas emissions from commercial dairy farms in a life cycle approach including effect of different allocation methods. *Livest. Sci.* 140:136-148.
- ³⁶⁰ Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines (livsmedelsverket.se)
- ³⁶¹ Towards Environmentally Sound Dietary Guidelines - Scientific Basis for Environmental Assessment of the Swedish National Food Agency's Dietary Guidelines (livsmedelsverket.se)
- ³⁶² Karlsson J, Rööf E, Sjunnestrand T, Pira K, Larsson M, Hesselund Andersen B, Sørensen J, Veistola T, Rantakokko J, Manninen S, Brubæk S. Future Nordic Diets Exploring ways for sustainably feeding the Nordics. *TemaNord* 2017:566.
- ³⁶³ van Selm, B., Frehner, A., de Boer, I.J.M. et al. Circularity in animal production requires a change in the EAT-Lancet diet in Europe. *Nat Food* 3, 66–73 (2022). <https://doi.org/10.1038/s43016-021-00425-3>
- ³⁶⁴ Springmann, M., Clark, M., Mason-D’Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>
- ³⁶⁵ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ³⁶⁶ Source: FAOSTAT production data, world total 2019 [FAOSTAT](#)
- ³⁶⁷ Poore, J. and Nemecek, T. (2018), ‘Reducing food’s environmental impacts through producers and consumers’, *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.

- ³⁶⁸ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ³⁶⁹ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives.
- ³⁷⁰ Springmann, M., Clark, M., Mason-D'Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>
- ³⁷¹ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ³⁷² Karlsson JO, Garnett T, Rollins NC, Rööf E. The carbon footprint of breastmilk substitutes in comparison with breastfeeding. *J Clean Prod.* 2019 Jun 10;222:436-445. doi: 10.1016/j.jclepro.2019.03.043. PMID: 31190697; PMCID: PMC6472111.
- ³⁷³ Andresen, E.C.; Hjelkrem, A.-G.R.; Bakken, A.K.; Andersen, L.F. Environmental Impact of Feeding with Infant Formula in Comparison with Breastfeeding. *Int. J. Environ. Res. Public Health* 2022, 19, 6397. <https://doi.org/10.3390/ijerph19116397>
- ³⁷⁴ Joffe N, Webster F, Shenker N. Support for breastfeeding is an environmental imperative *BMJ* 2019; 367 :l5646 doi:10.1136/bmj.l5646
- ³⁷⁵ Aoife Long, Kian Mintz-Woo, Hannah Daly, Maeve O'Connell, Beatrice Smyth, Jerry D. Murphy, Infant feeding and the energy transition: A comparison between decarbonising breastmilk substitutes with renewable gas and achieving the global nutrition target for breastfeeding, *Journal of Cleaner Production*, Volume 324, 2021, 129280, <https://doi.org/10.1016/j.jclepro.2021.129280>.
- ³⁷⁶ Production quantity data from FAOSTAT (2020) <https://www.fao.org/faostat/en/#data/QCL>
- ³⁷⁷ Assessing and reducing the environmental impact of tea cultivation. Thushari Lakmini Wijeratne, Tea Research Institute, Sri Lanka. February 2018.
- ³⁷⁸ Goldman, E., M.J. Weisse, N. Harris, and M. Schneider. 2020. "Estimating the Role of Seven Commodities in Agriculture-Linked Deforestation: Oil Palm, Soy, Cattle, Wood Fiber, Cocoa, Coffee, and Rubber." Technical Note. Washington, DC: World Resources Institute. Available online at: wri.org/publication/estimating-the-role-of-seven-commodities-in-agriculture-linked-deforestation
- ³⁷⁹ Source: Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216. Fig. S13. GHG emissions, acidification, and eutrophication by stage of the supply chain by product.
- ³⁸⁰ Assessing and reducing the environmental impact of tea cultivation. Thushari Lakmini Wijeratne, Tea Research Institute, Sri Lanka. February 2018.
- ³⁸¹ Source: Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ³⁸² Anna-Lena Klapp, Nico Feil, Antje Risius, A Global Analysis of National Dietary Guidelines on Plant-Based Diets and Substitutions for Animal-Based Foods, *Current Developments in Nutrition*, Volume 6, Issue 11, 2022, <https://doi.org/10.1093/cdn/nzac144>.
- ³⁸³ <https://www.prnewswire.com/news-releases/rising-trend-of-veganism-to-steer-plant-based-milk-market-past-us-32-billion-by-2031-301606534.html>
- ³⁸⁴ Storhaug CL, Fosse SK, Fadnes LT. Country, regional, and global estimates for lactose malabsorption in adults: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol.*, 2 (10) (2017), pp. 738-746
- ³⁸⁵ Singh-Povel, C., Van Gool, M., Gual Rojas, A., Bragt, M., Kleijnijenhuis, A., & Hettinga, K. (2022). Nutritional content, protein quantity, protein quality and carbon footprint of plant-based drinks and semi-skimmed milk in the Netherlands and Europe. *Public Health Nutrition*, 25(5), 1416-1426. doi:10.1017/S1368980022000453
- ³⁸⁶ Rööf E, Patel M, Spångberg J (2015) Miljöpåverkan från mjölk och havredryck. En scenarionanalys som inkluderar alternativt markanvändning samt olika infallsvinklar på behovet av nötkött och protein. Department of Energy and Technology, Swedish University of Agricultural Sciences, Uppsala, Sweden.
- ³⁸⁷ Elin Rööf, Mikaela Patel, Johanna Spångberg, Producing oat drink or cow's milk on a Swedish farm — Environmental impacts considering the service of grazing, the opportunity cost of land and the demand for beef and protein, *Agricultural Systems*, Volume 142, 2016, Pages 23-32, <https://doi.org/10.1016/j.agsy.2015.11.002>.
- ³⁸⁸ Westhoek, H., Rood, T., Van De Berg, M., Janse, J., Nijdam, D., Reudink, M., & Stehfest, E. (2011). The protein puzzle: The consumption and production of meat, dairy and fish in the European Union. *The Protein Puzzle: The Consumption and Production of Meat, Dairy and Fish in the European Union*.
- ³⁸⁹ Data based on a meta-analysis of life cycle assessment studies across the supply chain. Source: [Dairy vs. plant-based milk: what are the environmental impacts? - Our World in Data](#)
- ³⁹⁰ <https://ourworldindata.org/pollinator-dependence>
- ³⁹¹ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ³⁹² Source: FAOSTAT food balances available at [FAOSTAT](#)

-
- ³⁹³ Jakobsen et al 2023. Næringsstoffer I plantedrikke (Nutrient content in plant-based beverages) DTU National Food Institute Lyngby, 2023. Available at: <https://www.food.dtu.dk/publikationer/ernaering-og-kostvaner>
- ³⁹⁴ Westhoek, H., Rood, T., Van De Berg, M., Janse, J., Nijdam, D., Reudink, M., & Stehfest, E. (2011). The protein puzzle: The consumption and production of meat, dairy and fish in the European Union. *The Protein Puzzle: The Consumption and Production of Meat, Dairy and Fish in the European Union*.
- ³⁹⁵ Source: FAOSTAT production data, world total 2019 [FAOSTAT](#)
- ³⁹⁶ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ³⁹⁷ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ³⁹⁸ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives.
- ³⁹⁹ Food balance data for 2019: [FAOSTAT](#)
- ⁴⁰⁰ Springmann, M., Clark, M., Mason-D'Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>
- ⁴⁰¹ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ⁴⁰² Source: FAOSTAT production data, world total 2019 [FAOSTAT](#)
- ⁴⁰³ [Palm Oil - Our World in Data](#)
- ⁴⁰⁴ Goldman, E., M.J. Weisse, N. Harris, and M. Schneider. 2020. "Estimating the Role of Seven Commodities in Agriculture-Linked Deforestation: Oil Palm, Soy, Cattle, Wood Fiber, Cocoa, Coffee, and Rubber." Technical Note. Washington, DC: World Resources Institute. Available online at: wri.org/publication/estimating-the-role-of-seven-commodities-in-agriculture-linked-deforestation
- ⁴⁰⁵ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ⁴⁰⁶ Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ⁴⁰⁷ [Palm Oil - Our World in Data](#)
- ⁴⁰⁸ Land use per crop assumes that all vegetable oil is produced from that crop alone. Source: [Palm Oil - Our World in Data](#)
- ⁴⁰⁹ Springmann, M., Clark, M., Mason-D'Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>
- ⁴¹⁰ Includes: Rape and mustard (majority), palm, soyabean, sunflower, olive, coconut, groundnut, sesame, and ricebran.
- ⁴¹¹ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ⁴¹² Includes: cream (majority), butter, fish body oil and fish liver oil.
- ⁴¹³ Source: FAOSTAT food balances available at [FAOSTAT](#)
- ⁴¹⁴ Source: FAOSTAT production data, world total 2019 [FAOSTAT](#)
- ⁴¹⁵ Goldman, E., M.J. Weisse, N. Harris, and M. Schneider. 2020. "Estimating the Role of Seven Commodities in Agriculture-Linked Deforestation: Oil Palm, Soy, Cattle, Wood Fiber, Cocoa, Coffee, and Rubber." Technical Note. Washington, DC: World Resources Institute. Available online at: wri.org/publication/estimating-the-role-of-seven-commodities-in-agriculture-linked-deforestation
- ⁴¹⁶ Source: Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216. Fig. S13. GHG emissions, acidification, and eutrophication by stage of the supply chain by product.
- ⁴¹⁷ Source: Poore, J. and Nemecek, T. (2018), 'Reducing food's environmental impacts through producers and consumers', *Science*, 360(6392): pp. 987–92, doi: 10.1126/scienceaaq0216.
- ⁴¹⁸ <https://ourworldindata.org/pollinator-dependence>
- ⁴¹⁹ Global Nutrition Report 2021: Country Nutrition Profiles. Bristol, UK: Development Initiatives.
- ⁴²⁰ Food supply quantity (per person per year) of sugar (raw equivalent) in 2019, from FBS data: [FAOSTAT](#)
- ⁴²¹ Springmann, M., Clark, M., Mason-D'Croz, D. et al. Options for keeping the food system within environmental limits. *Nature* 562, 519–525 (2018). <https://doi.org/10.1038/s41586-018-0594-0>
- ⁴²² <https://www.fao.org/fsnforum/index.php/resources/trainings-tools-and-databases/ultra-processed-foods-diet-quality-and-health-using-nova>; <https://www.fao.org/3/ca5644en/ca5644en.pdf>

- ⁴²³ Kliemann, N., Al Nahas, A., Vamos, E.P. et al. Ultra-processed foods and cancer risk: from global food systems to individual exposures and mechanisms. *Br J Cancer* 127, 14–20 (2022). <https://doi.org/10.1038/s41416-022-01749-y>; Lawrence, M.A. & Baker, P.I. 2019. Ultra-processed food and adverse health outcomes. *British Medical Journal*, 365:l2289. DOI: 10.1136/bmj.l2289.
- ⁴²⁴ <https://www.fao.org/3/ca5644en/ca5644en.pdf>
- ⁴²⁵ Seferidi et al. The neglected environmental impacts of ultra-processed foods. *The Lancet Planetary Health*, ISSN: 2542-5196, Vol: 4, Issue: 10, Page: e437-e438 2020 [http://doi.org/10.1016/s2542-5196\(20\)30177-7](http://doi.org/10.1016/s2542-5196(20)30177-7)
- ⁴²⁶ Leite, F., Khandpur, N., Andrade, G. C., Anastasiou, K., Baker, P., Lawrence, M., & Monteiro, C. A. (2022). Ultra-processed foods should be central to global food systems dialogue and action on biodiversity. *BMJ global health*, 7(3), e008269. <https://doi.org/10.1136/bmjgh-2021-008269>
- ⁴²⁷ Monteiro, C., Cannon, G., Moubarac, J., Levy, R., Louzada, M., & Jaime, P. (2018). The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutrition*, 21(1), 5-17. doi:10.1017/S1368980017000234
- ⁴²⁸ Leite, F., Khandpur, N., Andrade, G. C., Anastasiou, K., Baker, P., Lawrence, M., & Monteiro, C. A. (2022). Ultra-processed foods should be central to global food systems dialogue and action on biodiversity. *BMJ global health*, 7(3), e008269. <https://doi.org/10.1136/bmjgh-2021-008269>
- ⁴²⁹ Baker, P, Machado, P, Santos, T, et al. Ultra-processed foods and the nutrition transition: Global, regional and national trends, food systems transformations and political economy drivers. *Obesity Reviews*. 2020; 21:e13126. <https://doi.org/10.1111/obr.13126>;
- ⁴³⁰ Baker, P, Machado, P, Santos, T, et al. Ultra-processed foods and the nutrition transition: Global, regional and national trends, food systems transformations and political economy drivers. *Obesity Reviews*. 2020; 21:e13126. <https://doi.org/10.1111/obr.13126>; <https://www.who.int/publications/i/item/9789240073616>; <https://apps.who.int/iris/rest/bitstreams/1501485/retrieve>
- ⁴³¹ Analysis includes all Nordic and Baltic countries except Iceland. Source: Baker, P, Machado, P, Santos, T, et al. Ultra-processed foods and the nutrition transition: Global, regional and national trends, food systems transformations and political economy drivers. *Obesity Reviews*. 2020; 21:e13126. <https://doi.org/10.1111/obr.13126>
- ⁴³² Seferidi et al. The neglected environmental impacts of ultra-processed foods. *The Lancet Planetary Health*, ISSN: 2542-5196, Vol: 4, Issue: 10, Page: e437-e438 2020 [http://doi.org/10.1016/s2542-5196\(20\)30177-7](http://doi.org/10.1016/s2542-5196(20)30177-7)
- ⁴³³ Hendrie, Gilly A., Danielle Baird, Brad Ridoutt, Michalis Hadjikakou, and Manny Noakes. 2016. "Overconsumption of Energy and Excessive Discretionary Food Intake Inflates Dietary Greenhouse Gas Emissions in Australia" *Nutrients* 8, no. 11: 690. <https://doi.org/10.3390/nu8110690>
- ⁴³⁴ Aceves-Martins M, Bates RL, Craig LCA, Chalmers N, Horgan G, Boskamp B, de Roos B. Nutritional Quality, Environmental Impact and Cost of Ultra-Processed Foods: A UK Food-Based Analysis. *International Journal of Environmental Research and Public Health*. 2022; 19(6):3191. <https://doi.org/10.3390/ijerph19063191>
- ⁴³⁵ Seferidi et al. The neglected environmental impacts of ultra-processed foods. *The Lancet Planetary Health*, ISSN: 2542-5196, Vol: 4, Issue: 10, Page: e437-e438 2020 [http://doi.org/10.1016/s2542-5196\(20\)30177-7](http://doi.org/10.1016/s2542-5196(20)30177-7)
- ⁴³⁶ Leite, F., Khandpur, N., Andrade, G. C., Anastasiou, K., Baker, P., Lawrence, M., & Monteiro, C. A. (2022). Ultra-processed foods should be central to global food systems dialogue and action on biodiversity. *BMJ global health*, 7(3), e008269. <https://doi.org/10.1136/bmjgh-2021-008269>
- ⁴³⁷ Benton et al 2021. [Food system impacts on biodiversity loss | Chatham House – International Affairs Think Tank](#)
- ⁴³⁸ Leite, F., Khandpur, N., Andrade, G. C., Anastasiou, K., Baker, P., Lawrence, M., & Monteiro, C. A. (2022). Ultra-processed foods should be central to global food systems dialogue and action on biodiversity. *BMJ global health*, 7(3), e008269. <https://doi.org/10.1136/bmjgh-2021-008269>
- ⁴³⁹ Gunnarsdottir S, Arnardottir R, Torfadottir JE, Thorgeirsdottir H, Geirsdottir OG, Gudjonsdottir M, Ogmundarson O, Halldorsson ThI, Gunnarsdottir I, Birgisdottir BE. Ultra-processed food consumption in Iceland. Abstract presented at the 14th European Nutrition Conference FENS, 2023, Belgrade, Serbia.
- ⁴⁴⁰ Please note: This information was provided by regional experts during a series of workshops and has not been verified.
- ⁴⁴¹ Monteiro, C., Cannon, G., Moubarac, J., Levy, R., Louzada, M., & Jaime, P. (2018). The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutrition*, 21(1), 5-17. doi:10.1017/S1368980017000234
- ⁴⁴² Trolle et al 2023 Integrating environmental sustainability into Food-Based Dietary Guidelines in the Nordic countries (in press).
- ⁴⁴³ Muller, A., Schader, C., El-Hage Scialabba, N., Brüggemann, J., Isensee, A., Erb, K.-H., Smith, P., Klocke, P., Liebe, F., Stolze, M. and Niggli, U. (2017), 'Strategies for feeding the world more sustainably with organic agriculture', *Nature Communications*, 8(1290), doi: 10.1038/s41467-017-01410-w; Benton, T. G., Bieg, C., Harwatt, H., Pudasaini, R., & Wellesley, L. (2021). Food system impacts on biodiversity loss. *Three levers for food system transformation in support of nature*; Benton, Tim G., and Helen Harwatt. "Sustainable agriculture and food systems." (2022). *Chatham House, London*; Billen, G., Aguilera, E., Einarsson, R., Garnier, J.,

Gingrich, S., Grizzetti, B., Lassaletta, L., Le Noë, J. and Sanz-Cobena, A., Reshaping the European agro-food system and closing its nitrogen cycle: the potential of combining dietary change, agroecology, and circularity, ONE EARTH, 2021, ISSN 2590-3322 (online),2590-3330 (print), 4 (6), p. 839-850, JRC123720.

⁴⁴⁴ Benton, T et al (2023) Forthcoming. Assessing the environmental sustainability of diets – an overview of approaches and identification of 5 key considerations for comprehensive assessments.

⁴⁴⁵ Harwatt, H., Wetterberg, K., Giritharan, A. and Benton, T. G. (2022), Aligning food systems with climate and biodiversity targets: Assessing the suitability of policy action over the next decade, Research Paper, London: Royal Institute of International Affairs, <https://doi.org/10.55317/9781784135416>.