



## First-Principles Based Kinetic Analysis of the Electrochemical Discharge of Water on Pt3M Skin Alloys

Viswanathan, Venkatasubramanian; Hansen, Heine Anton; Rossmeisl, Jan; Jaramillo, Thomas F.; Nørskov, Jens Kehlet; Pitsch, Heinz

*Published in:*  
Meeting Abstracts - Electrochemical Society

*Publication date:*  
2009

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Viswanathan, V., Hansen, H. A., Rossmeisl, J., Jaramillo, T. F., Nørskov, J. K., & Pitsch, H. (2009). First-Principles Based Kinetic Analysis of the Electrochemical Discharge of Water on Pt3M Skin Alloys. In *Meeting Abstracts - Electrochemical Society* (pp. Abstract 964). The Electrochemical Society.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

First-principles based kinetic analysis of the electrochemical discharge of water on Pt<sub>3</sub>M skin alloys  
Venkatasubramanian Viswanathan<sup>1</sup>, Heine Anton Hansen<sup>2</sup>, Jan Rossmeisl<sup>2</sup>, Thomas Jaramillo<sup>3</sup>, Jens Kehlet Nørskov<sup>2</sup>, Heinz Pitsch<sup>1</sup>

<sup>1</sup>Department of Mechanical Engineering, Stanford University, Stanford 94305

<sup>2</sup>Center for Atomic Scale Material Design, Department of Physics, Technical University of Denmark

<sup>3</sup>Department of Chemical Engineering, Stanford University

Polymer electrolyte membrane fuel cells (PEMFC) are promising energy conversion devices in which the chemical energy of low-carbon or zero carbon fuels (e.g. methanol or hydrogen) is converted to electricity at low temperature. A major obstacle for wide-scale deployment of PEMFC technology is the development of active, inexpensive, and stable catalysts for the oxygen-reduction reaction (ORR). Platinum and Pt-based alloys currently set the standard for ORR catalysts as they are active and stable, though there is room for improvement on both issues. Pt-skin alloys of Pt<sub>3</sub>M are particularly interesting; recent studies[1-2] have shown that Pt-skinned Pt<sub>3</sub>M alloys have shown improved activity for oxygen reduction reaction compared to pure Pt or bulk Pt alloys. As Pt skin alloys (Pt<sub>3</sub>M) consist of a pure Pt surface, they are in many ways similar to pure Pt, but the subtle changes in electronic and geometric structure arising from the alloyed second or third atomic layers with a transition metal M give rise to a significant improvement in ORR catalytic activity.

A detailed understanding of the relevant intermediates of ORR on these surfaces can aid in the design of better catalysts. Here, we present a first-principles based kinetic-analysis of the electrochemical discharge of water on Pt skin alloys.

Adsorbate-adsorbate interactions during relevant ORR conditions play a large role in ORR activity. The approach presented in this paper is based on a micro-kinetic micro-dynamics model which uses energies computed using Density Functional theory (DFT). DFT-calculated angular resolved adsorbate-adsorbate interactions account for the complex adsorbate structure and the non-linearity of the adsorbate interactions. These interactions are used in combination of equilibrium Monte Carlo and dynamic Monte Carlo to perform first principle simulations of the electrochemical discharge of water. The two electron transfer steps  $\text{H}_2\text{O} \leftrightarrow \text{OH} + \text{H}^+ + \text{e}^-$  and  $\text{OH} \leftrightarrow \text{O} + \text{H}^+ + \text{e}^-$  are treated kinetically while the fast steps like diffusion, rotation of adsorbed species are treated in equilibrium. The kinetics of the electron transfer reactions is computed using Local Reaction Center theory[3], as implemented in the Convergent Iterative Constrained Variation (CICV) method[4]. The effects of solvation, the electric field and entropy are accounted for by shifting the activation energy curves to their thermodynamic reversible potential computed from DFT[5].

The results of the adsorbate coverage from the electrochemical discharge of water on Pt<sub>3</sub>Ni(111) surface in 0.1 M HClO<sub>4</sub> solution is shown in Fig 1. Similar results will be presented for other Pt skin alloys (Pt<sub>3</sub>Fe, Pt<sub>3</sub>Ti). The results show that there is a coexistence of OH<sub>(ads)</sub> and O<sub>(ads)</sub> at high potentials. These adsorbates

adsorb on on-top and fcc sites, respectively, resulting in an adsorbate surface structure that impacts the nature of the ORR reaction; motivating the need for a theoretical framework that accurately describes adsorbate structure – not only to understand ORR kinetics on different surfaces but also to design improved materials.




Fig 1: Results of the simulation of the electrochemical discharge of water on Pt<sub>3</sub>Ni(111) in 0.1M HClO<sub>4</sub>.

## References

- [1] V. R. Stamenkovic, B. Fowler, B. S. Mun, G. Wang, P. N. Ross, C. A. Lucas, and N. M. Markovic. Improved Oxygen Reduction Activity on Pt<sub>3</sub>Ni(111) via Increased Surface Site Availability. *Science*, 315(5811):493–497, 2007.
- [2] V. R. Stamenkovic, B. S. Mun, M. Arenz, K.J.J. Mayrhofer, C. A. Lucas, G. Wang, P. N. Ross and N.M. Markovic, Trends in electrocatalysis on extended and nanoscale Pt-bimetallic alloy surfaces. *Nature Materials*, 6:241–247, 2007.
- [3] A.B.Anderson and T.V.Albu, Ab Initio Determination of Reversible Potentials and Activation Energies for Outer-Sphere Oxygen Reduction to Water and the Reverse Oxidation Reaction, *J. Am. Chem. Soc.*, 121(50):11855–11863, 1999.
- [4] M.Aryanpour, V.Rai, and H.Pitsch, Convergent Iterative Constraint Variation Algorithm for Calculation of Electron-Transfer Transition States, *J. Electrochem. Soc.*, 153:E52-E57, 2006.
- [5] J. Rossmeisl, G. S. Karlberg, T. Jaramillo and J. K. Nørskov, Steady state oxygen reduction and cyclic voltammetry, *Faraday Discussions*, 140: 337-346, 2008.