



Assessment of Lead Discrimination from CryoSat-2

Kildegaard Rose, Stine; Connor, L. N.; Newman, T.; Farrell, S. L.; Smith, W. H.; Forsberg, René

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Kildegaard Rose, S., Connor, L. N., Newman, T., Farrell, S. L., Smith, W. H., & Forsberg, R. (2012). *Assessment of Lead Discrimination from CryoSat-2*. Abstract from 2012 AGU Fall Meeting, San Francisco, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

2012 AGU Fall meeting, 3-8 dec.

Abstract number:

C51A-0763

Title:

Assessment of Lead Discrimination from CryoSat-2

Authors:

Stine K. Rose¹, Laurence N. Connor², Thomas Newman^{2,3}, Sinead L. Farrell^{2,3}, Walter H. Smith², Rene Forsberg¹

1. Geodynamics, DTU Space, Technical University of Denmark, Kgs. Lyngby, Denmark, 2. Laboratory for Satellite Altimetry, NOAA, College Park, MD, USA, 3. Cooperative Institute for Climate and Satellites, Earth system Science Interdisciplinary Center (ESSIC), University of Maryland, College Park, MD, USA

Abstract:

Sea ice is strongly affecting the global climate, and the sea ice extent has been monitored by satellites since 1979. To estimate the Arctic sea ice volume, ice thickness must be determined. The measurements of sea ice thickness are however more difficult to achieve, and encounter limitations due to spatial and temporal variability. The measurements of sea ice freeboard may be used to estimate sea ice thickness, when combined with examination of leads between ice floes to determine the local sea surface height. With CryoSat-2 (CS), we have the opportunity to measure much more of the Arctic Ocean due to its high sampling rate and geographical coverage to 88 °N/S. Validation of the CS retrievals are very important to verify the derived sea ice thickness and understand the associated error sources. We present a comparative analysis of CryoSat-2 elevations with the Operation IceBridge Airborne Topographic Mapper (ATM) laser altimeter data gathered on April 2, 2012, where the NASA P-3 completed an underflight of CS orbit number 10520, north of Alert, Nunavut, Canada. We present a new lead detecting algorithm which was developed using the CS Level1b (L1b) waveforms, and we analyze its capabilities via comparisons with IceBridge imagery and ATM elevations. In addition, using CS L1b waveforms we have developed a method to find misplaced CS Level 2 elevations and correct them to remove any elevation bias.