



Exploring methods for combining altimetry with other data to extend the 20-year altimetric record onto a 50 year timescale

Svendsen, Peter Limkilde; Andersen, Ole Baltazar; Nielsen, Allan Aasbjerg; Knudsen, Per

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Svendsen, P. L., Andersen, O. B., Nielsen, A. A., & Knudsen, P. (2012). *Exploring methods for combining altimetry with other data to extend the 20-year altimetric record onto a 50 year timescale*. Abstract from 2012 AGU Fall Meeting, San Francisco, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CONTROL ID: 1494218

TITLE: Exploring methods for combining altimetry with other data to extend the 20-year altimetric record onto a 50 year timescale

AUTHORS (FIRST NAME, LAST NAME): Peter Limkilde Svendsen¹, Ole Baltazar Andersen¹, Allan Aasbjerg Nielsen¹, Per Knudsen¹

INSTITUTIONS (ALL): 1. DTU Space, Kgs. Lyngby, Denmark.

ABSTRACT BODY: Ocean satellite altimetry has provided global sets of sea level data for the last two decades, allowing determination of spatial patterns in global sea level. For reconstructions going back further than this period, tide gauge data can be used as a proxy for the model. We examine different methods of combining satellite altimetry and tide gauge data using optimal weighting of tide gauge data, linear regression and EOFs, including automatic quality checks of the tide gauge time series.

We attempt to test the sensibility of reconstruction using known and existing datasets and to test the important of augmenting the model using various proxies such as climate indices like the NAO and PDO. We will also investigate alternative transformations such as maximum autocorrelation factors (MAF), which better take into account the spatio-temporal structure of the variation. Whereas a traditional EOF analysis tries to explain as much variance as possible, the MAF transform considers noise to be uncorrelated with a spatially or temporally shifted version of itself, unlike the desired signal which will exhibit autocorrelation. For the application to global dataset it is necessary to consider and account for wrap-around of spatial shifts. Parameters from physical oceanography will be incorporated using ocean models (i.e., DRAKKAR; SODA) for a preliminary reference. Our focus is on a timescale going back approximately 50 years, allowing reasonable global availability of model and tide gauge data. This allows for better sensitivity analysis with respect to spatial distribution, and tide gauge data.

KEYWORDS: [1641] GLOBAL CHANGE / Sea level change, [1637] GLOBAL CHANGE / Regional climate change.

(No Image Selected)

(No Table Selected)

Additional Details

Previously Presented Material:

Contact Details

CONTACT (NAME ONLY): Peter Svendsen

CONTACT (E-MAIL ONLY): psv@space.dtu.dk

TITLE OF TEAM:
