



## Nucleation in an ultra low ionization environment

Pedersen, Jens Olaf Pepke; Enghoff, Martin Andreas Bødker; Paling, Sean; Svensmark, Henrik

*Publication date:*  
2010

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Pedersen, J. O. P., Enghoff, M. A. B., Paling, S., & Svensmark, H. (2010). *Nucleation in an ultra low ionization environment*. Abstract from European Geosciences Union General Assembly 2010, Vienna, Austria.  
<http://adsabs.harvard.edu/abs/2010EGUGA..1210661O>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **Nucleation in an ultra low ionization environment**

Jens Olaf Pepke Pedersen<sup>1</sup>, Martin Bødker Enghoff<sup>1</sup>, Sean Paling<sup>2</sup>, and Henrik Svensmark<sup>1</sup>

(1) National Space Institute, Technical University of Denmark, Copenhagen, Denmark ,  
(2) Department of Physics and Astronomy, Sheffield University, Sheffield, U.K.

Atmospheric ions can enhance the nucleation of aerosols, as has been established by experiments, observation, and theory. In the clean marine atmosphere ionization is mainly caused by cosmic rays which in turn are controlled by the activity of the Sun, thus providing a potential link between solar activity and climate. In order to understand the effect ions may have on the production of cloud condensation nuclei the overall contribution of ion induced nucleation to the global production of secondary aerosols must be determined. One issue with determining this contribution is that several mechanisms for nucleation exist and it can be difficult to determine the relative importance of the various mechanisms in a given nucleation event when both ion induced and electrically neutral nucleation mechanisms are at work at the same time.

We have carried out nucleation experiments in the Boulby Underground Laboratory, located 1100 meters below ground, thus reducing the flux of ionizing cosmic radiation by six orders of magnitude. Similarly we have reduced the gamma background by shielding the experiment in lead and copper. Finally we have used air stored for several weeks and passed through an active charcoal filter in order to reduce the Radon concentration. In this way we have been able to make nucleation experiments with very low ionizing background, meaning that we can rule out ion induced nucleation as a contributing mechanism.

Our experimental setup is a 50 L electropolished stainless steel reactor at near atmospheric conditions. The chamber contains clean air with the addition of water vapour, ozone, and SO<sub>2</sub>. Using UV lights at 254 nm ozone is photolyzed, leading to the production of sulphuric acid and thus aerosols. An 18 MBq Caesium-137 gamma ray source with various amounts of lead in front allows us to alter the ionization in our chamber. By making series of nucleation bursts with varying amounts of ionizing radiation we then gauge the relative importance of ion induced nucleation in the events. This set of experiments is, to our knowledge, the first of its kind.