



Signals from beech (*Fagus sylvatica* L.) in response to precipitation extremes - flowering induction and reduced foliation

Callesen, Ingeborg

Publication date:
2010

[Link back to DTU Orbit](#)

Citation (APA):

Callesen, I. (2010). *Signals from beech (Fagus sylvatica L.) in response to precipitation extremes - flowering induction and reduced foliation*. Poster session presented at CARBO Extreme project meeting 2010, Roskilde, Denmark, .

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Signals from beech (*Fagus sylvatica* L.) in response to precipitation extremes – flowering induction and reduced foliation

Ingeborg Callesen^{1,2}

¹DTU Management Engineering, Section for Quantitative Sustainability Assessment

² Biosystems division, Risø DTU, Technical University of Denmark, Frederiksborgvej 399, 4000 Roskilde, Denmark. inca@risoe.dtu.dk.

Abstract :

Reduced foliation in older (but also young) beech (*Fagus sylvatica* L.) stands was observed in Denmark in the mid 1990ies and culminated with the 1996 summer drought and heat wave. Large differences in the degree of reduced foliation between regions and within stands were observed e.g. reflecting stand structure and exposure.

An analysis of monthly precipitation and temperature data from 1873-1996 in south-eastern Denmark revealed that the series of summers with low precipitation in 1994-1996 was rare but not unprecedented (Figure 1).

The precipitation data were compared with a data series of mast years (classes ranging from 1~no mast to 4~maximum mast production, Figure 2) showing that widespread flowering in beech occurred when the summer in the previous year had been warm and dry (Figure 3), and when a favorable growing season occurred two years prior to the mast year in terms of higher than average precipitation and lower than average temperature during June, July and August.

The different response could also be related to site factors that prevented deep and permanent rooting due to occasional lack of soil aeration. The permanent rooting depth should be devoid of signs of stagnant water (gley and severe pseudogley) caused by poor internal drainage (Figure 4) and minor depressions in micro relief.




Figure 1. Summer precipitation (may-aug), 3-yr averages in three regions of Denmark. Two extreme series of dry summers occurred in the mid 70ies and mid 90ies.




Figure 2. Mast production from the year 1832 to 1996. 1~no seed production, 2, 3.. to 4~maximum mast (seed) production.




Figure 3. Mean monthly temperature one (N-1) and two (N-2) years prior to mast year in June, July and August. Four mast classes: 1~no mast, 2~scarce mast 3~abundant mast, 4~maximum mast. *~significant correlation with mast class ($P<0.05$).




Figure 4. Beech stand with reduced foliation on clay soil with poor internal drainage (clay soil with pseudogley and gley).