



EU 1902+20 and EU 2017-01

Brandt, Søren; Castro-Tirado, A. J.; Lund, Niels

Published in:
International Astronomical Union Circulars (IAUC)

Publication date:
1992

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Brandt, S., Castro-Tirado, A. J., & Lund, N. (1992). EU 1902+20 and EU 2017-01. *International Astronomical Union Circulars (IAUC)*, (5659). <http://www.cbat.eps.harvard.edu/iauc/05600/05659.html#Item1>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Central Bureau for Astronomical Telegrams
INTERNATIONAL ASTRONOMICAL UNION
Postal Address: Central Bureau for Astronomical Telegrams
Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
Telephone 617-495-7244/7440/7444 (for emergency use only)
TWX 710-320-6842 ASTROGRAM CAM EASYLINK 62794505
MARSDEN@CFA or GREEN@CFA (.SPAN, .BITNET or .HARVARD.EDU)

EU 1902+20 AND EU 2017-01

S. Brandt, A. J. Castro-Tirado, and N. Lund, Danish Space Research Institute, communicate: "Two new x-ray transients, located at R.A. = 19h02m, Decl. = +20.0 and R.A. = 20h17m, Decl. = -1.0 (equinox 1950.0; estimated error radius 1 deg), have been discovered with the WATCH wide-field monitor on the European Space Agency's EURECA satellite. The new transients were detected in data from Nov. 22, but both appear at a lower level of significance already on Nov. 21. EU 2017-01 has a hard spectrum, whereas EU 1902+20 has only been detected between 6 and 8 keV, the flux from both sources being of the order 100 mCrab. Follow-up at other wavelengths is encouraged."

PERIODIC COMET SWIFT-TUTTLE (1992t)

J. Davies, Royal Observatory, Edinburgh, reports: "A 3.2- to 3.6-micron spectrogram of P/Swift-Tuttle was taken at the U.K. Infrared Telescope on Nov. 13.15 UT by S. Lumsden on behalf of Davies, M. Mumma, S. Hoban, H. Weaver, and collaborators. Preliminary reduction of the spectrum shows a broad emission feature between 3.3 and 3.45 microns. There may be substructure within this feature, most notably appearing as a local peak near 3.427 microns. There is a broad emission feature centered at 3.52 microns, with no evidence of an emission feature at 3.28 microns. Detailed modeling of fluorescence from methanol produces a spectrum that explains all of the flux from 3.3 to 3.4 microns, and also that of the 3.52-micron feature. Excess flux is seen from 3.4 to 3.48 microns, probably due to another species. The 3.52-micron feature (which we attribute to methanol) has a flux of $8 \times 10E-16$ W mE-2 summed over an area $3".1 \times 6".2$, implying a methanol production rate of $1.5 \times 10E28$ molecules sE-1. The flux in the 3.3- to 3.45-micron region is about $6 \times 10E-15$ W mE-2 in excess of the underlying continuum. The underlying continuum has a flux of $1.15 \times 10E-13$ W mE-2 micronE-1 near 3.25 microns, rising to $1.4 \times 10E-13$ near 3.58 microns."

NOVA IN THE LARGE MAGELLANIC CLOUD 1992

Photometry by A. C. Gilmore with the 0.6-m f/14 Cassegrain reflector at Mt. John University Observatory: Nov. 21.58 UT, V = 12.32 +/- 0.06, U-B = -0.86 +/- 0.07, B-V = +0.09 +/- 0.06, V-R = +0.73 +/- 0.07; Nov. 22.48, 12.39 +/- 0.02, -0.74 +/- 0.02, +0.05 +/- 0.01, +0.74 +/- 0.01.