



## X-Ray Activity in Cygnus and Aquila

**Brandt, Søren; Castro-Tirado, A. J.; Lund, Niels**

*Published in:*  
International Astronomical Union Circulars (IAUC)

*Publication date:*  
1992

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Brandt, S., Castro-Tirado, A. J., & Lund, N. (1992). X-Ray Activity in Cygnus and Aquila. *International Astronomical Union Circulars (IAUC)*, (5664). <http://www.cbat.eps.harvard.edu/iauc/05600/05664.html#Item1>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Central Bureau for Astronomical Telegrams  
INTERNATIONAL ASTRONOMICAL UNION  
Postal Address: Central Bureau for Astronomical Telegrams  
Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.  
Telephone 617-495-7244/7440/7444 (for emergency use only)  
TWX 710-320-6842 ASTROGRAM CAM EASYLINK 62794505  
MARSDEN@CFA or GREEN@CFA (.SPAN, .BITNET or .HARVARD.EDU)

**X-RAY ACTIVITY IN CYGNUS AND AQUILA**

S. Brandt, A. J. Castro-Tirado and N. Lund, Danish Space Research Institute, report: "A new x-ray transient, EU 2053+316, has been discovered with the WATCH wide field monitor on the ESA EURECA satellite.

First seen on Nov. 23, it has persisted since, but with varying intensity. The position (R.A. = 20h53m, Decl = +31.6 deg, equinox 1950.0; estimated error radius 1 deg) is inside the Cygnus Loop and near the bright northwestern rim as seen in the Einstein image of the Loop. Cyg X-2 flared on Nov. 14 and reached an intensity of 500 mCrab above 6 keV on Nov. 15, while Cyg X-3 flared to 300 mCrab on Nov. 17; both sources remain above the 100 mCrab level. The recurrent x-ray transient Aql X-1 (V1333 Aql) has been active since Nov. 20. GRS 1915+105 ([IAUC 5590](#)) has been monitored since the end of August and is still active and observed above 15 keV."

**PERIODIC COMET SWIFT-TUTTLE (1992t)**

G. Paubert, Institut de Radio Astronomie Millimetrique; P. Colom, D. Bockelee-Morvan, J. Crovisier and L. Jorda, Observatoire de Paris-Meudon; and D. Despois, Observatoire de Bordeaux, report: "The millimeter spectrum of this comet was observed on Nov. 21 with the IRAM 30-m telescope. The following lines were detected (signal-to-noise ranging from 7 to 30), preliminary evaluations of the corresponding molecular production rates being given in units of  $10^{27}$  s<sup>-1</sup>: HCN J(1-0), 89 GHz, 0.25; CH<sub>3</sub>OH, 12 lines around 145 and 165 GHz, 20; H<sub>2</sub>S 1(1,0)-1(0,1), 168.8 GHz, 1.5; H<sub>2</sub>CO 3(1,2)-2(1,1), 225.7 GHz, 2 (from a column density of  $6 \times 10^{12}$  cm<sup>-2</sup> for a 13" HPBW and assuming a parent-molecule distribution). All line profiles are asymmetric with a cusp at negative velocities. The production rates of HCN and CH<sub>3</sub>OH given on [IAUC 5653](#) were underestimated due to ephemeris uncertainties."

Jorda and J. Lecacheux, Paris-Meudon; and F. Colas, Observatoire de Paris, report: "Observations with the 1.05-m telescope and CCD camera at Pic du Midi during Nov. 20-26 showed a strong helicoidal jet extending from 1500 to 20 000 km. The jet rotates from one night to the next, and its motion is clearly seen between exposures taken over a 2-hr interval. Our observations are compatible with a nuclear rotation period of about 2.9 days and an emission from a high-latitude active zone. From the observations in 1862 Z. Sekanina (1981, A.J. 86, 1741) deduced a period of 2.77 days."