



The influence of digestibility on the allergenicity of food allergens

Bøgh, Katrine Lindholm; Madsen, Charlotte Bernhard

Published in:

The 7th meeting of the Immunotoxicology and Chemical Allergy Speciality Section ITCASS

Publication date:

2012

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Bøgh, K. L., & Madsen, C. B. (2012). The influence of digestibility on the allergenicity of food allergens. In *The 7th meeting of the Immunotoxicology and Chemical Allergy Speciality Section ITCASS* (pp. 9-9). Technical University of Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The influence of digestibility on the allergenicity of food allergens

Katrine L. Bøgh & Charlotte B. Madsen

National Food Institute, Division of Toxicology and Risk Assessment, Technical University of Denmark, Søborg, Denmark

Food allergy is a major health problem in the Western countries, affecting 3-8% of the population. What makes a dietary protein a food allergen has not yet been established, though several characteristics have been proposed to be shared by the food allergens. One of the features believed to be a general characteristic is resistance to digestion. This is based on studies showing that allergenic dietary proteins in general were more resistant to digestion than dietary proteins with no proven allergenicity, leading to the conclusion, that a correlation between stability to digestion and allergenic potential exist. Resistance to digestion is therefore a test parameter included in the safety assessment of the allergenic potential of novel proteins in genetically modified foods. In recent years, the association between resistance to digestion and allergenic potential has been challenged.

When reviewing existing data from digestibility studies on known food allergens, it becomes evident that food allergens do not necessarily resist digestion. However, the choice of assay conditions, the method used for detection of residual intact protein as well as fragments hereof greatly influences the outcome. Studies assessing the allergenicity of digestion products, by either IgE-binding, elicitation or sensitising capacity, shows that digestion may abolish, decrease, have no effect, or even increase the allergenicity of food allergens. However, this depends on the given allergen.

In conclusion, reviewing existing digestibility data shows that no absolute correlation between resistance to digestion and allergenic potential exist. Therefore stability to digestion may not necessarily be a good parameter for assessing the allergenic potential of novel proteins. Even very small peptides from food allergens may retain both IgE-binding, eliciting and sensitising capacity. As a consequence immunological studies should be performed when evaluating the digestibility of protein allergens.