



Improving volume loss estimate of the Greenland Ice Sheet between

Khan, Shfaqat Abbas; Wahr, John

Published in:
Geophysical Research Abstracts

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Khan, S. A., & Wahr, J. (2013). Improving volume loss estimate of the Greenland Ice Sheet between. *Geophysical Research Abstracts*, 15, EGU2013-2725-1.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Improving volume loss estimate of the Greenland Ice Sheet between 2002-2012

Shfaqat Abbas Khan (1) and John Wahr (2)

(1) DTU - Space, National Space Institute, Geodesy, Kgs. Lyngby, Denmark (abbas@space.dtu.dk, +45 459775), (2) Department of Physics and Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, USA

Ice loss from the Greenland Ice Sheet is dominated by loss in the marginal areas. Dynamic induced ice loss and its associated ice surface lowering is often largest close to the glacier calving front and may vary from rates of tens of meters per years to a few meters per year over relatively short distances. Hence, high spatial resolution data are required to accurately estimate volume changes. We estimate ice volume change rate of the Greenland ice sheet during 2003-2009 using Ice, Cloud and land Elevation Satellite (ICESat) laser altimeter data (Zwally et al., 2011). To improve the volume change estimate we supplement ICESat data with altimeter surveys from NASA's Airborne Topographic Mapper (ATM) during 2002-2010 (Krabill et al., 2011) and NASA's Land, Vegetation and Ice Sensor (LVIS) during 2007-2010 (Blair and Hofton, 2010). The Airborne data are mainly concentrated along the ice margin and therefore significantly improve the estimate of the total volume change. Our results show that adding ATM and LVIS data to the ICESat data increases the estimate of catchment-wide ice volume loss in northeast and southeast Greenland by 10-20 percent. This is mainly due to increased volume loss near the margin of the GrIS. Furthermore, we provide volume loss estimates during 2002-2009 and 2009-2012. Our 2009-2012 estimate suggest enhanced volume loss at the entire southern half of the GrIS.