



## 3D velocity structure of upper crust beneath NW Bohemia/Vogtland

Javad Fallahi, Mohammad; Mousavi, Sima; Korn, Michael ; Sens-Schönfelder, Christoph; Bauer, Klaus; Rössler, Dirk

*Published in:*  
Geophysical Research Abstracts

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Javad Fallahi, M., Mousavi, S., Korn, M., Sens-Schönfelder, C., Bauer, K., & Rössler, D. (2013). 3D velocity structure of upper crust beneath NW Bohemia/Vogtland. *Geophysical Research Abstracts*, 15, EGU2013-8678.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## **3D velocity structure of upper crust beneath NW Bohemia/Vogtland**

Mohammad Javad Fallahi (1), Sima Mousavi (1), Michael Korn (1), Christoph Sens-Schönfelder (2), Klaus Bauer (2), and Dirk Rößler (3)

(1) Institute for Geophysics and Geology, Leipzig University, Leipzig, Germany (fallahi@uni-leipzig.de), (2) GFZ German Research Centre for Geosciences, Potsdam, Germany, (3) DTU Space, Technical University of Denmark, Copenhagen, Denmark

The 3D structure of the upper crust beneath west Bohemia/Vogtland region, analyzed with travel time tomography and ambient noise surface wave tomography using existing data. This region is characterized by a series of phenomena like occurrence of repeated earthquake swarms, surface exhalation, CO<sub>2</sub> enriched fluids, mofettes, mineral springs and enhanced heat flow, and has been proposed as an excellent location for an ICDP drilling project targeted to a better understanding of the crust in an active magmatic environment.

We performed a 3D tomography using P- and S-wave travel times of local earthquakes and explosions. The data set were taken from permanent and temporary seismic networks in Germany and Czech Republic from 2000 to 2010, as well as active seismic experiments like Celebration 2000 and quarry blasts. After picking P and S wave arrival times, 399 events which were recorded by 9 or more stations and azimuthal gap < 160° were selected for inversion.

A simultaneous inversion of P and S wave 1D velocity models together with relocations of hypocenters and station corrections was performed. The obtained minimum 1D velocity model was used as starting model for the 3D V<sub>p</sub> and V<sub>p</sub>/V<sub>s</sub> velocity models. P and S wave travel time tomography employs damped least-square method and ray tracing by pseudo-bending algorithm. For model parametrization different cell node spacings have been tested to evaluate the resolution in each node. Synthetic checkerboard tests have been done to check the structural resolution. Then V<sub>p</sub> and V<sub>p</sub>/V<sub>s</sub> in the preferred 3D grid model have been determined. Earthquake locations in iteration process change till the hypocenter adjustments and travel time residuals become smaller than the defined threshold criteria. Finally the analysis of the resolution depicts the well resolved features for interpretation. We observed lower V<sub>p</sub>/V<sub>s</sub> ratio in depth of 5-10 km close to the foci of earthquake swarms and higher V<sub>p</sub>/V<sub>s</sub> ratio is observed in Saxothuringian zone and surrounding area.

Surface wave tomography using ambient noise provides additional constraints on shear velocities. The detailed knowledge of the 3D structure is essential to select the optimal future borehole locations. We use the vertical and transverse component ambient noise data to estimate both Rayleigh and Love waves from ambient noise cross-correlation waveforms to investigate the crustal seismic structure of W-Bohemia/Vogtland.

More than 2000 Rayleigh and Love group-velocity dispersion curves are obtained by time-frequency analysis of stacked ambient noise cross-correlation functions between station pairs. We used the data between 2002 and 2004 recorded at 43 seismic stations from BOHEMA experiment and between 2006 and 2008 recorded at 79 seismic stations from permanent station networks of Germany, Czech Academy of Sciences (WEBNET) and PASSEQ experiments.

At each period between 1 and 10 s, group velocity maps are constructed, all corresponding to different sampling depths, and thus together giving an indication of the 3D shear wave velocity structure extending to a depth of about 15 km.