



Quantifying the effect of natural microflora on growth of *Salmonella* Typhimurium DT104 and *Salmonella* Derby in fresh pork

Møller, Cleide; Ilg, Yvonne; Aabo, Søren; Dalgaard, Paw; Christensen, Bjarke Bak; Hansen, Tina Beck

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Møller, C., Ilg, Y., Aabo, S., Dalgaard, P., Christensen, B. B., & Hansen, T. B. (2011). *Quantifying the effect of natural microflora on growth of Salmonella Typhimurium DT104 and Salmonella Derby in fresh pork*. Poster session presented at 9th International Conference on the Epidemiology and Control of Biological Chemical and Physical Hazards in Pigs and Pork, Maastricht, Netherlands.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Title

Quantifying the effect of natural microflora on growth of *Salmonella* Typhimurium DT104 and *Salmonella* Derby in fresh pork

Cleide O.A. Møller¹, Yvonne Ilg², Søren Aabo¹, Paw Dalgaard³, Bjarke B. Christensen⁴, Tina B. Hansen¹

¹National Food Institute, Technical University of Denmark, Mørkhøj Bygade 19, DK-2860 Søborg, Denmark

²Institute of Animal Science, Department of Preventive Health Management, University of Bonn, Katzenburgweg 7-9, 53115 Bonn, Germany

³National Food Institute, Technical University of Denmark, Sølvtofts Plads, Building 221, DK-2800 Kgs. Lyngby, Denmark

⁴Department of Food Science, Faculty of Life Sciences, University of Copenhagen, Rolighedsvej 30, DK-1958 Frederiksberg C., Denmark

Objective


To build predictive growth rate models for *Salmonella* Typhimurium DT104 and *Salmonella* Derby in fresh pork in the temperature area between 4 and 20°C using

- sterile meat (irradiated at 5 kGy for 523 min)
- meat with a natural microflora


Conclusion

- Temperature abuse of fresh pork, in the chilled temperature area, induced critical *Salmonella* growth before spoilage occurred (**Result I**).
- At temperatures below 20°C, the natural background flora in pork slowed down growth of *Salmonella* Typhimurium DT104 and *Salmonella* Derby (**Result II**).



Experimental set up



Result I



Result II



Models

PRIMARY: Baranyi and Roberts model (DMFit web edition)

SECONDARY:

$$\sqrt{\mu_{\max}} = b \cdot (T - T_{\min})$$

where b is a constant, T is the temperature in °C and T_{\min} is the intercept between the model and the temperature axis.

Estimates

<i>Salmonella</i>	Pork	b	T_{\min}	R^2
<i>S. Derby</i>	Sterile	0.036	2.84	0.974
	Natural flora	0.043	6.86	0.978
<i>S. Typhimurium DT104</i>	Sterile	0.038	3.70	0.979
	Natural flora	0.045	7.34	0.990