



FISPRO - an ALGOL procedure for calculation of fission products

Højerup, C.F.

Publication date:
1976

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)


Citation (APA):
Højerup, C. F. (1976). *FISPRO - an ALGOL procedure for calculation of fission products*. Risø National Laboratory. Risø-M No. 1899

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

<p>Title and author(s)</p> <p>FISPRO - an ALGOL procedure for calculation of fission products</p> <p>by</p> <p>C.F. Højerup</p>	<p>Date October 1976</p> <p>Department or group</p> <p>Department of Reactor Technology</p> <p>Group's own registration number(s)</p> <p>RP-6-76</p>
<p>13 pages + 1 tables + 3 illustrations</p>	
<p>Abstract</p> <p>FISPRO is an ALGOL procedure performing calculations of the concentrations of a large number of fission products. All FP's with halflives ≥ 10 minutes are included. The procedure can easily be extended to include more FP's. Cross sections have been generated in 76 groups for 189 FP's, and can be condensed to arbitrary energy structures.</p>	<p>Copies to</p> <p>Library 100 C.F. Højerup 16</p>
<p>Available on request from: Risø Library, Research Establishment Risø, DK-4000 Roskilde, Denmark. Risø Bibliotek, Forsøgsanlæg Risø, 4000 Roskilde. Telephone: (03) 35 51 01, ext. 334, telex: 43116.</p>	<p>RISØ BIBLIOTEK</p> <p>5100023949699</p> 

INTRODUCTION

FISPRO is an ALGOL procedure, which performs point burn up calculations of the concentrations of a large number of fission products. The fission products may originate from fissioning of Th 232, Pa 233, U 233, U 234, U 235, U 236, U 238, Np 239, Pu 239, Pu 240, Pu 241 and Pu 242.

The cross sections of the products have been taken from UKNDL and a 76-groups library has been generated for 189 isotopes. Decay constants and yields have been taken from ref. 1.

As yields can only be found for the six "important" nuclides, i.e. Th 232, U 233, U 235, U 238, Pu 239 and Pu 241, these yields were ascribed to the remaining six in this way:

Yield for	Used for
Th 232	Th 232, Pa 233
U 233	U 233, U 234
U 235	U 235, U 236
U 238	U 238, Np 239
Pu 239	Pu 239, Pu 240
Pu 241	Pu 241, Pu 242

Two supporting procedures, FPDATA and FPCOND, are indispensable when using FISPRO. FPDATA establishes the reading of the necessary data (from the disk-file LIBDATA/CFHFPDATA) and FPCOND condenses the 76 group cross sections to any chosen group structure (which must be a sub-set of the 76 structure) by means of a 76-group neutron flux spectrum.

THEORY

The concentration $X(Z, M, I, t)$ of some fission product is governed by the production and losses, thus:

$$\frac{dX(Z, M, I, t)}{dt} = S(Z, M, I) - \lambda'(Z, M, I) \times X(Z, M, I, t)$$

Here the fission product concentration X is characterized by the atomic number Z , the mass number M , and a number I ($= 1$ or 0) distinguishing between two kinds of excited state, (excited or not excited). t is the time, and S is the production term, which may

contain contributions from:

direct yield from fission

neutron absorptions in X (Z, M-1, I = $\begin{Bmatrix} 0 \\ 1 \end{Bmatrix}$)

β -decays of X (Z-1, M, I = $\begin{Bmatrix} 0 \\ 1 \end{Bmatrix}$)

isomeric transition of X (Z, M, I = 1 \rightarrow 0)

(other forms of transformations, f.ex. β^+ and α decays could easily be incorporated, but have not, due to their very minor importance).

λ' is a removal constant, taking care of the removal of X (Z, M, I) by neutron absorption and by radioactive decay.

For a time interval in which S can be regarded as constant, the equation has the solution

$$X(Z, M, I, t) = S(Z, M, I) * (1 - \exp(-\lambda' t)) / \lambda' +$$

$$X(Z, M, I, t_0) * \exp(-\lambda' * (t - t_0))$$

Now, an inspection of the mechanisms contained in S readily reveals, that S in general has terms varying exponentially with time, with time constants as short as the shortest one included (in this version all isotopes with half-lives longer than 10 minutes are included).

Fortunately, however, short-lived isotopes saturate, so that their timevariation is only violent from $t = 0$ and for 2 - 3 half-lives thereafter.

If we, therefore, choose a series of timesteps, starting with a very short one, say $\frac{1}{2}$ * the shortest half-life, and the following ones increasing by a factor of 2 from step to step, we will get a reasonably good coverage of the growth of the isotope with the shortest half-life and better coverage of those with longer half-lives. This is illustrated by fig. 1. The doubling of the time steps also ensures that their number will be reasonable. (To cover 100 days total with 10 minutes as the first sub-step requires

$$n = \frac{\log(100 * 24 * 60/10)}{\log 2} \approx 14 \text{ sub time steps}$$




Figure 1

Growth of 3 isotopes with half-lives = 1, 2 and 10 times "shortest half-life". The true growth (the "1-exponentials") and the assumed growth when used as source terms, (the step functions) over the first 5 timesteps.

The full expression for the timevariation of the fission products in FISPRO is:

$$X(Z, M, I, t) = X(Z, M, I, t_0) * E + S * (1 - E) / \lambda'$$

where:

$$E = \text{EXP}(-\lambda' * (t - t_0))$$

$$\lambda' = \lambda [Z, M, I] + \text{ABS} [Z, M, I] ;$$

$$\lambda [Z, M, I] = \text{decay constant}$$

$$\text{ABS} [Z, M, I] = \sum_{K=1}^{\text{NG}} \text{FI}[K] * \text{SA} [Z, M, I, K] ;$$

FI [K] = Neutron flux in group K

SA [Z, M, I, K] = absorption cross section in group K

NG = number of groups

$$S = \text{GAM} [Z, M, I] +$$

(IF I = 1 = EXCITED THEN

$$\text{BRZM1M} [Z, M, I] * \lambda [Z-1, M, 1] * \bar{X} [Z-1, M, 1] +$$

$$\text{BRZM1G} [Z, M, I] * \lambda [Z-1, M, 0] * \bar{X} [Z-1, M, 0] +$$

$$\text{BRABS} [Z, M-1, 1] * \text{ABS} [Z, M-1, 1] * \bar{X} [Z, M-1, 1] +$$

$$\text{BRABS} [Z, M-1, 0] * \text{ABS} [Z, M-1, 0] * \bar{X} [Z, M-1, 0])$$

ELSE IF I = 0 = GROUNDSTATE THEN

$$\text{BRZM1M} [Z, M, I] * \lambda [Z-1, M, 1] * \bar{X} [Z-1, M, 1] +$$

$$\text{BRZM1G} [Z, M, I] * \lambda [Z-1, M, 0] * \bar{X} [Z-1, M, 0] +$$

$$\text{BRZM} [Z, M, I] * \lambda [Z, M, 1] * \bar{X} [Z, M, 1] +$$

$$(1 - \text{BRABS} [Z, M-1, 1]) * \text{ABS} [Z, M-1, 1] * \bar{X} [Z, M-1, 1] +$$

$$(1 - \text{BRABS} [Z, M-1, 0]) * \text{ABS} [Z, M-1, 0] * \bar{X} [Z, M-1, 0] ;$$

$$\text{GAM} [Z, M, I] = \sum_{\substack{J=\text{fissile} \\ \text{nuclides}}} \sum_{K=1}^{\text{NG}} \bar{X} [J] * \text{FI} [K] * \text{SF} [J, K] * \text{YIELD} [Z, M, I, J] ;$$

SF [J, K] = fission cross section of heavy nuclide J in group K

YIELD [Z, M, I, J] = fission yield of (Z, M, I) from fission of nuclide J

\bar{X} denotes the average concentration of the nuclide in question during the time step ($\bar{X} = \frac{1}{2} * (X(t) + X(t_0))$)

BRZM1M [Z, M, I] = the factor of the decays of nuclide (Z-1, M, I) that leads to formation of (Z, M, I)

BRZM1G [Z, M, I] = the fraction of the decays of nuclide (Z-1, M, 0) that leads to formation of (Z, M, I)

BRZM [Z, M, I] = the fraction of the decays of nuclide (Z, M, 1) that leads to formation of (Z, M, I)
This branching ratio obviously only exists for I = 0.

BRABS [Z, M, I] = the fraction of absorptions in nuclide (Z, M, I) that leads to formation of (Z, M+1, 1)
Note the different definitions of the two types of branching ratios. The branching ratios associated with radioactive decays are characterized by the products they decay into, while the branching ratio for absorption is characterized by the absorbing nucleus itself, and says, which fraction of the absorptions that leads to the formation of a nucleus in an excited state.

CALL OF THE PROCEDURE

The procedure head is

FISPRO (FP, LAM, YIELD, BRZM1M, BRZM1G, BRZM, BRABS, SA, EXIST, ZL, ZU, NG, FPSA, NDF, NDFP, FI, SF, DT, TMIN, PRINTER, FIL6);

```
VALUE DT, ZL, ZU, NG, TMIN, PRINTER;  
REAL DT, TMIN;  
INTEGER ZL, ZU, NG, PRINTER;  
ARRAY NDF, NDFP, FI, FPSA [0], FP, LAM, SF [0,0],  
SA, YIELD [0,0,0].  
ARRAY BRZM1M, BRZMIG, BRZM, BRABS [0,0] ;  
INTEGER ARRAY EXIST [0,0] ;  
FILE FIL6;
```

DT is the time step in days
TMIN is the length of the shortest (starting) sub timestep
in days.
ZL is the lowest atomic number of fission products
treated (= 30)
ZU is the highest at. number of F.P. treated (=67)
NG is the number of energy groups
PRINTER governs the amount of printed information:
if PRINTER \geq 9 the quantities ABS and GAM defined in
the preceding chapter will be printed; if PRINTER
 \geq 8 also the fission product concentrations in the
sub timesteps will be printed;
if PRINTER \geq 7 also the macroscopic absorption cross
section (Σ PFSA) of all the fission products combined
will be printed.
This printing takes place on the file
FIL6 Printer file.
NDF, NDFP [0:12] contain the concentrations at the end (NDF)
and at the beginning (NDFP) of the timestep DT, of
the heavy nuclides Th 232 (=1), Pa 233 Pu 242
(= 12).
FI [0:NG] = The flux spectrum in NG groups (FI [0] is not
used).
FPSA [0:NG] = The poisoning cross section of the fission
products, i.e. usually the real result of the use of
the procedure.
FP [0:ZU, 0:60] contains the fission product concentrations.
Only the rows from ZL to ZU are used. In each row (Z)
the places (N) are allocated according to the algorithm:
$$N: = 2 \times M - 4 \times Z - 12 + 1$$

where M is the mass number and I takes the value 0 for an excited state and 1 for the groundstate.

LAM [0:ZU, 0:60] contains the decay constants.

SF [0:12, 0:NG] contains the fission cross sections of the heavy nuclides.

SA [0:ZU, 0:60, 0:NG] contains the absorption cross sections of the fission products.

YIELD [0:ZU, 0:60, 0:12] contains the yields of the fission products from the 12 heavy nuclides.

BRZM1M,

BRZM1G,

BRZM,

BRABS [0:ZU, 0:60] contain the various branching ratios described in the preceding chapter.

EXIST [0:ZU, 0:61] contains:

in position [Z,0] : - NL, where NL is the lowest value, for which there exists a fission product with position [Z,NL] in FP.

in position [Z,1] : - NU, where NU is the highest etc. etc.

in position [Z,N] = K, where K is a non-negative integer meaning that:

K = 0: No fission product exists in this position (Z,N).

K = 1: The fission product (Z,N) exists, and has an absorption cross section $\neq 0$.

K = 2: The fission product (Z,N) exists, but has a cross section = 0.

The arrays

EXIST, LAM, YIELD, SA, BRZM1M, BRZM1G, BRZM, BRABS

will be supplied with data by calling the procedure

FPDATA (EXIST, FPNAME, LAM, BRZM1G, BRZM1M, BRZM, BRABS, YIELD, SA, NG, ZL, ZU, PRINTER, FIL5, FIL6);

where, in addition to quantities already mentioned, ALPHA ARRAY FPNAME [0:ZU, 0:60] will contain the names of the fission products f.ex. "XE135G",

ZL, ZU and NG must have the values 30, 67 and 76, respectively, and for PRINTER ≥ 9 the data will be printed out on FIL6. The data are read from FIL5, which in the procedure call must have the name LIBDATA/CFHFPPDATA.

If another group structure than the 76 group RISØ standard structure is to be used, the cross sections for the F.P.'s can be condensed by calling the procedure

```
CONDFP(FPSAFG, FPSACG, FI, EXIST, GMAX, GLOW, ZL, ZU, NG);
```

where

ARRAY FPSAFG [0:ZU, 0:60, 0:76] are the Fine Group cross sections (obtained by Fpdata)

ARRAY FPSACG [0:ZU, 0:60, 0:NG] are the Coarse Group cross sections.

ARRAY FI [0:76] must contain a 76-group flux spectrum and INTEGER ARRAY GMAX, GLOW [0:NG] must contain the connection between the 76-group structure and the NG-group structure. The rules are such that:

Coarse group no. I will consist of Fine Groups nos. GMAX [I] to and including GLOW [I].

It follows that usually

$$GMAX [I+1] = GLOW [I] + 1.$$

The procedures FISPRO, Fpdata and CONDFP are made available in an ALGOL programme by the command $\$$ INCLUDE "RRPMS/FISPRO". In figure 2. are shown the fission products that are presently contained in the FISPRO data library.

THE ALGOL TEXT

A listing of the three procedures CONDFP, FISPRO and FPDATA is shown below.

```
PROCEDURE CONDFP(FPSAFG,FPSACG,FI, EXIST,GMAX,GLOW,ZL,ZU,NG);
VALUE ZL,ZU,NG;
INTEGER ZL,ZU,NG;
ARRAY FPSAFG,FPSACG[0,0,0],FI[0];
INTEGER ARRAY EXIST[0,0],GMAX,GLOW[0];
BEGIN
REAL S; INTEGER I,AI,BI,G,Z,N,NL,NU;

ARRAY F[0:NG];
FOR I:=1 STEP 1 UNTIL NG DO BEGIN AI:=GMAX[I]; BI:=GLOW[I];
S:=0;
FOR G:=AI STEP 1 UNTIL BI DO S:=S+FI[G];
F[I]:=S;
END;
FOR Z:=ZL STEP 1 UNTIL ZU DO BEGIN
NL:=-EXIST[Z,0]; NU:=-EXIST[Z,1];
FOR NI=NL STEP 1 UNTIL NU DO IF EXIST[Z,NI]=1 THEN
FOR I:=1 STEP 1 UNTIL NG DO BEGIN AI:=GMAX[I]; BI:=GLOW[I];
S:=0;
FOR G:=AI STEP 1 UNTIL BI DO S:=S+FI[G]*FPSAFG[Z,N,G];
FPSACG[Z,N,I]:=S/FI[I];
END ELSE
FOR I:=1 STEP 1 UNTIL NG DO FPSACG[Z,N,I]:=0;
END;
END;

PROCEDURE FISPRO(FP,LAM,YIELD,BRZM1M,BRZM1G,BRZM,BRABS,SA,EXIST,ZL,ZU,
NG,FPSA,NDP,NDFP,FI,SF,DT,TMIN,PRINTER,FILE);
VALUE DT,ZL,ZU,NG,TMIN,PRINTER; REAL DT,TMIN; INTEGER ZL,ZU,NG,PRINTER;
ARRAY NDP,NDFP,FI,FPSA[0],FP,LAM,SF[0,0],SA,YIELD[0,0,0];
ARRAY BRZM1M,BRZM1G,BRZM,BRABS[0,0];
INTEGER ARRAY EXIST[0,0];
FILE FILE;
BEGIN
ARRAY ABS,GAK,FPC[0:ZU,0:60];

INTEGER Z,NL,NU,N,ZM1,NP4,NP5,NP3,NM1,NM2,NM3,K,I,J;
REAL S,A,B,C,LAMP,X,E,TSEC,TS,T;
FOR Z:=ZL STEP 1 UNTIL ZU DO BEGIN
NL:=-EXIST[Z,0]; NU:=-EXIST[Z,1];
FOR NI=NL STEP 1 UNTIL NU DO
IF EXIST[Z,NI]>0 THEN BEGIN
S:=0;
IF EXIST[Z,NI]=1 THEN
FOR K:=1 STEP 1 UNTIL NG DO S:=S+FI[K]*SA[Z,N,K]*B-24;
ABS[Z,NI]:=S;
S:=0;
FOR I:=1 STEP 1 UNTIL 12 DO BEGIN F:=(NOF[I]+NOFP[I])/2*YIELD[Z,N,I];
FOR K:=1 STEP 1 UNTIL NG DO S:=S+FI[K]*E*SFI[I,K];
END;
GAK[Z,NI]:=S;
END;
END;
END;
```

```
IF PRINTER GEO 9 THEN BEGIN
WRITE(FILE, <"ABS">);
FOR Z:=ZL STEP 1 UNTIL ZU DO
WRITE(FILE, <10F12.3>,
ABS(Z,*1));
WRITE(FILE, <"GAM">);
FOR Z:=ZL STEP 1 UNTIL ZU DO
WRITE(FILE, <10F12.3>,
GAM(Z,*1));
END;
IF PRINTER GEO 8 THEN
WRITE(FILE, <"FP ">);
TS:=TMIN/2; T:=0;
FOR TS:=2*TS WHILE T < DT DO BEGIN T:=T+TS;
IF T>DT*.9 THEN BEGIN TS:=TS-T+DT; T:=DT; END;
TSEC:=TS*8400;
FOR Z:=ZL STEP 1 UNTIL ZU DO BEGIN
NL:=-EXISTIZ,01; NU:=-EXISTIZ,11;
FOR N:=NL STEP 1 UNTIL NU DO BEGIN
IF EXISTIZ,N1>0 THEN BEGIN
%METASTABIL KERNE
FP0(Z,N1):=B:=FP(Z,N1);
ZM1:=Z-1; NP4:=N+4; NP5:=N+5;
NM1:=N-1; NM2:=NM2;
S:=GAM(Z,N1)+
(IF EXISTIZM1,NP4)>0 THEN BRZM1(Z,N1)*LAM(ZM1,NP4)*
(FP(ZM1,NP4)+FP0(ZM1,NP4))/2 ELSE 0)+
(IF EXISTIZM1,NP5)>0 THEN BRZM1(Z,N1)*LAM(ZM1,NP5)*
(FP(ZM1,NP5)+FP0(ZM1,NP5))/2 ELSE 0)+
(IF EXISTIZ ,NM1)=1 THEN ABS(Z,NM1)*BRABS(Z,NM1)*
(FP(Z ,NM1)+FP0(Z ,NM1))/2 ELSE 0)+
(IF EXISTIZ ,NM2)=1 THEN ABS(Z,NM2)*BRABS(Z,NM2)*
(FP(Z ,NM2)+FP0(Z ,NM2))/2 ELSE 0);
LAMP:=LAM(Z,N1)+ABS(Z,N1); X:=LAMP*TSEC;
E:=IF X<.6 THEN (2-X)/(2+X) ELSE EXP(-X);
IF LAMP=0 THEN FP(Z,N1):=B+S*TSEC ELSE
FP(Z,N1):=B*E+S*(1-E)/LAMP;
END;
N:=N+1;
%KERNE I GRUNDFILSTAND
IF EXISTIZ,N1>0 THEN BEGIN
FP0(Z,N1):=B:=FP(Z,N1); ZM1:=Z-1; NP3:=N+3; NP4:=N+4; NM1:=N-1;
NM2:=N-2; NP3:=N-3;
S:=GAM(Z,N1)+
(IF EXISTIZM1,NP3)>0 THEN BRZM1(Z,N1)*LAM(ZM1,NP3)*
(FP(ZM1,NP3)+FP0(ZM1,NP3))/2 ELSE 0)+
(IF EXISTIZM1,NP4)>0 THEN BRZM1(Z,N1)*LAM(ZM1,NP4)*
(FP(ZM1,NP4)+FP0(ZM1,NP4))/2 ELSE 0)+
(IF EXISTIZ ,NM1)>0 THEN BRZM (Z,N1)*LAM(Z ,NM1)*
(FP(Z ,NM1)+FP0(Z ,NM1))/2 ELSE 0)+
(IF EXISTIZ ,NM2)=1 THEN ABS(Z,NM2)*(1-BRABS(Z,NM2))*
(FP(Z ,NM2)+FP0(Z ,NM2))/2 ELSE 0)+
(IF EXISTIZ ,NM3)=1 THEN ABS(Z,NM3)*(1-BRABS(Z,NM3))*
(FP(Z ,NM3)+FP0(Z ,NM3))/2 ELSE 0);
LAMP:=LAM(Z,N1)+ABS(Z,N1); X:=LAMP*TSEC;
E:=IF X<.6 THEN (2-X)/(2+X) ELSE EXP(-X);
IF LAMP=0 THEN FP(Z,N1):=B+S*TSEC ELSE
FP(Z,N1):=B*E+S*(1-E)/LAMP;
END;
END;
END;
```

```
IF PRINTER GEO 6 THEN BEGIN
WRITE(FILE, <"/"TIMESTEP=" , F15.5, "          TIME=" , F15.5, >, TS, T);
```

```
FOR Z:=ZL STEP 1 UNTIL ZU DO
BEGIN
NL := -EXIST[Z, 0]; NU := -EXIST[Z, 1];
WRITE(FILE, <"Z, NL, NU: " , 3I5, >, Z, NL, NU);
WRITE(FILE, <10F12.3, >, FOR K:=NL STEP 1 UNTIL NU DO FP[Z, K]);
END;
END;
END;
```

```
FOR K:=1 STEP 1 UNTIL NG DO FPSA[K]:=0;
FOR Z:=ZL STEP 1 UNTIL ZU DO BEGIN
NL := -EXIST[Z, 0]; NU := -EXIST[Z, 1];
FOR N:=NL STEP 1 UNTIL NU DO
IF EXIST[Z, N]=1 THEN BEGIN
A:=FP[Z, N]*G*24;
FOR K:=1 STEP 1 UNTIL NG DO FPSA[K]:=FPSA[K]+A*SA[Z, N, K];
END;
END;
END;
```

```
IF PRINTER GEO 7 THEN BEGIN
WRITE(FILE, <"ABSORPTION IN FISSION PRODUCTS: "/>);
WRITE(FILE, <10F12.5, >);
FOR K:=1 STEP 1 UNTIL NG DO FPSA[K]);
END;
END;
```

```
PROCEDURE Fpdata (EXIST, FPNAME, LAM, BRZM1G, BRZM1M, BRZY, LRABS, YIELD, SA, NG,
ZL, ZU, PRINTER, FIL5, FIL6);
VALUE NG, PRINTER, ZL, ZU;
INTEGER NG, PRINTER, ZL, ZU;
ARRAY LAM, BRZM1G, BRZM1M, BRZY, LRABS[0, 0], YIELD, SA[0, 0, 0];
INTEGER ARRAY EXIST[0, 0];
ALPHA ARRAY FPNAME[0, 0];
FILE FIL5, FIL6;
BEGIN
INTEGER Z, N, J, I, K;
```

```
ALPHA NAME;
LABEL L, L1;
FOR I:=ZL STEP 1 UNTIL ZU DO BEGIN
FOR K:=0 STEP 1 UNTIL 65 DO EXIST[I, K]:=0;
EXIST[I, 0]:=60;
END;
```

```
IF PRINTER GEO 9 THEN
WRITE(FILE, <"          LAMBDA   BR Z-1 G   BR Z-1 M   BR Z-0 M YIELD%: TH
U233      U235      U238          PU239          PU241">);
```

```
L:
READ(FIL5, /,
Z, N, J);
IF PRINTER GEO 9 THEN
WRITE(FILE, <"/"Z, N, J: " , 2(I5, " , " , 15, >,
Z, N, J);
```

```
IF Z>0 THEN BEGIN
  READ(FILE,/,
  NAME);
  IF PRINTER GEG 9 THEN
  WRITE(FILE,<A6>,
  NAME);
  N:=2*N-4*Z-12+J;
  FPNAME[Z,N]:=NAME;
  IF N<-EXIST[Z,0] THEN EXIST[Z,0]:=-N+J;
  IF N>-EXIST[Z,1] THEN EXIST[Z,1]:=-N;
  READ(FILE,/,
  LAM[Z,N],BRZM1G[Z,N],BRZM1M[Z,N],BRZM[Z,N],
  FOR K:=5,7,9,11,3,1 DO YIELD[Z,N,K]);
  IF PRINTER GEG 9 THEN
  WRITE(FILE,<I1F11.3>,
  LAM[Z,N],BRZM1G[Z,N],BRZM1M[Z,N],BRZM[Z,N],
  FOR K:=5,7,9,11,3,1 DO YIELD[Z,N,K]);
  FOR K:=1 STEP 2 UNTIL 11 DO YIELD[Z,N,K]:=YIELD[Z,N,K+1]:=YIELD[Z,N,K]*
  .01;
  EXIST[Z,N]:=2;
  GO TO L1;
END;
L1:
READ(FILE,/,Z,N,J);
IF PRINTER GEG 9 THEN
WRITE(FILE,</"Z,N,J:",3I5>,Z,N,J);
IF Z>0 THEN BEGIN
  N:=2*N-4*Z-12+J;
  READ(FILE,<6E12.5>,
  FOR K:=1 STEP 1 UNTIL NG DO SA[Z,N,K]);
  READ(FILE,Z,BRABS[Z,N]);
  IF EXIST[Z,N]>0 THEN BEGIN
  EXIST[Z,N]:=-1;
  IF PRINTER GEG 9 THEN
  WRITE(FILE,<A6>,FPNAME[Z,N]);
  END ELSE
  IF PRINTER GEG 9 THEN
  WRITE(FILE,<"CROSS SECTION, BUT NO FP">);
  IF PRINTER GEG 9 THEN
  WRITE(FILE,<6E15.5>,
  FOR K:=1 STEP 1 UNTIL NG DO SA[Z,N,K]);
  IF PRINTER GEG 9 THEN
  WRITE(FILE,<"BRABS=",F12.5>,BRABS[Z,N]);
  GO TO L1;
END;
END;
```

COMPARISON WITH FIPO

The fission product poisoning has up to now been calculated with a procedure FIPO (2) in Risø reactor physics codes. FIPO has its own 10 groups data library, and thus locks all calculations to the same 10 groups.

In order to compare FISPRO and FIPO, they were both used with the same group structure on the same example, a BWR calculated with the programme CCC (3).

The burn up curve for k_{eff} is shown in fig. 3. The poisoning turns out to be significantly less, when calculated by FISPRO, amounting to about 1% on k_{eff} per 10 MWd/kg U.

A closer analysis of the reasons for this revealed, that the cross sections of a few, but important, FP's happened to be somewhat larger in FIPO than in FISPRO. This may either originate in differences in the basic data sets (which are different editions of UKNDL fission product library) or from the fact that FIPO's cross sections have been generated once and for all by means of some characteristic LWR flux spectrum, while FISPRO's cross sections are generated by CONDFP with the actual spectrum.

A more detailed comparison of the two procedures is made in table 1, where the activities of the most important radioactive isotopes are listed for a PWR core at 18 MWd/kg U and 40 kW/kg U. In the same table are also given values deduced from WASH-1400 (4).

Table 1

Comparison of radioactive inventories (Mci) in a reactor core burned to 18 MWd/kg U, as calculated in WASH 1400, with FIPO, and with FISPPO

ISOTOPE	WASH 1400	FIPO	FISPPO
KR 85	0.56	0.49	0.50
KR 85M	24	-	26
KR 87	47	-	49
KR 88	68	-	68
RB 86	0.026	-	0.053
SR 89	94	93	95
SR 90	3.7	3.7	3.7
SR 91	110	-	120
Y 90	3.9	3.8	3.7
Y 91	120	120	120
ZR 95	150	150	150
ZR 97	150	-	150
NB 95	150	150	150
MO 99	160	160	160
TC 99M	140	-	140
RU 103	110	120	120
RU 105	72	-	73
RU 106	25	22	22
RH 105	49	62	69
TE 127	5.9	-	4.1
TE 127M	1.1	0.60	0.62
TE 129	31	-	23
TE 129M	5.3	4.2	4.3
TE 131M	13	12	11
TE 132	120	110	120
SB 127	6.1	3.2	4.2
SB 129	33	-	24
I 131	85	87	81
I 132	120	-	120
I 133	170	170	140

Table 1 (continued)

ISOTOPE	WASH 1400	FIPO	FISPRO
I 134	190	-	190
I 135	150	160	160
XE 133	170	170	140
XE 135	34	43	45
CS 134	7.5	3.0	2.9
CS 136	3.0	1.1	2.3
CS 137	4.7	4.5	4.6
BA 140	160	150	150
LA 140	160	160	160
CE 141	150	140	140
CE 143	130	140	140
CE 144	85	86	86
PR 143	130	140	140
ND 147	60	58	57
NP 239	1600	1500	1500
PU 241	3.4	4.0	4.0

REFERENCES

1. Compilation of Fission Product Yields. M.E. Meck and B.F. Rider. NEDO-12154-1, 74 NED6, 1974.
2. The Fission Product Treatment in the CEB Unit Cell Burn-up Programme. Leif Mortensen. Risø-M-1356. 1976.
3. An Example of Burn Up Calculations in Three Dimensions for a CANDU Core. C.F. Højerup and Torben Petersen. RISØ-M-1773. 1974.
4. Reactor Safety Study. App. VI. US NRC. (Normann Rasmussen). WASH 1400. 1975.

FIG. 3.

k_{eff} for a BWR
calculated with
EIPO and FISPRO

