



## Current and emerging techniques for contaminant mapping and data visualization at DNAPL sites

Wealthall, Gary; Durant, Neal; Grosen, Bernt; Jørgensen, Torben H.; Broholm, Mette Martina; Janniche, Gry Sander; Christensen, Anders G.; Kerrn-Jespersen, Henriette

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Wealthall, G., Durant, N., Grosen, B., Jørgensen, T. H., Broholm, M. M., Janniche, G. S., Christensen, A. G., & Kerrn-Jespersen, H. (2013). *Current and emerging techniques for contaminant mapping and data visualization at DNAPL sites*. Abstract from AquaConSoil 2013 — 12th International UFZ-Deltares Conference on Groundwater-Soil-Systems and Water Resource Management, Barcelona, Spain.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **Current and emerging techniques for DNAPL site characterization in sediments and fractured bedrock**

Gary Wealthall<sup>1</sup>, Neal Durant (Geosyntec Consultants, Inc.), Bernt Grosen , Torben Højbjerg Jørgensen, (COWI A/S), Mette Martine Broholm, Gry Sander Janniche (DTU Environment ), Anders G. Christensen (NIRAS) Henriette Kerrn-Jespersen (Capital Region of Denmark)

<sup>1</sup>130 Research Lane, Guelph, Ontario, Canada, N1G 5G3 Phone: +1.519.822.2230 ext. 333  
Mobile: +1.519.400.8033, gwealthall@geosyntec.com

Recent advances in the development of contaminant specific site investigation tools has significantly enhanced our ability to characterize the spatial architecture of DNAPL source zones in both sedimentary and fractured bedrock environments. The application of innovative site characterization methods will be discussed in the context of contaminant delineation, remediation design, technology verification and regulatory acceptance.

We present a range of site investigation tools, based on the principle of combined lines of evidence and the premise that a single technique is not available to fully delineate DNAPL distribution in the subsurface. Key to this strategy is the selection of technologies with multiple scales of measurement and data quality, of which there are two main categories. The first category provides qualitative, dense spatial data, often with higher detection limits over a preset value. These methods are generally of lower cost, produce vertical profiles of real-time data and are primarily used to identify site areas that require further investigation. Examples of such "decision quality" methods are drive point tools, such as laser-induced fluorescence profiling and membrane interface probing in unconsolidated sediments, and the NAPL FLUTE™ system in fractured bedrock. These first category methods produce qualitative data that are used to guide the sampling strategy for the application of the second category of technologies that are typically compound-specific and generate quantitative, precise data that have low detection limits. Examples include the use of fully cored boreholes (and associated subsampling) and high resolution multilevel well completions. Where these two categories of measurement technologies are used in tandem, a more complete and accurate dataset is achieved and additional site mobilizations are limited. These second category methods tend to be higher cost with longer turnaround times that preclude on-site decision making, hence applying them to quantify, rather than produce a conceptual model, facilitates a key cost saving.

The use of this 'toolbox' approach is described for a comprehensive fractured bedrock case study in Denmark (funded by the Capital Region of Denmark), where a chlorinated solvent DNAPL has impacted underlying glacial till fractured limestone bedrock. The findings of the study and recommendations for future investigations will be discussed in the context of published literature and current and emerging best practice guidance.