



## Ion acoustic waves in the presence of high frequency oscillators

Pécsele, Hans

*Publication date:*  
1975

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Pécsele, H. (1975). *Ion acoustic waves in the presence of high frequency oscillators*. Risø-M No. 1790

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Risø - M - 1790

<p><b>Title and author(s)</b></p> <p>Ion acoustic waves in the presence of high frequency oscillations</p> <p>by</p> <p>H.L. Pécseli</p>	<p><b>Date</b> February 1975</p> <p><b>Department or group</b></p> <p>Physics</p> <p><b>Group's own registration number(s)</b></p>
<p>19 pages + tables + illustrations</p>	
<p><b>Abstract</b></p> <p>Spatially varying high frequency oscillations will influence the electron motion in plasmas (through ponderomotive forces) and in turn affect the characteristics of ion acoustic oscillations. In this report we derive the dielectric function for long wavelength, low frequency ion acoustic waves in the presence of high frequency, short wavelength oscillations. Apart from reproducing well known results this dielectric function shows the possibility of unstable ion acoustic waves for extreme intensities of high frequency oscillations. It is demonstrated that moderate intensities have a stabilizing effect on linearly unstable double humped ion distribution functions and it is argued that this effect may be of importance in laboratory experiments. Finally we point out some generalisations.</p>	<p><b>Copies to</b></p>
<p>Available on request from the Library of the Danish Atomic Energy Commission (Atomenergi-kommissionens Bibliotek), Risø, DK-4000 Roskilde, Denmark Telephone: (08) 35 51 01, ext. 334, telex: 43116</p>	

CONTENTS

	Page
Introduction.....	1
The dielectric function.....	5
Discussion .....	11
Generalization and conclusion.....	13
References.....	17

Ion acoustic waves in the presence  
of high frequency oscillations

H.L. Pécseli

Introduction

In this preliminary report we consider the influence of high frequency short wavelength electron oscillations on the propagation on low frequency long wavelength ion acoustic oscillations.<sup>1,2</sup> The high frequency oscillations are denoted by  $(\omega, k)$  and the ion acoustic oscillations by  $(\Omega, q)$ . We describe the ions by the collisionfree Vlasov equation

$$\frac{\partial f_i}{\partial t} + v \frac{\partial f_i}{\partial x} + \frac{e}{M} E \frac{\partial f_i}{\partial v} = 0 \quad (1)$$

The electrons are considered as an isothermal fluid and we let the electron motion be determined by<sup>1</sup>

$$m \left( \frac{\partial v_e}{\partial t} + v_e \frac{\partial v_e}{\partial x} \right) = -eE - \frac{\kappa T_e}{n_e} \frac{\partial n_e}{\partial x} + F \quad (2)$$

The force  $F$  arises from the high frequency oscillations (plasmons) and is determined by the following argument<sup>1,3</sup>: Let  $E(x, t)$  be a high frequency field that varies slowly in amplitude with position. The equation of motion for an electron is then

$$\ddot{x} = \frac{e}{m} E(x, t) \quad (3)$$

We look for solutions in the form  $x = y + \xi$  where  $y$  is a slow displacement of the particle due to spatially varying amplitude of  $E$  and  $\xi$  is a small displacement in the high frequency field. Expanding  $E(x, t)$  in powers of  $\xi$  we obtain

$$\dot{y} + \ddot{\xi} = \frac{e}{m} \left( E(y, t) + \xi \frac{\partial}{\partial y} E(y, t) \right) \quad (4)$$

Taking the rapidly oscillating term in (4) we obtain

$$\xi = -\frac{e}{m} \sum_k \frac{E_k(y)}{\omega^2} \exp(-i\omega t) \quad (5)$$

Inserting (5) in (4) and averaging over at time  $\tau$  where  $\frac{1}{\omega} \ll \tau \ll \frac{1}{\Omega}$  we obtain

$$F \equiv m \langle \dot{y} \rangle = -\frac{2}{\Omega} \sum_k \frac{e^2 E_k^2}{2m\omega^2} \quad (6)$$

A similar problem is considered in ref. 4 in a more general form. The high frequency oscillations obeys the dispersion relation

$$\omega = \omega_p \left(1 + \frac{1}{2} \frac{k^2 \kappa T_e}{m \omega_p^2}\right) = \omega_p \left(1 + \frac{1}{2} (k\lambda_d)^2\right) \quad (7)$$

where  $\omega_p = \omega_p(y)$  is the local electron plasma-frequency. We prefer to use  $k$  as a summation index. Transforming the sum in (6) into an integral (with proper change in the dimensions of the integrand since  $\sum_k \rightarrow \frac{1}{2\pi} \int dk$ ) we obtain

$$F = -\frac{e^2}{2m} \frac{\partial}{\partial y} \int \frac{E_k^2}{\omega_k^2} dk \quad (8)$$

Introducing  $N_k = \frac{E_k^2}{\omega_k^2}$  equation (2) reduces to

$$m \left( \frac{\partial N_k}{\partial t} + v_k \frac{\partial N_k}{\partial x} \right) = -eE - \frac{\kappa T_e}{n_0} \frac{\partial n_0}{\partial x} - \frac{e^2}{2\epsilon_0 m} \frac{\partial}{\partial x} \int \frac{N_k}{\omega_k} dk \quad (9)$$

The variation of  $N_k$  is determined by

$$\frac{\partial N_k}{\partial t} + \frac{2\omega}{\partial k} \frac{\partial N_k}{\partial x} - \frac{2\omega}{\partial x} \frac{\partial N_k}{\partial k} = -2\gamma_k N_k \quad (10)$$

where  $\gamma_k$  is the appropriate decay time for the distribution of high frequency oscillations. Equation (10) corresponds to the zero order W.K.B. approximation for the evolution of a wave packet of high frequency oscillations. This approximation is valid since we assume  $k \gg q$  and  $\frac{\partial}{\partial x}$  refers to the spatial variation of the ion density.

Using (7) we reduce (10) to:

$$\frac{\partial N_k}{\partial t} + 3 \frac{\kappa T_e k}{m \omega_p} \frac{\partial N_k}{\partial x} - \left(1 - \frac{1}{2} (k\lambda_d)^2\right) \frac{\omega_p}{2n_0} \frac{\partial n_0}{\partial x} \frac{\partial N_k}{\partial k} = -2\gamma_k N_k \quad (11)$$

Finally we introduce Poisson's equations

$$\frac{\partial E}{\partial x} = \frac{e}{\epsilon_0} (n_i - n_e) \quad (12)$$

Equations (1), (9), (11) and (12) form a closed set of equations.

In the following we will use the terms "radiation pressure" for  $F$  and "number of quasiparticles" (or plasmons) for  $N_k$  although this terminology may be slightly misleading.

We linearize the equations (1), (9) and (11) and obtain

$$\frac{\partial f_i}{\partial t} + v \frac{\partial f_i}{\partial x} + \frac{e}{m} E f'_i(v) = 0 \quad (13)$$

$$m \frac{\partial N_k}{\partial t} = -eE - \frac{\kappa T_e}{n_0} \frac{\partial n_0}{\partial x} - \frac{N_k}{2n_0} \frac{\partial}{\partial x} \int N_k dk \quad (14)$$

$$\frac{\partial N_k}{\partial t} + \frac{3\kappa T_e k}{m \omega_p} \frac{\partial N_k}{\partial x} - \left(1 - \frac{1}{2} (k\lambda_d)^2\right) \frac{\omega_p}{2n_0} \frac{\partial n_0}{\partial x} \frac{\partial N_k}{\partial k} = 0 \quad (15)$$

In the integrand in the last term of (9) we have replaced  $1/\omega_k$  with  $1/\omega_p$ . This is a reasonable approximation for the case of weak dispersion (see eq. (7)). The zero order distribution of quasiparticles,  $N_0$ , is a function of time  $N_0(t) = N_0(k) \exp(-2\gamma_k t)$ . In the following we assume  $\gamma_k = 0$  corresponding to a stationary spectrum. We may for inst. assume that  $N_0(k)$  qualitatively represents a fluctuation level due to the thermal fluctuations of the electrons.

In a more interesting case  $N_0(k)$  is determined by a turbulent spectrum.

The following features of  $F$  and  $N_k$  may be noted: the force  $F$  tries to push the electrons to regions at low fluctuation level i.e. small  $N_k$  thus justifying the term "radiation pressure". The

force acting on the quasi-particles,  $-\frac{\partial \omega}{\partial x}$  tries to condense the quasi-particles in regions of low ion density, a feature obtained also by physical intuition when the dispersion relation (7) is taken into account.

In eqs. (13)-(15)  $\omega_p$  and  $d$  refer to the unperturbed state.

### The dielectric function

By considering the adiabatic response to a test charge we find the dielectric function for the medium. In the calculations we neglect electron inertia i.e. the term on the left side of (14).

$$\epsilon(q, \Omega) = \frac{1}{(qd)^2} \left[ (1 + (qd)^2) - \left( C_0^2 + \frac{\omega_p^2}{2\pi} \int \frac{Q(k)N_0'(k)}{v_0(k) - \frac{\Omega}{q}} dk \right) \int \frac{f_0(v)}{v - \frac{\Omega}{q}} dv \right] \quad (15)$$

where

$$C_0^2 = \frac{\kappa T_e}{m}$$

$$Q(k) = \left( 1 - \frac{1}{2} (kd)^2 \right) \frac{\omega_p^2}{2\pi}$$

and

$$v_0(k) = \frac{\partial \omega}{\partial k} = 3 \frac{\kappa T_e}{m \omega_p} k$$

The sign  $\int$  indicates that the integration path runs below the pole. Since we have assumed that  $q$  is small ( $q \ll 1/d$ ) we will neglect the term  $(qd)^2$  in the parenthesis in (15).

The validity of (16) for the case where  $N_0=0$  is well established experimentally.<sup>6,7</sup> We will now consider the effect of the high frequency oscillations.

We consider three special cases since eq. (16) is somewhat confusing for arbitrary  $f_0(v)$  and  $N_0(k)$ .

I. Assume  $f_0(v) = \delta(v)$ .

Then

$$\text{Re } \epsilon(q, \Omega) = 1 - \left( \frac{\Omega}{2\pi} \right)^2 \left( C_0^2 + \frac{\omega_p^2}{2\pi} \int \frac{Q(k)N_0'(k)}{v_0(k) - \frac{\Omega}{q}} dk \right) \quad (17a)$$

$$\text{Im } \epsilon(q, \Omega) = - \frac{\omega_p^2}{6\pi \kappa T_e} \pi Q \left( \frac{\Omega \omega_p m}{3q \kappa T_e} \right) N_0' \left( \frac{\Omega \omega_p m}{3q \kappa T_e} \right) \left( \frac{\Omega}{2\pi} \right)^2 \quad (17b)$$

For  $N_0=0$  eq. (17) describes undamped oscillations. When  $N_0 \neq 0$ , these oscillations are damped (if  $N_0' < 0$  since  $Q > 0$  for cases of interest) on quasiparticles with distribution  $N_0(k)$  in a way much similar to the conventional Landau damping (This effect is discussed in Ref. 2). Conversely a distribution of quasiparticles for which  $N_0' > 0$  for some  $k$  may lead to instability. (Note however that for  $\Omega/q \approx C_s$  the quantity  $(\Omega \omega_p m)/(3q \kappa T_e)$  corresponds to a very small  $k$ -value. We have assumed  $q \ll k$ . For small  $k$  we may let  $Q \sim \frac{\omega_p}{2 n_0}$ ). The imaginary part of the frequency is determined by the well known expression

$$Im \omega \approx + \frac{Im \mathcal{E}(K, \Omega_K)}{\frac{\partial Re \mathcal{E}(K, \Omega_K)}{\partial \Omega_K}} \quad (18)$$

where  $\Omega_K$  is the solution to  $Re \mathcal{E}(K, \Omega_K) = 0$ . Eq. (18) is valid for small  $Im \omega$ .

II. As a second case assume  $N_0(k) = N_0 K \delta(k-K)$ . This assumption corresponds to the case of a monochromatic high frequency oscillation. Then:

$$Re \mathcal{E} = 1 - \left( C_s^2 + \frac{\omega_p^2}{24 M \kappa T_e} \left( \frac{3 K^2 \kappa T_e + 2 \omega_p^2 m - 2 K m \omega_p \frac{\Omega}{q}}{n_0 \left( K - \frac{\Omega \omega_p m}{3 q \kappa T_e} \right)^2} \right) \right) P \int \frac{f_0'}{v - \frac{\Omega}{q}} dv \quad (19a)$$

$$Im \mathcal{E} = -P \int \frac{f_0'(\frac{\Omega}{q})}{\left( C_s^2 + \frac{\omega_p^2}{24 M \kappa T_e} \left( \frac{3 K^2 \kappa T_e + 2 \omega_p^2 m - 2 K m \omega_p \frac{\Omega}{q}}{\left( K - \frac{\Omega \omega_p m}{3 q \kappa T_e} \right)^2} \right) \right)} dv \quad (19b)$$

It is well known that this particular  $N_0(k)$  leads to instability if we also choose  $f_0(v) = \delta(v)$ . ("Instability of a cold plasmon gas" see ref. 2 and 3). In the case where  $f_0(v) \neq \delta(v)$  (assume  $f_0$  to be a Maxwellian for definiteness) we consider:

$$3 K^2 \kappa T_e + 2 \omega_p^2 m - 2 K m \omega_p \frac{\Omega}{q} = 0$$

$$K = \frac{m \omega_p \Omega}{3 q \kappa T_e} \pm \sqrt{\left( \frac{m \omega_p \Omega}{3 q \kappa T_e} \right)^2 - \frac{2 \omega_p^2 m}{3 \kappa T_e}}$$

When  $K$  has two real solutions the paranthesis in (19b) becomes negative for sufficiently large  $N_0$ , and  $K$  chosen appropriately. Obviously there is a threshold value for  $N_0$ . If  $Re \mathcal{E}(q, \Omega) = 0$  has solutions also we get unstable oscillations. Since the case where  $f_0 = \delta(v)$  is unstable for arbitrary  $K$  we notice a stabilizing effect of the thermal spread in ion velocities. This instability belongs to the class of parametric instabilities. For a review see for inst. F.F. Chen.<sup>8</sup>

III. Finally we consider the case where

$$N_0(k) = \begin{cases} 0 & \text{for } |k| > K \\ N_0 & \text{for } |k| \leq K \end{cases}$$

Then:

$$Re \mathcal{E}(q, \Omega) = 1 - \left( C_s^2 - \frac{\omega_p^2 m Q}{6 M \kappa T_e} \frac{2 M K}{K^2 - \left( \frac{\Omega \omega_p m}{3 q \kappa T_e} \right)^2} \right) P \int \frac{f_0'}{v - \frac{\Omega}{q}} dv \quad (20a)$$

$$Im \mathcal{E}(q, \Omega) = -P \int \frac{f_0'(\frac{\Omega}{q})}{\left( C_s^2 - \frac{\omega_p^2 m Q}{6 M \kappa T_e} \frac{2 M K}{K^2 - \left( \frac{\Omega \omega_p m}{3 q \kappa T_e} \right)^2} \right)} dv \quad (20b)$$

<sup>8</sup> and  $N_0$  large enough<sup>2</sup>

For physical reasons  $K$  is of the order  $1/d$  ( $d$  the Debye-length).

Then

$$\frac{\Omega}{q} \frac{\omega_p m}{3 \kappa T_e} \frac{1}{K} \approx \frac{\Omega}{q} \frac{1}{3 d \omega_p} = \frac{\Omega}{q} \frac{1}{3 v_{therm}} \ll 1$$

where  $v_{therm}$  is the electron thermal velocity. Equations (20a) and (20b) then reduces to

$$\operatorname{Re} \epsilon(q, \Omega) = 1 - \left( C_0^2 - \frac{\omega_p^2 m Q N_0}{3 M \kappa T_e K} \right) P \int \frac{f_0(v)}{v - \frac{\Omega}{q}} dv \quad (21a)$$

$$\operatorname{Im} \epsilon(q, \Omega) = -\pi f_0' \left( \frac{\Omega}{q} \right) \left( C_0^2 - \frac{\omega_p^2 m Q N_0}{3 M \kappa T_e K} \right) \quad (21b)$$

Assume for definiteness that  $f_0(v)$  is a Maxwellian. We may then make use of the plasma dispersion function<sup>9</sup> in calculating the integral in (21a). For small  $\Omega/q$  this integral is negative. If

$$N_0 \frac{\omega_p^2 m (1 - \frac{1}{2} (Kd)^2)}{6 M n_0 \kappa T_e K} > C_0^2 \quad (22)$$

then a Nyquist contour encircles the origin and the plasma is unstable.<sup>10,11</sup>

If  $K < 0.82 d^{-1}$  and

$$N_0 > \frac{(\kappa T_e)^2 C_0 \kappa}{\omega_p^2 m (1 - \frac{1}{2} (Kd)^2)} \quad (23)$$

the inequality (22) is satisfied.

The physical mechanism for this instability is easily understood: In a plasma where  $N_0 \neq 0$  an ion acoustic oscillation is driven by the E-field that builds up because of the electron pressure. This

E-field points from the high density region to the low density region of wave. When  $N_0 \neq 0$  the radiation pressure will hinder the electron motion from crest to trough of the wave. If the radiation pressure is strong enough it will overcome the electron pressure thus giving rise to an E-field in the opposite direction. This E-field will in turn force the ions towards regions of higher density and instability sets in. In the case where the two pressure terms cancel (equality in eq. (23)) an initial perturbation of the ions is damped solely due to ion free streaming.

We may introduce an effective electron temperature

$$\kappa T_{eff} = \left( \kappa T_e - N_0 \frac{\omega_p^2 m (1 - \frac{1}{2} (Kd)^2)}{6 n_0 \kappa T_e K} \right) \quad (24)$$

although this definition refers to a particular choice for  $N_0(k)$ .

The influence of a radiation pressure on double humped ion distributions, which are unstable when  $N_0=0$  is immediately realized by considering the stability diagram shown in for inst. ref. 12. This diagram shows that a distribution consisting of two ion groups with a sufficiently large velocity difference is unstable for large enough electron temperatures,  $T_e$ . Replacing  $T_e$  by  $T_{eff}$  from eq. (24) we see that the radiation pressure reduces the instability (reduces  $T_e$ ), eventually stabilizes the plasma. This fact may be of importance for the study of ion acoustic instabilities in for inst. Q-machines. In these devices double humped ion distributions are created by charge-exchange processes and the electron temperature is increased by RF- or  $\mu$ -wave heating<sup>13,14,15</sup>. These methods of electron heating are known to give rise to a high noise level<sup>16</sup> (high  $N_0$ ), considerably above the level of thermal fluctuations. The effect of "radiation pressure" should therefore be



taken into account in the analysis.

The condition (3) means (roughly) that the energy in the fluctuations should be larger than the thermal energy of the electrons thus rendering the instability mentioned somewhat academic. If we replace (7) by the dispersion relation for electromagnetic waves the basic physical mechanisms remains unchanged. The instability described may be observed for inst. when an intense laser beam interacts with a plasma.

### Discussion

In the foregoing sections we considered the influence of high frequency short wavelength oscillations on long wavelength ion acoustic waves. The question remains whether their effect is of any importance compared with the influence of the long wavelength electron oscillations. In this connection we draw the attention to the case of a cylindrical plasma column confined by a homogeneous magnetic field, for inst. the plasma in a conventional Q-machine. Geometrical effects are negligible for ion-acoustic oscillations. The dispersion relation for electron oscillations is on the other hand drastically changed for  $k \ll D^{-1}$  ( $D$  is diameter of plasma column), the dispersion curve (for the mode  $m = 0$ ) going through  $(\omega, k) = (0, 0)$  instead of  $(\omega, k) = (\omega_p, 0)$  as for infinite plasmas.<sup>17</sup> (For  $D^{-1} \ll k \ll d^{-1}$  eq. 7 is still a good approximation). We may now excite electron oscillations in the plasma around the plasma frequency. (For inst. by band limited white noise around  $\omega_p$ ). Because of this particular dispersion relation only large  $k$  will be excited and our analysis is appropriate (with properly chosen  $N_0(k)$ 's). Example II of the previous section has obvious applications also for an infinite plasma. Note that the instability in example III will be stabilized as time goes since the constant input of energy in the oscillations will heat up the electrons through nonlinear processes.

It should be noted that a force similar to  $F$  in eq. (6) acts on the ions also. This force is small since  $m \ll M$ , but might be taken into account in a more sophisticated theory. It is worth noting, however, that this force will try to move ions from low density regions towards regions of higher density thus having a

destabilizing effect (rather than stabilizing) on the instability discussed in example III of the previous section.

Finally we mention that the force on the quasi particles should properly read

$$-\frac{\partial \omega}{\partial x} = -\left(1 - \frac{1}{2}(kd)^2\right) \frac{\omega_p}{2n_0} \frac{\partial n_0}{\partial x} \quad (25)$$

rather than the expression used in eq. (15) where  $\frac{\partial n_i}{\partial x}$  was inserted. This distinction is irrelevant since we later assumed  $(dq)^2 \ll 1$  corresponding to the assumption of quasi-neutrality ( $n_e \approx n_i$ ). The expression in eq. (15) facilitates the calculations. If we apply (25) we obtain:

$$\begin{aligned} \mathcal{E}(q, \Omega) &= \frac{1}{(dq)^2} \left[ 1 + (dq)^2 - C_0^2 \int \frac{f_0(v)}{v - \frac{\Omega}{q}} dv \right. \\ &\quad \left. - \frac{1}{2m} \int \frac{Q(\omega)M_0'(k)}{2\gamma(k) - \frac{\Omega}{q}} dk \left( \omega_p d^2 + \frac{1}{2m} \int \frac{Q(\omega)M_0'(k)}{2\gamma(k) - \frac{\Omega}{q}} dk \right)^{-1} \right] \quad (26) \end{aligned}$$

It is easily shown that for  $(dq)^2 \ll 1$  the equation  $\mathcal{E}(q, \Omega) = 0$  will have the same solutions independent of which expression for  $\mathcal{E}(q, \Omega)$  is used: (26) or (16).

Note that we nowhere in the calculations make use of resonance conditions as  $k_1 = k_2 + k_3$  and  $\omega_1 = \omega_2 + \omega_3$ . These conditions can be satisfied for the dispersion relation mentioned earlier in this section if we let for inst.  $(\omega_2, k_2)$  represent an ion acoustic oscillation.

### Generalization and Conclusion

The analysis in the preceding sections was based on the dispersion relation (7) for the high frequency oscillations. We emphasize that the physical arguments holds equally well if electromagnetic oscillations are considered. The expressions for F in eq. (6) remains unchanged<sup>1,8</sup> and eq. (7) should be replaced by

$$\omega = \sqrt{\omega_p^2 + c^2 k^2} \quad (27)$$

Then

$$v_g = kc^2 (\omega_p^2 + c^2 k^2)^{-3/2}$$

and

$$\frac{\partial \omega}{\partial x} = \omega_p^2 (\omega_p^2 + c^2 k^2)^{-3/2} \frac{1}{n_0} \frac{\partial n}{\partial x}$$

The only important change is that now  $k \perp E_k$ . It is immediately realized (compare with the discussion p. 8-9) that a sufficiently strong radiation may drive an ion acoustic instability. Self-focusing of an intense laser beam is a related effect but does not require ion motion. This effect is easily understood using the dielectric function  $\mathcal{E} = 1 - (\omega_p/\omega)^2$ . A decrease in electron density (caused by the radiation pressure) is accompanied by an increase in  $\mathcal{E}$  thus causing the self-focusing. An intense beam of electromagnetic radiation with a large cross section (much larger than the Debye-length) and a uniform energy distribution is thus unstable when passing through a plasma ("filamentation"). The theory for these phenomena is well established (e.g. ref. 18,19).

We would like to point out that the previous analysis also

applies to the influence of high frequency ion oscillations (around  $\omega_{pi}$ ) on the propagation on low frequency, long wavelength-ion acoustic oscillations. We make use of the following set of equations:

The ion Vlasov equation with the radiation pressure (ponderomotive force) included

$$\frac{\partial f}{\partial t} + v \frac{\partial f}{\partial x} + \left( \frac{e}{m} E - \frac{e^2}{2em} \frac{\partial}{\partial x} \int \frac{N_k}{\omega} dk \right) f'_0(v) = 0 \quad (28)$$

The variation of  $N_k$  is given by

$$\frac{\partial N_k}{\partial t} + v_g \frac{\partial N_k}{\partial x} - \frac{\partial v}{\partial x} \frac{dN_k}{dk} = 0 \quad (29)$$

Since the frequency of the ion oscillations is low we may assume that the electrons are in equilibrium and use the Boltzmann law

$$E = - \frac{k T_e}{e} \frac{1}{n_0} \frac{\partial n}{\partial x} \quad (30)$$

In (30) we have used the assumption of quasi-neutrality i.e.  $n_e \approx n_i = n$ , since our assumptions require  $q \ll d^{-1}$  as in the previous analysis ( $q$  being the wavenumber of the low frequency, long wavelength oscillations). For the high frequency oscillations we use the dispersion relation

$$\omega = C_0 k (1 + (kd)^2)^{-1/2} \quad (31)$$

(We assume  $T_e \gg T_i$  in order to make (31) meaningful. Otherwise the waves will be damped within a period of oscillation for  $k \sim d^{-1}$  thus making the concept of a wavelike motion questionable). Using (31) we get

$$\frac{\partial v}{\partial k} \approx v_g = C_0 (1 + (kd)^2)^{-3/2} \quad (32)$$

and

$$\begin{aligned} \frac{\partial v}{\partial x} &= \frac{1}{2} C_0 k^2 d^2 (1 + (kd)^2)^{-3/2} \frac{1}{n_0} \frac{\partial n}{\partial x} \\ &\equiv A(k) \frac{\partial n}{\partial x} \end{aligned} \quad (33)$$

Since we assume that the wavelengths of the high frequency oscillations are small ( $k \sim d^{-1}$ ) we may replace  $\omega_k^{-1}$  in (28) by  $\omega_{pi}^{-1}$  so eq. (28) reduces to

$$\frac{\partial f}{\partial t} + v \frac{\partial f}{\partial x} + \left( \frac{e}{m} E - \frac{4\pi e^2}{2} \frac{\partial}{\partial x} \int N_k dk \right) f'_0(v) = 0 \quad (34)$$

We find that the zeroes of the dispersion relation are given by

$$1 - (C_0^2 + \frac{\omega_{pi}^2}{2} \int \frac{A(k) N'_k(k)}{v_g - \omega_k} dk) \int \frac{f'_0(v)}{v - \omega_k} dv = 0 \quad (35)$$

This equation is formally the same as (16) with  $(qd)^2 \ll 1$ . Apart from the difference in  $\frac{\partial v}{\partial k}$  and  $\frac{\partial v}{\partial x}$  the only change is that  $\omega_{pi}$  in (16) is replaced by  $\omega_{pi}$ . The effect of the ponderomotive forces is therefore rather weak in this case but otherwise it gives rise to the same effects as in the case considered before, in particular it has a stabilizing effect on unstable double humped ion-distribution functions. In spite of the similarity between the two problems in a linear treatment their nonlinear behaviour is entirely different, since electron oscillations are trapped in regions of low plasma density (wave troughs) while ion acoustic oscillations around  $\omega_{pi}$  will be trapped in regions of high density (wave crests). A correct treatment of these phenomena must take into account the problem of turning points in the WKB-approximation used in eq. (10). Such a treatment is of particular interest with

reference to soliton stability<sup>20</sup>. We shall not consider this problem here.

The influence of the turbulent spectrum of electron oscillations on low frequency long wavelength drift waves is investigated by Satya and Kaw<sup>21</sup>. One finds also in this case that the influence of the ponderomotive forces may lead to strong modifications of the wave behaviour. In particular the dispersion properties of the drift waves are drastically altered by even moderate amounts of short wavelength electron oscillations. We expect high frequency ion acoustic oscillations to give a similar, although weaker, effect (when  $T_e \gg T_i$ ).

Ponderomotive forces will, since they modify the linear dielectric function, also affect wave-wave interaction between the low frequency, long wavelength waves, through a modification of the coupling coefficients. This effect can be investigated by applying the theory of ref. 22 to the dielectric function (16). The coupling coefficients calculated in ref. 23 are easily modified using (24) for the case of a rectangular distribution for  $N_0(k)$ .

In conclusion we draw attention to the fact that the radiation pressure due to high frequency oscillations is a nonlinear effect. Fortunately it is possible to render the equations describing this effect linear by the introduction of  $N_k$ .

## References

- 1) A.A. Vedenov; Theory of Turbulent Plasma (Iliffe 1968)
- 2) R.Z. Sagdeev and A.A. Galeev; Nonlinear Plasma Theory (W.A. Benjamin 1969)
- 3) A.A. Vedenov and L.I. Rudakov, Sov.Phys. Doklady 9, 1073 (1965)
- 4) L.P. Pitaevskii, Sov.Phys. JETP 12, 1008 (1961)
- 5) B.B. Kadomtsev, Plasma Turbulence (Academic Press 1965)
- 6) S.A. Andersen, G.B. Christoffersen, V.O. Jensen, P. Michelsen and P. Nielsen, Phys.Fluids 14, 990 (1971)
- 7) P. Michelsen and H.L. Pécseli, Phys.Fluids 16, 221 (1973)
- 8) F.F. Chen; Introduction to Plasma Physics (Plenum 1974)
- 9) B. Fried and S. Conte, The Plasma Dispersion Function (Academic Press, 1961)
- 10) O. Penrose, Phys.Fluids 3, 258 (1960)
- 11) J.D. Jackson, J.Nucl.Energy, Part C: Plasma Physics, 1, 171 (1960)
- 12) B.D. Fried and A.Y. Wong, Phys.Fluids 9, 1084 (1966)
- 13) D.R. Baker, Phys.Rev.Lett. 28, 1189 (1972)
- 14) G.B. Christoffersen and L.P. Prahm, Phys.Fluids 16, 708 (1973)
- 15) G.B. Christoffersen and L.P. Prahm, Plasma Physics 14, 1140 (1972)
- 16) H.L. Pécseli and P.I. Petersen, Risø Rep. No. 290 (1973)
- 17) P.I. Barrett, H.G. Jones and R.N. Franklin, Plasma Phys. 10, 911 (1968)
- 18) M.S. Sodha, D.P. Tewari, Awdhesh Kumar and V.K. Tripathi, J.Phys. D: Appl. Phys. 7, 345 (1974)

- 19) K.B. Dysthe, Phys. Letters 27A, 59 (1968)
- 20) V.E. Zakharov, Sov. Phys. JETP, 35, 908 (1972)
- 21) Y.S. Satya and P.K. Kaw, Phys. Rev. Lett. 31, 1453 (1973)
- 22) L. Stenflo, Plasma Phys. 12, 509 (1970)
- 23) H.L. Pécseli, Physica Scripta (in press).