



The range of attraction for light traps catching *Culicoides* biting midges

Græsbøll, Kaare; Kirkeby, Carsten; Bødker, Rene; Stockmarr, Anders; Christiansen, Lasse Engbo; Lind, Peter

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Græsbøll, K., Kirkeby, C., Bødker, R., Stockmarr, A., Christiansen, L. E., & Lind, P. (2013). *The range of attraction for light traps catching Culicoides biting midges*. Poster session presented at GeoVet 2013, London, United Kingdom.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The range of attraction for light traps catching *Culicoides* biting midges

Kaare Græsbøll¹, Carsten Kirkeby², René Bødker², Anders Stockmarr¹, Lasse E. Christiansen¹, and Peter Lind²
¹Department of Applied Mathematics and Computer Science. ²National Veterinary Institute. Technical University of Denmark (DTU)


View from above:
The light tube generates an asymmetric light-field around a trap.



Lighter colors indicate higher light intensity.

Conclusion

We present a model that explains observed catches of *Culicoides* by assuming that they evaluate light sources in the horizon and fly towards the perceived strongest source of light. The model implicates that the range of attraction for a light trap is dependent on the intensity of the light emitted from the trap. Furthermore, traps placed closely together will collaborate to attract *Culicoides* from farther away. The maximum range of attraction for a single 4W CDC light trap is estimated to be 15.25 meters.




Two traps in the same location results in increased light intensity and therefore also increased range of attraction.

Experiment




10 light traps were placed on a line in a symmetric pattern with less distance towards the middle. In total 10,150 *Culicoides* were caught in 242 individual trap catches in 16 catch periods of one hour. Surprisingly the two central traps caught the highest fractions of *Culicoides*.



Result

Data was transformed to fraction of catch per catch period, thus avoiding influence of fluctuations in the level of abundance. The symmetrized experimental data is plotted as a boxplot with fractional catch as a function of trap number.

The red line shows the best fit of the model to the data, where the maximum range of attraction for a single trap is estimated to be 15.25 m.





Model

We developed a simulation model to fit the observed experimental data. The model simulates the flight of *Culicoides* towards the light traps. The light intensity from the traps is calculated up to at least 100 m away from any trap. Included in this calculation is that light is emitted asymmetrically from the light tubes in the traps.

Left figure: The experimental field visualized from above, with trap locations indicated by black dots: The colors show the maximum perceived light intensity by the *Culicoides* around the transect of light traps.

The direction of flight for each *Culicoides* is determined by the direction where they perceived the brightest light to originate, which most often is not in a straight line towards the traps.

Right figure: The experimental field visualized from above: The catch area for each trap is marked with an individual color. The center traps catch *Culicoides* from a large area because of the synergistic effect. The white lines represent the three different light intensity cutoff values used to determine the fits in the 'Result' figure.



For more information contact:



Modeler:

Field expert:

Kaare Græsbøll
kagr@dtu.dk

Carsten Kirkeby
ckir@vet.dtu.dk

Background picture is of the experimental setup
Taken by: Carsten Kirkeby

The research presented in this poster is published as:
"The range of attraction for light traps catching biting midges (Diptera: Ceratopogonidae: *Culicoides*)"
Parasites & Vectors, in press 2013

Try our WebApp!

It calculates catch from a setup of traps that you decide.

