



## A Two-Dimensional Electron Gas at the Spinel/Perovskite Interface of $\gamma\text{-Al}_2\text{O}_3/\text{SrTiO}_3$ with Carrier Mobility Exceeding $100,000 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$

Chen, Yunzhong; Bovet, N.; Trier, Felix; Christensen, Dennis Valbjørn; Qu, F.M.; Andersen, Niels Hessel; Kasama, Takeshi; Zhang, Wei; Giraud, R.; Dufouleur, J.

Total number of authors:

19

Publication date:

2013

[Link back to DTU Orbit](#)

### Citation (APA):

Chen, Y., Bovet, N., Trier, F., Christensen, D. V., Qu, F. M., Andersen, N. H., Kasama, T., Zhang, W., Giraud, R., Dufouleur, J., Jespersen, T. S., Sun, J. R., Smith, A., Nygård, J., Lu, L., Büchner, B., Shen, B. G., Linderoth, S., & Pryds, N. (2013). A Two-Dimensional Electron Gas at the Spinel/Perovskite Interface of  $\gamma\text{-Al}_2\text{O}_3/\text{SrTiO}_3$  with Carrier Mobility Exceeding  $100,000 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$ . Abstract from 20th Workshop on Oxide Electronics<sup>3</sup> (WOE20), Singapore, Singapore.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

O-21

## A Two-Dimensional Electron Gas at the Spinel/Perovskite Interface of $\gamma\text{-Al}_2\text{O}_3/\text{SrTiO}_3$ with Carrier Mobility Exceeding $100,000 \text{ cm}^2\text{V}^{-1}\text{s}^{-1}$

**Y. Z. Chen<sup>1\*</sup>, N. Bovet<sup>2</sup>, F. Trier<sup>1</sup>, D. V. Christensen<sup>1</sup>, F. M. Qu<sup>3</sup>, N. H. Andersen<sup>4</sup>, T. Kasama<sup>5</sup>, W. Zhang<sup>1</sup>, R. Giraud<sup>6,8</sup>, J. Dufouleur<sup>6</sup>, T. S. Jespersen<sup>7</sup>, J. R. Sun<sup>3</sup>, A. Smith<sup>1</sup>, J. Nygård<sup>7</sup>, L. Lu<sup>3</sup>, B. Büchner<sup>6</sup>, B. G. Shen<sup>3</sup>, S. Linderoth<sup>1</sup>, and N. Pryds<sup>1</sup>**

<sup>1</sup>Department of Energy Conversion and Storage, Technical University of Denmark, Risø Campus, 4000 Roskilde, Denmark

<sup>2</sup>Nano-ScienceCenter, Department of Chemistry, University of Copenhagen, 2100 Copenhagen, Denmark

<sup>3</sup>Institute of Physics, Chinese Academy of Sciences, 100190 Beijing, China

<sup>4</sup>Department of Physics, Technical University of Denmark, 2800 Lyngby, Denmark

<sup>5</sup>Center for Electron Nanoscopy, Technical University of Denmark, 2800 Lyngby, Denmark

<sup>6</sup>Leibniz Institute for Solid State and Materials Research, IFW Dresden, D-01171 Dresden, Germany

<sup>7</sup>Niels Bohr Institute, University of Copenhagen, 2100 Copenhagen, Denmark

<sup>8</sup>Laboratoire de Photonique et de Nanostructures-CNRS, Route de Nozay, 91460 Marcoussis, France

\*E-mail: [yunc@dtu.dk](mailto:yunc@dtu.dk)

The discovery of two-dimensional electron gases (2DEGs) at the heterointerface between two insulating perovskite-type oxides, such as  $\text{LaAlO}_3$  and  $\text{SrTiO}_3$  [1], provides opportunities for a new generation of all-oxide electronic devices [2]. Key challenges remain for achieving interfacial electron mobilities much beyond the current value of approximately  $1000 \text{ cm}^2\text{V}^{-1}\text{s}^{-1}$  (at low temperatures). Here, we create a new type of 2DEG at the heterointerface between  $\text{SrTiO}_3$  and a spinel  $\gamma\text{-Al}_2\text{O}_3$  epitaxial film with compatible oxygen ions sublattices [3]. Electron mobilities more than one order of magnitude higher than those of hitherto investigated perovskite-type interfaces are obtained. Particularly, electron Hall mobilities as large as  $1.4 \times 10^5 \text{ cm}^2\text{V}^{-1}\text{s}^{-1}$  and  $n_s$  as high as  $3.7 \times 10^{14} \text{ cm}^{-2}$  at 2 K is obtained at this spinel/perovskite complex oxide interface [3]. The spinel/perovskite 2DEG, where the two-dimensional conduction character is revealed by quantum magnetoresistance oscillations, is found to be strongly coupled to interface-stabilized oxygen vacancies confined within a layer of 0.9 nm in proximity to the interface. Our findings pave the way for studies of mesoscopic physics with complex oxides and design of high-mobility all-oxide electronic devices.

A. Ohtomo & H. Y. Hwang, *Nature* 427, 423-426 (2004).

J. Mannhart & D. G. Schlom, *Science* 327, 1607-1611 (2010).

Y. Z. Chen et al. *Nat. Commun.* 4:1371, doi: 10.1038/ncomms2394 (2013).