



## Maternal high-fat/high-sucrose diet during lactation results in increased adipose tissue mass, and altered hepatic fatty acid metabolism at weaning

Ingvorsen, Camilla; Hellgren, Lars

*Publication date:*  
2012

[Link back to DTU Orbit](#)

*Citation (APA):*

Ingvorsen, C., & Hellgren, L. (2012). *Maternal high-fat/high-sucrose diet during lactation results in increased adipose tissue mass, and altered hepatic fatty acid metabolism at weaning*. Abstract from Benzon Symposium No. 58 Adipose Tissue in Health and Disease, Copenhagen, Denmark.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **Maternal high-fat/high-sucrose diet during lactation results in increased adipose tissue mass at weaning.**

**Camilla Ingvorsen<sup>1,2</sup> and Lars I. Hellgren<sup>1</sup>**

<sup>1</sup>Technical University of Denmark, Department of Systems Biology

<sup>2</sup>Centre for Fetal Programming, Copenhagen, Denmark

**Aim** Maternal nutrition during gestation and lactation has been implicated as a factor that might modulate the metabolic risk of the offspring. Therefore, we have investigated how maternal high-fat/high sucrose diet during gestation and lactation effect weight and weight gain during lactation as well as metabolic status at weaning.

**Methods** Female obese-prone rats were fed a conventional (C) or a high fat/high sucrose (HFHS) diet for 17 weeks before and during gestation and lactation. All female pups are sacrificed at day 0 and male pups are cross-fostered during lactation giving 4 groups of offspring: (maternal diet during gestation/maternal diet during lactation) C/C, C/HF, HF/C and HF/HF. Male pups are then sacrificed at weaning at 4 weeks of age.

**Results** There were no difference in weight and metabolic status of the dams before or after gestation and lactation. The C born pups had a higher birth weight ( $p=0.06$ ), and a significantly higher weight gain the first week resulting in a higher body weight at 1 and 2 weeks of age. From 2 weeks of age, the diet of the lactating dam dominates the effect on the offspring's growth rate. Although there were no difference in fat content of the milk, pups lactated by HFHS dams have an increased weight gain during 3th and 4th week of lactation, resulting in increased adipose tissue mass and circulating leptin at weaning. Interestingly, at four weeks of age, all groups that had been either born or lactated by HFHS-fed dam's had increased blood glucose, but only pups both born and lactated by HFHS-dams had significantly increased HOMA-IR. Furthermore, based on substrate/product ratios, it is evident that both pre- and postnatal HFHS-intake by the dams affected the activity of hepatic  $\Delta 5$ - and  $\Delta 6$ -desaturases, and hence potentially the long-chained n-6 and n-3 PUFA levels in the offspring.