



Expression of immune-related genes during wound healing in fish

Schmidt, Jacob; Nielsen, Michael Engelbrecht

Published in:
DAFINET Workshop and PhD Course

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Schmidt, J., & Nielsen, M. E. (2013). Expression of immune-related genes during wound healing in fish. In *DAFINET Workshop and PhD Course: Book of Abstracts*

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Expression of immune-related genes during wound healing in fish

Schmidt J.G., Nielsen M.E.

*National Food Institute, Biological Quality Research Group,
Technical University of Denmark, Søborg, Denmark*

Sterile tissue damage triggers the same pathways through damage-associated molecular patterns (DAMPs) that an infection does through pathogen-associated molecular patterns (PAMPs), albeit with different kinetics of expression of the involved genes. The cascade of reactions initiated by tissue damage starts with a series of non-transcriptional responses that leads to vasoconstriction and hemostasis. This is usually followed by an inflammatory response also initiated in the absence of transcription, but later greatly enhanced by expression of genes coding for proinflammatory cytokines and proteases. Inflammation is aimed at eradicating invading pathogens but causes concomitant tissue damage and this compromises regeneration. A strong inflammatory response often results in fibrosis and poor repair of the wound site, and is thus disadvantageous if the wound is uninfected. However, the inflammatory response is limited in early life stages of most animals, and these heal with a much higher degree of regeneration.

I will be presenting a selection of results from my PhD studies with a focus on the response to wounding during ontogeny of carp larvae and juveniles aged 7-56 days post-fertilization. The wounds of larvae completely regenerate within 3 days, whereas the wounds in juveniles are still visible 7 days post-wounding. Wound-induced changes in gene expression are limited, which may be ascribed to similarities between natural morphogenesis and tissue repair and that the investigated genes are thus already transcribed at adequate levels. However, expression of the mucosal immunoglobulin IgZ1 is upregulated in wounded larvae, indicating important innate effector functions of this isoform during early ontogeny.

Presenting author: Jacob Schmidt, jacsc@food.dtu.dk