



## Furan in food including homemade and ready-to-eat food products

Fromberg, Arvid; Granby, Kit; Mariotti Celis, M.; Pedreschi, F.

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Fromberg, A., Granby, K., Mariotti Celis, M., & Pedreschi, F. (2013). *Furan in food including homemade and ready-to-eat food products*. Abstract from Årsmøde i Den Nordiske Metodikkomité for Levnedsmidler (NMKL) 2013, Kalmar, Sweden.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Furan in food including homemade and ready-to-eat food products

A. Fromberg<sup>1</sup>, K. Granby<sup>1</sup>, M. Mariotti Celis<sup>2</sup>, F. Pedreschi<sup>2</sup>

<sup>1</sup>Technical University of Denmark, National Food Institute, Copenhagen, Denmark.

<sup>2</sup>Pontificia Universidad Catolica de Chile. Department of Chemical Engineering and Bioprocesses, Santiago, Chile

**Furan is formed in canned, jarred or browned food items. As furan is carcinogenic in animal experiments, attention has been drawn to the presence in commercial and home-cooked foods. The formation of furan in home cooked dishes and snacks (in between meals) were studied as well as the stability of furan during cooking, saving and reheating of meals.**

## Introduction

Furan is formed e.g. in Maillard reactions. Several compounds may be precursors of furan formation during food processing and preparation e.g. sugars, amino acids, ascorbic acid (vitamin C) and polyunsaturated fatty acids<sup>1</sup>.

Although exposure studies show that coffee is the main source of furan for adult coffee consumers<sup>2,3</sup>, mitigation in other foods may be relevant to reduce the intake of this possible carcinogenic compound.

As

- furan is formed in canned, jarred and roasted foods
- furan has a low boiling points (31.4°C)

the question was

**What is the stability of furan during cooking, saving or reheating of meals?**

## Materials & methods

Recipes representative for European dietary habits and with potential of furan contents were selected to study the occurrence of furan in freshly prepared home cooked dishes. Dishes with preprocessed ingredients and snack (between meals) were also included in the survey. Furthermore the stability of furan during cooking, saving and reheating of meals were studied. Analyses of furan was made by GC-MS with a headspace autosampler, vials heated to 60°C for 30 min. Furan was detected at  $m/z$  68 and 39; The internal standard  $d_4$ -furan at  $m/z$  72 and 42<sup>2</sup>

## Results

Most of the homemade dishes which did not include browning of carbohydrates, did not generate furan<sup>2</sup>. When carbohydrate rich foods e.g. potato crisps were browned, furan was generated. For example when deep frying potato crisps (diameter 40 mm x width 2 mm) in oil at 150°, 170° and 190°C to the same water content (crispiness) of ~ 2% simulating industrial conditions, the furan content increased from 12-15 ng/g up to 31-52 ng/g (Fig. 1).

### Stability of furan during cooking and reheating

Industrially made soups with furan, which were cooked, left to cool for 60 min and reheated, showed some loss of furan during cooking (up to the half), but during cooling furan was more stable and still ~ 50% remains when the meals were reheated (Fig. 2).

### Snacks and breakfast cereals

Furan was found in some sweets and snacks. The presence in chocolate, crisps, cookies and popcorn may be due to the Maillard reactions during heat-treatment. However, Maillard reactions also take place during drying of e.g. for fruits at moisture contents ~ 5-30%. Hence this may explain the furan conc. of up to 83 ng/g in raisins and also traces in dried dates, pineapples, plums and bananas. For breakfast cereals one sample contained 387 ng/g furan while the others were below 87 ng/g (n=11). Children with high consumption of breakfast cereals may get a relatively large proportion of the furan exposure from this food category.

## Conclusion

Dishes prepared from industrial products with furan may contain furan, while homemade dishes prepared without browning contain less. Browning of carbohydrates may generate furan and dried fruit is a furan source.

Fig.1. The occurrence of furan in potato crisps increased with increasing frying temperature.




Fig. 2. Furan is stable in the foods after preparation and even after reheating >50% is still present<sup>2</sup>




Table 1. Furan conc. in sweets, snacks & breakfast cereals

Food item	ng/g
Chocolate(2)	3.8-11
Cookies(2)	5.3-8.8
Macaroon	3.6
Potato crisps(7)	7.6-39 (mean 20)
Corn crisps(5)	5.5-26 (mean 15)
Popcorn(2)	36-91
Banana crisps	10.8
Peanuts	2.9
Hazelnut	<2.4
Raisins(2)	2.5-83
Plums dried	9.6
Pineapple dried	6.3
Dates sun dried	2.4
Oat, müsli(2)	3.0-5.9
Whole grain cereals(3)	6.1-43 (mean 29)
Corn flakes	87
Honey cereals(2)	61-387

## References:

<sup>1</sup>Limacher A., Kerler J., Conde-Petit B., Blank I. 2007. Formation of furan and methylfuran from ascorbic acid in model systems and food. Food Additives and Contaminants, 24 (S1): 122-135

<sup>2</sup>Fromberg A., Fagt S., Granby K. 2009. Furan in heat processed food products including home cooked food products and ready-to-eat products, Rep. made to EFSA-Q-2009-00846.

<sup>3</sup>EFSA 2011 Update on furan levels in food from monitoring years 2004-2010 and exposure assessment. Rep. 9(9):2347 33pp available from [www.efsa.europa.eu/efsajournal](http://www.efsa.europa.eu/efsajournal)