



Only small fractions of soluble β -glucan modulate the mucosal immune system in carp (*Cyprinus carpio* L.)

Przybylska, Dominika Alicja; Nielsen, Michael Engelbrecht

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Przybylska, D. A., & Nielsen, M. E. (2011). *Only small fractions of soluble β -glucan modulate the mucosal immune system in carp (*Cyprinus carpio* L.)*. Poster session presented at 12th Fish Immunology Workshop, Wageningen, Netherlands.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Dominika Przybylska and Michael Engelbrecht Nielsen

DTU, National Food Institute, Section for Aquatic Protein Biochemistry,
Søltofts Plads, Building 221, DK – 2800 Kgs. Lyngby, Denmark

Introduction

β -glucans are well known for their ability to modulate the piscine immune system. The discrepant effects of different origins and solubility of β -glucans on immune function possibly relates to factors as administration route, molecular weight and water solubility.

Results




Fig. 1a Water - soluble unbranched 6.3kD β -glucan.

Fig. 1b Water - soluble unbranched 15kD β -glucan.

Expression of interleukin 1 β (IL-1 β), interleukin 6 (IL-6) and heat shock protein 70 (hsp70) in skin and gill collected from juvenile carps. Fish were immersed with soluble β -glucan (1 μ g/ml) for 0h, 2h, 4h, 8h and 24h. Results are obtained by RQ-PCR and expressed relative to control at each time point and corrected for 40S expression. NS – no samples.