



EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA); Scientific Opinion on principles for deriving and applying Dietary Reference Values

EFSA Publication

Link to article, DOI:
[10.2903/j.efsa.2010.1458](https://doi.org/10.2903/j.efsa.2010.1458)

Publication date:
2010

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
EFSA Publication (2010). *EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA); Scientific Opinion on principles for deriving and applying Dietary Reference Values*. European Food Safety Authority. the EFSA Journal No. 1458 <https://doi.org/10.2903/j.efsa.2010.1458>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

SCIENTIFIC OPINION

Scientific Opinion on principles for deriving and applying Dietary Reference Values¹

EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA)^{2, 3}

European Food Safety Authority (EFSA), Parma, Italy

ABSTRACT

This Opinion of the EFSA Panel on Dietetic products, Nutrition, and Allergies (NDA) deals with the general principles for development and application of Dietary Reference Values (DRVs). These quantitative reference values for nutrient intakes for healthy individuals and populations are based on health criteria. Derived from DRVs, nutrients goals and recommendations take into account other criteria such as food composition or dietary habits, and may be used for assessment and planning of diets. It is proposed to derive the following DRVs: 1) Population Reference Intakes (PRI), 2) Average Requirement (AR), 3) Lower Threshold Intake (LTI), 4) Adequate Intake (AI), 5) Reference Intake ranges for macronutrients (RI). Nutrient requirements differ with age, sex and physiological condition. The Panel proposes to define the age ranges used for each nutrient on a case-by-case basis depending on the available data. For the age group < 6 months requirements are considered to be equal to the supply from breast- milk, except in those cases where this does not apply. Separate reference values will be established for pregnant and lactating women. Interpolation or extrapolation between population groups will be used in instances where no data are available for defined age and sex groups.

KEY WORDS

Dietary Reference Values (DRVs), Population Reference Intakes (PRI), Reference Intake ranges for macronutrients (RI), Average Requirement (AR), Lower Threshold Intake (LTI), Adequate Intake (AI), principles.

1 On request from the European Commission, Question No EFSA-Q-2008-463, adopted on 04 December 2009.

2 Panel members: Carlo Agostoni, Jean-Louis Bresson, Susan Fairweather-Tait, Albert Flynn, Ines Golly, Hannu Korhonen, Pagona Lagiou, Martinus Løvik, Rosangela Marchelli, Ambroise Martin, Bevan Moseley, Monika Neuhäuser-Berthold, Hildegard Przyrembel, Seppo Salminen, Yolanda Sanz, Sean (J.J.) Strain, Stephan Strobel, Inge Tetens, Daniel Tomé, Hendrik van Loveren and Hans Verhagen.

Correspondence: nda@efsa.europa.eu

3 Acknowledgement: The Panel wishes to thank for the preparation of this Opinion: Wulf Becker, Henk van den Berg, Karin Hulshof, Albert Flynn, Alan Jackson, Ambroise Martin, and Hildegard Przyrembel.

Suggested citation: EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA); Scientific Opinion on principles for deriving and applying Dietary Reference Values. EFSA Journal 2010; 8(3):1458. [30 pp.]. doi:10.2903/j.efsa.2010.1458. Available online: www.efsa.europa.eu

SUMMARY

The European Commission has requested EFSA to review the existing advice of the Scientific Committee on Food on Population Reference Intakes for energy, nutrients and other substances with a nutritional or physiological effect. These reference values date from 1993. Since then new scientific data have become available for some of the nutrients, and scientific advisory bodies in many European Union Member States and in the United States have reported on recommended dietary intakes.

This Opinion focuses on the general principles for development and application of Dietary Reference Values (DRVs) - quantitative reference values for nutrient intakes for healthy individuals and populations which may be used for assessment and planning of diets.

Similarly to the earlier Scientific Committee on Food (SCF) report in 1993 the Panel proposes to derive the following Dietary Reference Values:

- *Population Reference Intakes (PRI)*: the level of (nutrient) intake that is adequate for virtually all people in a population group.
- *Average Requirement (AR)*: the level of (nutrient) intake that is adequate for half of the people in a population group, given a normal distribution of requirement.
- *Lower Threshold Intake (LTI)*: the level of intake below which, on the basis of current knowledge, almost all individuals will be unable to maintain “metabolic integrity”, according to the criterion chosen for each nutrient.

In addition, the Panel also proposes to derive the following Dietary Reference Values :

- *Adequate Intake (AI)*: the value estimated when a Population Reference Intake cannot be established because an average requirement cannot be determined. An Adequate Intake is the average observed daily level of intake by a population group (or groups) of apparently healthy people that is assumed to be adequate.
- *Reference Intake ranges for macronutrients (RI)*: the intake range for macronutrients, expressed as % of the energy intake. These apply to ranges of intakes that are adequate for maintaining health and associated with a low risk of selected chronic diseases.

The Panel will not address the Tolerable Upper Intake Level (UL) as this has been assessed previously. The Tolerable Upper Intake Level is the maximum level of total chronic daily intake of a nutrient (from all sources) judged to be unlikely to pose a risk of adverse health effects to humans⁴.

Some of the Reference Values - the Average Requirement, Population Reference Intake and the Lower Threshold Intake - relate to nutrient requirements that are defined by specific criteria of nutrient adequacy. In defining nutrient requirements the selection of criteria to establish nutrient adequacy is an important step. For most nutrients a hierarchy of criteria for nutrient adequacy can be established, ranging from prevention of clinical deficiency to optimisation of body stores, or status. Which criterion, or combination of criteria, will be the most appropriate will be decided on a case-by-case basis.

Within any lifestage group, nutrient requirements vary between individuals and the Average Requirement, Population Reference Intake and Lower Threshold Intake represent different points on

⁴ An opinion on tolerable upper intake levels for vitamins and minerals was published in February 2006 (http://www.efsa.europa.eu/en/science/nda/nda_opinions.html).

the distribution of individual requirements. Nutrient requirements also differ with age, sex and physiological condition, due to differences in the velocity of growth for the younger age groups, and age-related changes in nutrient absorption and body functions and/or functional capacity, such as renal function. Especially in older subjects, variability in functional capacity and in energy expenditure appears higher than in younger adults, particularly for elderly above 75 years.

Because of this, Dietary Reference Values are developed for different life stage and sex groups. The Panel proposes to define the age ranges used for each nutrient on a case-by-case basis depending on the available data. For the age group <6 months requirements are considered to be equal to the supply from breast-milk, except on a case-by-case basis where this does not apply. Separate reference values will be established for pregnant and lactating women, taking into account the additional nutrient requirement for the formation of new tissues, or to compensate for the nutrients lost to the body in the form of human milk, respectively, and considering the physiological adaptations that occur during these conditions.

Interpolation or extrapolation between population groups will be used in instances where no data are available for defined age and sex groups. Scaling methods using isometric (linear with body weight) or allometric (body weight to the power of a chosen exponent) or interpolation based on other non predefined parameters are being used. Which method is the most appropriate will be decided on a case-by-case basis.

Reference heights and weights are useful when more specificity about body size and nutrient requirements are needed than that provided by life stage categories. In the absence of more recent data, reference weights will be the same as in the SCF report, and for children <1 year, as established by the WHO for fully breastfed infants.

Dietary reference values can be used for different purposes, such as in diet assessment and diet planning, both at the population and individual level, but also as a basis for reference values in food labelling, and in establishing food based dietary guidelines.

In dietary assessment of groups the Average Requirement can be used to estimate the prevalence of inadequate intakes of micronutrients (the Average Requirement cut-point method), if the distribution of nutrient intakes is normal, and intakes are independent from requirements. The Population Reference Intake should not be used for this purpose as this would result in overestimation of the proportion of the group at risk of inadequacy. Probabilistic methods, taking into account both the intake and requirement variation might be used as an alternative, and in case distributions are skewed.

For macronutrients with a defined reference intake range for individuals, the distribution of usual intake of individuals may be assessed to ascertain what proportion of the group lies outside the reference lower and upper limits of the range. In case of energy, the mean usual intake of energy of a defined group, relative to the average requirement, may be used in assessing the adequacy.

For assessment of adequacy of nutrient intakes in individuals Dietary Reference Values are of limited use. Usual intakes below the AR are likely inadequate, and below the Lower Threshold Intake very probably inadequate, while chronic intakes above the Tolerable Upper Intake Level may be associated with an increased risk of adverse effects. For a valid assessment of the adequacy of an individual's usual intake, combined information with anthropometric, biochemical (status) and clinical data is needed.

In dietary planning for groups the usual intake distribution should be between the AR and UL to avoid inadequate, respectively excessive intakes. For nutrients such as vitamins, minerals, and protein, the PRI can be a practical starting point. However, target median intakes higher than the Population Reference Intake might be considered, especially in case of a skewed intake distribution. For macronutrients the distribution of usual intake of individuals should be such as to minimise the

proportion of the group that lies outside the reference lower and upper limits of the range. For energy, the reference intake (estimated average energy requirement) of the group based on sex, age, height, weight, and physical activity level of the group may be used as a planning goal.

The goal of planning diets for individuals is to have a low probability of inadequacy while minimising potential risk of excess for each nutrient. For nutrients such as vitamins, minerals, and protein, this is done by ensuring that the usual intake meets the Population Reference Intake or Adequate Intake while not exceeding the Tolerable Upper Intake Level. Population Reference Intakes would be an overestimation for most individuals. For macronutrients which have a reference intake range, the usual intake of individuals should be between the lower and upper bounds of the reference range. For energy, the reference intake (average energy requirement) based on an individual's sex, age, height, weight, and physical activity level may be used as an initial planning goal; however, body weight must be monitored and intake adjusted as appropriate.

TABLE OF CONTENTS

Abstract	1
Summary	2
Table of contents	5
Background as provided by the European Commission	6
Terms of reference as provided by European Commission	6
Assessment	8
1. Introduction	8
2. General principles for deriving dietary recommendations	9
2.1. Estimating the physiological requirement and metabolic demand	9
2.2. Establishing the dietary requirement of nutrients	9
2.3. Establishing dietary reference values	9
2.4. Establishing nutrient goals and recommendations	10
2.5. Establishing food-based dietary guidelines	10
3. Terminology and definitions	10
4. Conceptual basis for derivation of dietary reference values	12
4.1. Population Reference Intakes (PRI)	12
4.2. Average Requirement (AR)	13
4.3. Lower Threshold Intake (LTI)	13
4.4. Adequate Intake (AI)	13
4.5. Reference Intake ranges for macronutrients (RI)	14
4.6. Tolerable Upper intake Level (UL)	14
5. Methods for determining dietary reference values – types of data used	14
5.1. Criteria for assessment of nutrient requirement	14
5.2. Methods for determining nutrient requirements	15
5.3. Diet and health relationship: risk of chronic disease	16
5.4. Interpolation or extrapolation between population groups	17
5.5. Factors that affect nutrient requirements	17
5.6. Lifestage and sex	17
6. Application of dietary reference values for nutrients	18
6.1. Assessment of the risk of (in)adequacy of nutrient intake	18
6.1.1. Assessment of the risk of (in)adequacy of nutrient intake in populations	18
6.1.2. Assessment of the risk of (in)adequacy of nutrient intake in individuals	20
6.2. Dietary planning	20
6.2.1. Dietary planning for groups	20
6.2.2. Dietary planning for individuals	21
6.3. Labelling reference intake values	21
6.4. Food-based dietary guidelines	22
Conclusions	22
References	24
Appendices	28
Glossary / Abbreviations	30

BACKGROUND AS PROVIDED BY THE EUROPEAN COMMISSION

The scientific advice on nutrient intakes is important as the basis of Community action in the field of nutrition, for example such advice has in the past been used as the basis of nutrition labelling. The Scientific Committee for Food (SCF) report on nutrient and energy intakes for the European Community dates from 1993. There is a need to review and if necessary to update these earlier recommendations to ensure that the Community action in the area of nutrition is underpinned by the latest scientific advice.

In 1993, the SCF adopted an opinion on the nutrient and energy intakes for the European Community⁵. The report provided reference intakes for energy, certain macronutrients and micronutrients, but it did not include certain substances of physiological importance, for example dietary fibre.

Since then new scientific data have become available for some of the nutrients, and scientific advisory bodies in many European Union Member States and in the United States have reported on recommended dietary intakes. For a number of nutrients these newly established (national) recommendations differ from the reference intakes in the SCF (1993) report. Although there is considerable consensus between these newly derived (national) recommendations, differing opinions remain on some of the recommendations. Therefore, there is a need to review the existing EU reference intakes in the light of new scientific evidence, and taking into account the more recently reported national recommendations. There is also a need to include dietary components that were not covered in the SCF opinion of 1993, such as dietary fibre, and to consider whether it might be appropriate to establish reference intakes for other (essential) substances with a physiological effect.

In this context the EFSA is requested to consider the existing population reference intakes for energy, micro- and macronutrients and certain other dietary components, to review and complete the SCF recommendations, in the light of new evidence, and in addition advise on a population reference intake for dietary fibre.

For communication of nutrition and healthy eating messages to the public it is generally more appropriate to express recommendations for the intake of individual nutrients or substances in food-based terms. In this context the EFSA is asked to provide assistance on the translation of nutrient based recommendations for a healthy diet into food based recommendations intended for the population as a whole.

TERMS OF REFERENCE AS PROVIDED BY EUROPEAN COMMISSION

In accordance with Article 29 (1)(a) and Article 31 of Regulation (EC) No. 178/2002, the Commission requests EFSA to review the existing advice of the Scientific Committee for Food on population reference intakes for energy, nutrients and other substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

In the first instance the EFSA is asked to provide advice on energy, macronutrients and dietary fibre. Specifically advice is requested on the following dietary components:

- Carbohydrates, including sugars;

⁵ Scientific Committee for Food, Nutrient and energy intakes for the European Community, Reports of the Scientific Committee for Food 31st series, Office for Official Publication of the European Communities, Luxembourg, 1993.

- Fats, including saturated fatty acids, poly-unsaturated fatty acids and mono-unsaturated fatty acids, *trans* fatty acids;
- Protein;
- Dietary fibre.

Following on from the first part of the task, the EFSA is asked to advise on population reference intakes of micronutrients in the diet and, if considered appropriate, other essential substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

Finally, the EFSA is asked to provide guidance on the translation of nutrient based dietary advice into guidance, intended for the European population as a whole, on the contribution of different foods or categories of foods to an overall diet that would help to maintain good health through optimal nutrition (food-based dietary guidelines).

ASSESSMENT

A draft of this Opinion, agreed by the NDA Panel on 11 April 2008, was published on the EFSA website⁶ for public consultation between 8 August and 15 December 2008. The draft Opinion was also discussed at a National Expert Meeting with Member States on Dietary Reference Values held in Barcelona on 7 and 8 September 2009. All the public comments received and comments from Member States that related to the remit of EFSA were assessed and the Opinion has been revised taking relevant comments into consideration. The comments received, a report on the outcome of the public consultation, and the minutes of the meeting with Member States have been published on the EFSA website.

1. Introduction

Acknowledging the magnitude of the task involved for this request and considering the resource implication and time constraints, the Panel decided to lay down the general principles for deriving and applying dietary reference values (DRVs) and then, for each nutrient/dietary component:

- to review existing DRVs and the criteria used in their derivation, though existing DRVs are only indicative and cannot be a basis for a full scientific assessment;
- to review and summarise existing data on sources of nutrients and on their intakes in European countries;
- to review existing literature, particularly focusing on human studies that can provide useful figures, and to select appropriate studies based on their quality and their pertinence;
- to identify criteria (endpoints) and key evidence / data on which to base the DRV and, based on the above;
- to give a critical and independent judgement on the appropriate DRVs.

The main objective of DRVs is to provide the scientific bases that are one of the key elements on which nutrition recommendations are built, expressed either as nutrient recommendations or food recommendations. Nutrition recommendations are made to ensure that a diet provides energy and nutrients for optimal growth, development, function and health during the whole life. Nutrient requirements can be deduced from physiological needs and metabolic demand. There is also increasing evidence that nutrients and diet composition influence the risk of chronic diseases (WHO, 2003). In more recent editions of (national) nutrient recommendations, epidemiological and other data on disease risk have therefore been used, if available, as an additional criterion in estimating nutrient requirements, such as for calcium, vitamin D and dietary fibre, but also in establishing recommendations for safe and adequate intake ranges for macronutrients.

In this task, the Panel will focus on establishing Dietary Reference Values for energy and nutrients. The Panel assumes that, in general, for any given population group, physiological requirements will not vary across populations in Europe and therefore allow defining a common set of dietary reference values for the European population.

⁶ http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1211902045161.htm

2. General principles for deriving dietary recommendations

As nutrients are consumed as foods, nutrient requirements are translated into nutrient reference values, then into recommended nutrient intakes and finally into food-based dietary guidelines to help consumers to meet recommended intakes (FAO/WHO, 1998; WHO, 2003; Eurodiet, 2000; USDA, 2005). In setting dietary recommendations, the following five separate steps can be identified:

2.1. Estimating the physiological requirement and metabolic demand

Metabolism embraces the processes through which life is maintained, and there are on-going needs for energy and nutrients to support these processes. The rate at which a nutrient has to be provided to the body to support metabolism and maintain functions represents the physiological requirement or metabolic demand for that nutrient. The physiological requirement varies between individuals dependent upon genetic and epigenetic differences, age, sex, physiological state such as pregnancy, and also varies in the same individual in response to environmental stress such as infection, trauma, behavioural or social stressors. Metabolic demands are related to adaptive metabolic responses in the body, such as post-exercise replenishment of fuel stores and the regeneration of damaged muscle (Millward, 1998). For a group of people of similar age, sex and physiological state, the variation in the requirement for a nutrient has been presumed to approximate a normal distribution which can be characterised by its central tendency (mean), and its distribution (standard deviation). The metabolism is regulated and its adaptation, by modifications of absorption, elimination, recycling, degradation, constitution or mobilisation of body stores, allows individuals to cope with the large variations of day to day intakes, in the short and medium term. This metabolic adaptation should be taken into account, when appropriate, in setting recommendations for populations and in managing dietary assessment or planning for individuals.

2.2. Establishing the dietary requirement of nutrients

Some nutrients need to be provided in the diet, and for others metabolism can contribute to their formation and availability to the body. The amount of a nutrient which has to be provided preformed in the diet, or the amount of the precursor which has to be taken in the diet in order to meet the physiological requirement is known as the dietary requirement. The dietary requirement has been identified as that amount of a nutrient which must be consumed on a regular basis to maintain health in an otherwise healthy individual, on the assumption that the requirements for energy and all other nutrients have already been satisfied. The dietary requirement will differ from the physiological requirement depending on a range of factors, such as the efficiency or effectiveness of absorption and utilisation of the nutrient, or the extent to which the nutrient can be formed as a part of metabolic interchange, i.e. its bioavailability and respective bioefficacy. The requirement is a characteristic of an individual and the magnitude of the dietary requirement varies amongst individuals of similar age, sex and physiological state. This variation has been presumed to approximate a normal distribution which can be characterised by its central tendency (mean), and its distribution (standard deviation).

2.3. Establishing dietary reference values

A number of different terms have been used to characterise a habitual level of intake of a nutrient which will satisfy the needs of nearly all the members of a population group. For a number of purposes, it is desirable to be able to identify this level of intake. For nutrients for which an average dietary requirement can be established, this level of intake has been set as the mean plus two standard deviations of the average requirement. The assumption is that if the intake of all members of the group were at this level, the risk of an inadequate intake for any member of the group would be small (<2.5%).

Dietary Reference Values are scientific references based on health criteria, taking into account dietary requirements and health outcomes. DRVs include the Average Requirement (AR), the Lower Threshold of Intake (LTI), the Population Reference Intake (PRI), and the Reference Intake range (RI) used for macronutrients. When no average requirement can be established, the Adequate Intake (AI) can be used. They represent one of the bases for establishing nutrient recommendations and food based dietary guidelines.

2.4. Establishing nutrient goals and recommendations

Nutrient goals are typically targets for nutrient intake at population level that may be established by policy makers for public health planning and assessment. For example, a population average intake of 30 E% as total fat might be a nutrient goal for some populations. Nutrient recommendations are typically targets for nutrient intake of individuals to ensure adequate nutrition and health status. Examples of recommended intakes of nutrients for individuals in specific populations might be an intake \geq PRI for a micronutrient or an intake of \leq 10 E% for saturated fatty acids.

These goals and recommendations may not always be communicated directly to the consumer, but are rather for use by healthcare professionals and policymakers. The definition of nutrient goals and recommendations has found great practical utility in the planning of food provisions, food-based dietary guidelines and in the characterisation of the nutritional status of populations. Nutrient goals and recommendations may differ between countries depending on health needs, nutritional status and known patterns of intake of foods and nutrients in specific populations and the actual composition of available foods.

There is a general consensus that recommendations for energy should be based on average requirements for a group and should not be used for individuals without adaptation to anthropometric parameters and the actual level of physical activity. Since the AR is higher than the requirements of half of the individuals in a group, its use for these individuals may induce a positive energy balance that, even when moderate, but chronic, has a long-term impact on body weight.

2.5. Establishing food-based dietary guidelines

Food-based dietary guidelines represent the form in which advice is provided to people to assist them in selecting a diet to meet their needs for health.

Nutrients are consumed as food. A wide range of foodstuffs together comprise the diet of the individual, group or population. Advice to people about selecting foods based on their nutritional composition is not helpful without giving advice on how to select a suitable mixture of foods to meet their nutritional requirements on the basis of foods. Increasingly it is being found that particular patterns of food consumption appear to confer health benefits that cannot simply be ascribed to their containing an adequate mixture of individual nutrients, suggesting that there are unidentified factors which may contribute to the health benefit conferred.

3. Terminology and definitions

A wide range of terminologies have been used by different national Agencies. In 2007, an expert group from the United Nations University (UNU), Food and Agriculture Organization (FAO), the World Health Organization (WHO), and the United Nations Children's Fund (UNICEF) discussed the concept of nutrient based dietary standards. They proposed a harmonisation of terminology used by national and international groups, and to limit the number of nutrient intake values (King and Garza, 2007). The Panel decided to derive a set of dietary reference values adapting the terminology from an

earlier European report (SCF, 1993) to the five step process proposed for formulating nutritional advice to people.

The complete set of dietary reference values and definitions is given below:

Dietary Reference Values (DRVs): the complete set of nutrient reference values such as the adequate intake level, the lower threshold and upper intake levels.

Population Reference Intakes (PRI): the level of (nutrient) intake that is enough for virtually all healthy people in a group.

Average Requirement (AR): the level of (nutrient) intake that is enough for half of the people in a healthy group, given a normal distribution of requirement.

Lower Threshold Intake (LTI): the level of intake below which, on the basis of current knowledge, almost all individuals will be unlikely to maintain “metabolic integrity”, according to the criterion chosen for each nutrient (see section 5.1).

In addition to these values defined by the SCF (1993), the Panel will also use the following:

Adequate Intake (AI): it is the value estimated when a PRI cannot be established because an average requirement cannot be determined.

Reference Intake ranges for macronutrients (RI): the reference intake range for macronutrients, expressed as % of the daily energy intake, defined by a lower and an upper bound.

Tolerable Upper intake Level (UL): the maximum level of total chronic daily intake of a nutrient (from all sources) judged to be unlikely to pose a risk of adverse health effects to humans.

Similar terms and definitions are being used by other authorities and in other countries, although the terminology may vary (Table 1).

Table 1: Terminology used for dietary reference values used by different bodies

	Population reference intake (PRI)	Average requirement (AR)	Lower threshold intake (LTI)	Adequate intake (AI)	Reference intake range (RI)	Tolerable upper intake level (UL)
UK (DoH, 1991)	Recommended intakes (RI)	Average requirement	Lower level of intake			
SCF, 1993	Population reference intake	Average requirement	Lower threshold intake (LTI)	Adequate intake	-	-
United States (IoM 1997, IoM 1998, IoM 2002)	Recommended dietary allowance (RDA)	Estimated average requirement (EAR)	-	Adequate Intake	Acceptable macronutrient distribution ranges (AMDR)	Tolerable upper intake level
Germany, Austria, Switzerland (D-A-CH, 2008)	Empfohlene Zufuhr	-	-	Schätzwerte Richtwerte	-	-

The Netherlands (GR, 2001)	Aanbevolen dagelijkse hoeveelheid (ADH)	Gemiddelde behoefte	-	Adequate inneming	-	Aanvaardbare bovengrens
France (AFSSA, 2001)	Apport nutritionnel conseillé (ANC)	Besoin nutritionnel moyen	-	Apport nutritionnel conseillé (ANC)	Apport nutritionnel conseillé (ANC)	Limite de sécurité
Nordic countries (NNR, 2004)	Recommended intakes (RI)	Average requirement	Lower limit of intake (LI)	-	-	Upper intake level (UL)

4. Conceptual basis for derivation of dietary reference values

Three dietary reference values (AR, PRI, LTI) describe the form of a requirement distribution when average requirements can be determined (Figure 1). When the requirement cannot be determined, two other values (AI, RI) can be proposed. An UL is also established whenever possible for nutritional safety purposes (Figure 2).

4.1. Population Reference Intakes (PRI)

The PRI is derived from the average requirement (AR) of a defined group of individuals in an attempt to take into account the variation of requirements between individuals.




Figure 1: Population reference intakes (PRI) and average requirement (AR), if the requirement has a normal distribution and the inter-individual variation is known.

By convention and on the assumption that the individual requirements for a nutrient are normally distributed within a population and the inter-individual variation is known, the PRI is calculated on the basis of the AR plus twice its standard deviation (SD) (Figure 1). This will meet the requirements of 97.5% of the individuals in the population. When the distribution of the requirement among individuals is not normal, data may be transformed to normality. The magnitude of the PRI in relation to the AR depends on the estimated variation between individuals.

However, data on inter-individual variation in requirements are limited to a few nutrients. For nutrients for which the variation in requirement is unknown, a default coefficient of variation (CV) of

10% to 20% is used assuming a normal distribution (IoM: 10%; AFSSA: 15%; GR: 10-20%). Depending on the CV used the PRI is set at 1.2 (CV=10%), 1.3 (CV=15%), or 1.4 (CV=20%) times the estimated AR. In this task, the coefficient of variation will be decided on a case-by case basis.


PRI are expressed on a daily basis, but are applied to usual intakes averaged over longer periods of time.

4.2. Average Requirement (AR)

The average requirement is the level of intake of a defined group of individuals estimated to satisfy the physiological requirement or metabolic demand, as defined by the specified criterion for adequacy for that nutrient, in half of the healthy individuals in a life stage or sex group, on the assumption that the supply of other nutrients and energy is adequate. For energy, the average requirement is generally the only reference value provided, because of the very large variations in requirements due to variations in anthropometric characteristics and in the levels of physical activity.

4.3. Lower Threshold Intake (LTI)

The LTI is the lowest estimate of the requirement from the normal distribution curve, and is generally calculated on the basis of the AR minus twice its SD (Figure 1). This will meet the requirements of only 2.5% of the individuals in the population. When the distribution of the requirement among individuals is not normal, data may be transformed to normality.



From: Health and Welfare, Canada, 1983; as adapted by Netherlands Health Council, 2000

Figure 2: Relationship between individual intake and risk of adverse effects due to insufficient or excessive intake.

4.4. Adequate Intake (AI)

An AI is the average (median) daily level of intake based on observed, or experimentally determined approximations or estimates of nutrient intake, by a group (or groups) of apparently healthy people that is assumed to be adequate. The practical implication of an AI is similar to that of a PRI, i.e. to

describe the level of intake that is considered adequate for health reasons. The terminological distinction relates to the different way in which these values are derived and to the resultant difference in the ‘firmness’ of the value.

4.5. Reference Intake ranges for macronutrients (RI)

For some energy-yielding macronutrients, reference intake ranges are expressed as the proportion (percentage) of energy derived from that macronutrient. These are usually derived from data on intake in healthy populations and on data on risk of chronic disease and apply to ranges of intakes that are adequate for maintaining health and that are associated with a low risk of selected chronic diseases.

4.6. Tolerable Upper intake Level (UL)

The tolerable upper intake level (UL) is the maximum level of total chronic daily intake of a nutrient (from all sources) judged to be unlikely to pose a risk of adverse health effects to humans⁷.

5. Methods for determining dietary reference values – types of data used

A variety of data and methods can in principle be used in deriving estimates on (average) nutrient requirements and establishing nutrient-diet-health relationships. These may vary between *in vitro* studies, animal studies, human experimental nutrition studies, and epidemiological studies. Which methods and data are most appropriate depends on the criterion or criteria chosen.

5.1. Criteria for assessment of nutrient requirement

In defining nutrient requirement the selection of criteria to establish nutrient adequacy is an important step. An obvious criterion would be the risk of deficiency, i.e. the amount of a nutrient needed to prevent, or reverse clinical deficiency symptoms. This criterion has been used in earlier recommendations, as well as criteria based upon functional competence, i.e. the amount needed to maintain a critical nutrient-related function, biochemical or physiological, such as maintenance of a desirable plasma level or nutrient-dependent enzyme activity over a certain time period, in otherwise healthy people. Maintenance of cell (organ) integrity or data on body composition, nutrient body stores or pool size are other typical criteria that can be used. In the more recently published recommendations, epidemiological and clinical data on health outcome, such as disease risk in relation to the nutrient intake pattern and/or diet composition, are also taken into consideration, if available. Although clinical endpoints, e.g. data on morbidity or mortality, are considered the most relevant, surrogate markers of health outcomes can provide useful information (IoM 1997, 1998, and 2002; Diplock et al., 1999; Asp et al., 2003, 2004).

For most nutrients a hierarchy in criteria for nutrient adequacy can be established, ranging from prevention of clinical deficiency to optimisation of body stores or status, to minimising chronic disease risk.

Which criterion, or combination of criteria, is the most appropriate remains a matter of scientific judgement, taking into account all available data and weighing of the evidence, and should be decided on a case-by-case basis.

⁷ An opinion on tolerable upper intake levels for vitamins and minerals was published in February 2006 (http://www.efsa.europa.eu/en/science/nda/nda_opinions.html). In this task, therefore, only UL's for macronutrients will be established, if appropriate.

5.2. Methods for determining nutrient requirements

Experimental feeding studies in humans, such as depletion-repletion and nutrient balance studies, are classical approaches for assessing the basic (average) requirement for a nutrient. Due to the limited time of observation they may not reflect the steady state and they may underestimate the possibility of adaptation to lower intakes.

In balance studies under- or overestimation of excretory losses and/or retention are a problem due to incomplete collection of urine and faeces, and/or through the skin, depending on the level of intake. This is especially true for nutrients that are highly conserved in the body, such as iron and calcium. Balance studies indicated higher iron retentions than observed in the more reliable radio isotope studies (Green, 1968). In the case of calcium, apparent nutrient balance has been observed, even at relatively high intakes (around 2 g/day), and this approach likely overestimates actual requirements (Heaney, 1977; Matkovic, 1990).

For some B-vitamins, metabolic conversion (or synthesis) by the intestinal bacteria might complicate balance studies. Results from depletion-repletion studies tend to overestimate requirements, most likely due to incomplete equilibration, as was shown for vitamin B₆ (see IoM report, 1998).

Biochemical markers of the nutritional (nutrient) status, or combinations of different markers, are widely used to assess nutritional adequacy. However, there is a lack of suitable biomarkers of status for some micronutrients.

Data on the dose response between nutrient intake and status indicators, such as the nutrient (or metabolite) blood or tissue level, enzyme activity, are useful, but also data on physiological (functional) response measures, such as immune competence, nerve conduction velocity, or physical or cognitive performance, are potentially valuable but are difficult to standardise. In some nutrients, the intake-status relationship may be confounded by uncertainties with respect to bioavailability.

When biochemical markers of nutritional (nutrient) status are not available and/or there are no good intake-status relationships, the so-called factorial method is used in establishing nutrient needs, which involves adding up the various factors that determine the requirement for maintenance of a defined plasma level or body store, and that is associated with the absence of adverse health effects, respectively normal tissue or body function. This requires measuring of the amounts of nutrients that leave the body via the faeces, urine and skin, either unchanged or as metabolites and also estimating the amounts that are required for growth, pregnancy or lactation. State of the art isotope techniques now enable accurate estimation of nutrient kinetics, i.e. nutrient fluxes and measurement of “true” absorption and retention rates.

The extra requirement during growth and pregnancy is due to the amounts of nutrients that become incorporated into newly formed tissues, an effect which is referred to as ‘retention’.

During pregnancy body weight increases by 10 to 15 kg, with a stage-specific rate, which is low in the first half, highest in the second trimester and slightly lower in the third trimester (0.15 to 0.69 kg/wk during wks 13 to 20; 0.31 to 0.65 kg/wk during wks 20 to 30, and 0.18 to 0.61 kg/wk during wks 30 to 36, respectively) (IoM, 1990). The total weight gain includes the weight of the foetus (appr. 25%), the placenta (appr. 5%) and the amniotic fluid (6%). In setting reference values for growing children and pregnant women consideration is also given to the many physiological adaptations that occur during these conditions, such as an increased absorption, and/or greater nutrient retention or losses.

Extra requirements during lactation are determined in part by the amounts of nutrients lost from the body in the form of human milk.

For infants during the first half year of life (0 to 6 months) the nutrient intake of healthy well-growing exclusively (fully) breast-fed infants is generally considered to be adequate and, therefore, to correspond to the requirement for nutrients. The average measured concentration of the nutrient in milk obtained from a representative sample of adequately nourished women is multiplied by the average intake volume of breast-fed infants over the first six months of life and regarded to be the AI over this period.

Exceptions to this rule are vitamin D and K, the intake of which is generally insufficient with human milk feeding. It is, moreover, important to ascertain the adequacy of the nutritional status of the lactating woman providing milk samples, especially with respect to those nutrients whose contents in milk depend on the dietary intake of the mother (i.e. iodine, selenium, vitamin A, vitamin B₆, vitamin B₁₂).

For older infants between 7 and 11 months a combination of the nutrient intake from human milk plus the intake from typical complementary food can be used, if no other data are available.

Basic research in animal models can produce valuable knowledge on mechanisms and/or dose-response relationship. However, due to inter-species differences, extrapolation from animal models to humans is subject to considerable uncertainties and data from animal models are rarely used in the setting of DRVs, e.g. in case of ULs, when data from human studies are insufficient.

5.3. Diet and health relationship: risk of chronic disease

In addition to data on basic nutrient requirements based on maintenance of body stores and function, evidence is accumulating on the relationship between dietary intake and risk of chronic disease and obesity (WHO, 2003). For some nutrients such as sodium, calcium and vitamin D, but especially for the energy-yielding macronutrients such as fatty acids, data from human studies, including both observational and intervention studies, are now used in deriving nutrient reference values and nutrient or food recommendations.

Observational studies can provide valuable data on the association between dietary nutrient intakes and health effects, such as disease risk or mortality, in free-living subjects, but by themselves cannot prove causal relationships. Additional pieces of evidence are therefore needed to establish causality. Other limitations include confounding and systematic bias in nutrient and energy intake estimates from self-reporting.

Data from randomised, controlled intervention studies with “hard” clinical endpoints, such as morbidity and mortality, are considered as the most reliable, and to provide the highest strength of evidence.

However, also intervention trials have their limitations. How much the data is representative of, and to which extent can be extrapolated to, the general population can be a problem, as intervention studies are often conducted in selected subgroups, such as individuals with an already increased risk of disease or history of the disease. Follow-up periods are relatively short compared to the long periods of disease development, especially in chronic diseases, such as cancers. Intervention with relatively high dosages and using isolated or purified nutrients will also affect the outcome, and represent a different situation as in studies with complete diets and nutrients consumed as food at usual levels of intake.

Judgement should therefore be based on the consistency, strength and quality of the studies, and take into account all the available evidence obtained with the various methods, including the knowledge on the mechanism linking nutrient intake and the occurrence of chronic disease which is often obtained from animal experiments.

5.4. Interpolation or extrapolation between population groups

Interpolation or extrapolation will be used in instances where no data are available for defined age and sex groups which permit proceeding according to sections 4.1 to 4.3. Interpolation between different population groups using a predefined hypothesis, such as isometric (linear with body weight) or allometric (body weight to the power of a chosen exponent) scaling, has limitations and will be made according to scientific judgement on a case-by-case basis (see Appendix 1).

Because of the assumptions made in interpolation or extrapolation, due to lack of knowledge on the proportionality of nutrient requirements with parameters such as body or metabolic weight during growth, reference values for the age groups between 6 months and 18 years, might have a higher uncertainty than those for adults. Metabolic weight has been defined as 0.75 power of body mass (weight), and is also used as the preferred method by some authorities to adjust for metabolic differences between age groups (see Appendix 1).

5.5. Factors that affect nutrient requirements

Nutrient requirements vary between individuals due to differences in absorption and utilisation of nutrients from the diet. This may be related to the food matrix, i.e. liberation of nutrients from the food matrix in the course of the digestive process, as well as to diet composition and interactions with other diet-related factors.

Bioavailability has been defined as the fraction of an ingested nutrient which is absorbed from the gastro-intestinal tract and utilised for normal physiological function or storage (Jackson, 1997). Host-related factors may play a role, such as stomach acidity, and in some cases nutrient status, but food/diet composition and food preparation are the main determinants of nutrient bioavailability. Iron in the form of haem-iron is more available than non-haem iron due to a different absorption mechanism. The absorption of iron also depends on the size of the body iron store.

Nutrient-nutrient interactions may result from competition at the level of absorption, such as between copper and zinc. Vitamin C enhances the absorption of non-haem iron through formation of a soluble complex with ferrous iron in the stomach.

Dietary protein requirements depend on the protein quality of the diet, and present another example of a food-related factor affecting dietary requirements (Millward and Jackson, 2004). Dietary protein requirements may be slightly higher for groups consuming mainly vegetable proteins, e.g. vegans.

5.6. Lifestage and sex

In this task, the reference weights for different age and sex groups from the SCF reports (1993 and 2000) will be considered as an initial proposal (Table 2), until more recent data become available. Reference weights may be used in interpolation, or extrapolation of nutrient requirements for population (sub)groups for which insufficient or no data are available to set a reference value. Reference weights are also used in the calculation of the average energy requirements. The Panel recognises that the reference weights used in the SCF report were based upon pooling of national data from a limited number of EU Member States. These data are relatively old and not necessarily representative for the newer EU Member States, but will be used unless more recent and representative data becomes available. The Panel would recommend the development of a database with reference weights and heights representative for the total European population.

The age group <6 months will not be considered given that requirements will refer to amounts of nutrients provided by breast-milk, except on a case-by-case basis where this does not apply. The choice of life stage groups will be based upon differences in requirements related to velocity of

growth, change in endocrine status, such as in puberty, and age-related changes in nutrient absorption and body functions and/or functional capacity, such as renal function. In older subjects, especially in elderly above 75 years, variability in functional capacity and in energy expenditure appears higher than in younger adults. Separate recommendations will be made for pregnant and lactating women. The Panel proposes to define the age ranges used for each nutrient on a case-by-case basis depending on the available data.

Table 2: Reference weights of population groups in Europe (source: SCF, 1993 and 2000).

Age (yr)	Reference weight (kg) ¹	
	Males	Females
0-6 months	6.0	5.5
7-11 months	9.0	8.3
1-3	13.0	12.5
4-6	20.0	19.0
7-10	28.5	29.0
11-14	44.5	45.0
15-17	61.5	53.5
18-29	74.6	62.1
30-59	74.6	62.1
60-74	73.5	66.1
≥75	73.5	66.1

¹ Based upon median weights of European men and women from various studies (for references see SCF, 1993). Weights for the age groups up to 1 year are based upon median values from the WHO Multicentre Growth Reference Study (WHO, 2006a).

6. Application of dietary reference values for nutrients

Dietary Reference Values are the basis for formulating nutrient recommendations that are applied in different areas, especially diet assessment and diet planning for individuals or for population(s) (groups) but are also the basis for deriving reference values for food labelling.

Diet assessment applications involve determining the probable (in)adequacy or excess of observed intakes. Diet planning applications involve using dietary reference values to develop recommendations for what food intakes should be in order to achieve nutritional balance. Rather than focussing on a fixed recommended or advised intake level, the distribution of nutrient intakes among the population is now considered of more importance. For that purpose a set of dietary reference values is derived that can be used both for diet assessment and planning. These dietary reference values are intended for healthy populations and cannot be applied without caution to patients.

6.1. Assessment of the risk of (in)adequacy of nutrient intake

6.1.1. Assessment of the risk of (in)adequacy of nutrient intake in populations

Intakes of individuals vary from day-to-day, and thus many days of intake are needed to estimate a person's usual intake. Because it is seldom practical to collect data for many days, statistical methods are available to remove the effect of day-to-day variation from the intake distribution for a group. This general approach was proposed by NRC (1986), was further developed by Nusser et al. (1996) and shown to be robust (Hofmann et al., 2002). It can also be adapted to take into account the effects of age (Waijers et al., 2006). To adjust intake distributions it is necessary to have at least two

independent days of dietary intake data for a representative subsample of individuals in the group (IoM, 2000).

Most dietary assessment methods result in some underreporting of food (and nutrient) intakes at the group level, and the methods of removing day-to-day variation do not remove this negative effect. Therefore, the prevalence of inadequacy in populations may be overestimated for some nutrients (Murphy et al., 2006).

The proportion of the population with an inadequate intake may be estimated using the AR cut-point method, originally proposed by Beaton (1994) and adopted by the US Food and Nutrition Board (IoM, 2000). This method requires knowledge of the AR and the distribution of habitual nutrient intakes and has been shown to be effective in obtaining a realistic estimate of the prevalence of dietary inadequacy (Carriquiry, 1999). The percentage of the population with a habitual daily nutrient intake that is lower than the AR is taken as an estimate of the percentage of the population with probable inadequate intakes. For example, at a median intake equal to the AR, 50% of a population group will have intakes that may be inadequate for the chosen criterion of nutritional status. At a median intake level around the PRI, intakes are considered adequate for 97.5% of the population group. Higher intakes convey no additional health benefit. For nutrients for which a UL has been established intakes above the UL may be associated with increased prevalence of adverse effects in the population.

The AR cut-point method only holds if the distribution of nutrient intakes is normal and requirements and intakes are independent. It is generally assumed that the average individual daily intakes of vitamins and minerals are not related to requirements, and that the average requirements for vitamins and minerals, except iron, are symmetrically distributed (SCF, 1993; IoM, 1997). However, in the case of skewed distributions, such as in the case of iron, the cut-point method would underestimate the prevalence of inadequacy in menstruating women (SCF, 1993; IoM, 2000). The SD of the habitual daily intakes of vitamins and minerals are generally greater than 30% of the mean (O'Brien et al., 2001; Hannon et al., 2001) and are almost always more than twice the commonly assumed variance of requirement of 10 to 15% of the mean.

Energy and protein maintenance requirements are correlated with energy expenditure and body protein mass, respectively, and energy and protein intakes, together with energy and protein requirements may therefore behave collinear. Actually, this relationship may involve other nutrients, because their intakes are often correlated (Day et al., 2001 and 2004).

In these cases, the AR cut-point method may overestimate the risk of inadequacy.

Probabilistic methods, which take into account both the intake and requirement variability might be a useful alternative for the AR cut-point method, and give a better estimation of the real prevalence of inadequacy (de Lauzon et al., 2004).

The PRI is an intake level that covers the requirement of 97-98% of all individuals when requirements of the group have a normal distribution. The PRI should therefore not be used as a cut-point for assessing nutrient intakes of groups because a certain overestimation of the proportion of the group at risk of inadequacy would result.

The prevalence of inadequacy of intake of a nutrient in a population cannot be estimated using as comparator the less precise estimates of recommended intake, e.g. nutrients with “adequate intake” (IoM, 1997), ‘acceptable range of intakes’ (SCF, 1993) or “safe and adequate intake” (DoH, 1991), because the relationship of such reference values to the requirement for the nutrient is not known. Groups with mean intakes at or above the AI can be assumed to have a low prevalence of inadequate intakes for the defined criterion of nutritional status.

For macronutrients which have a reference intake range for individuals, the distribution of usual intakes of individuals may be assessed to ascertain what proportion of the group lies outside the reference lower and upper bounds of the range.

Reference intakes of energy for groups are based on estimated average energy expenditure associated with the sex, age, height, weight, and physical activity level of the group. Mean usual intake of energy relative to the average requirement may be used in assessing the adequacy of energy intakes of groups. Because there is a high correlation between energy intake and energy expenditure (requirement), median intake of food energy should be close to the requirement for there to be low risk of inadequate or excessive intake. However, because of an expected correlation between energy intakes and energy needs at the group level, it is not possible to generate an unbiased estimate of the prevalence of inadequate or excessive intakes.

6.1.2. Assessment of the risk of (in)adequacy of nutrient intake in individuals

Usual nutrient intakes of individuals may be compared with specific Dietary Reference Values, even though dietary intake data alone cannot be used to ascertain an individual's nutritional status. However, such comparisons are of limited use because of inherent problems in the validity of the assessment of usual dietary intake in individuals. Ideally, usual intake data should be combined with anthropometric, biochemical (status), and clinical information to provide a valid assessment of an individual's nutritional status.

For nutrients with an AI (i.e. without an AR), if an individual's usual intake equals or exceeds the AI, it can be concluded that the diet is almost certainly adequate. If, however, an individual's intake falls below the AI, no quantitative (or qualitative) estimate can be made of the probability of nutrient inadequacy (IoM, 2000).

Observed intakes below the LTI have a very high probability of inadequacy.

Observed intakes of an individual below the AR very likely are inadequate because the probability of inadequacy is up to 50 percent, and an intake between the AR and the PRI may be adequate because the probability of adequacy is higher than 50 percent. For nutrients for which a UL has been established chronic intakes above the UL may be associated with an increased risk of adverse effects, and should therefore be avoided.

For macronutrients which have a reference intake range, the usual intake of individuals may be assessed to ascertain whether the intake lies between the reference range lower and upper bounds.

Reference intakes of energy for individuals are based on estimated average energy expenditure associated with an individual's sex, age, height, weight, and physical activity level. As such, it exceeds the needs of half the individuals with specified characteristics, and is below the needs of the other half. Because it is difficult to determine energy balance, even from several days of intake, recent weight history is used as an indicator of the likely adequacy of energy intake.

6.2. Dietary planning

6.2.1. Dietary planning for groups

Planning diets for groups may include institutional food planning, military food and nutrition planning, planning for food assistance programs, food fortification, and assuring food safety.

For groups, the goal of planning is to determine a usual intake distribution that results in a low prevalence of intakes that are inadequate (i.e. less than the AR) or at risk of being excessive (i.e. greater than the UL). For nutrients such as vitamins, minerals, and protein, the PRI could be a practical starting point. However, target median intakes higher than the PRI might be considered, especially in case of a skewed intake distribution (WHO, 2006b). The AI may be used as an appropriate target average when no AR has been established. For macronutrients which have a reference intake range, the distribution of usual intakes of individuals should be such as to minimise the proportion of the group whose intake lies outside the lower and upper bounds of the reference range. For energy, the reference intake (average energy requirement) of the group based on sex, age, height, weight, and physical activity level of the group may be used as a planning goal (IoM, 2005).

6.2.2. Dietary planning for individuals

Planning diets for individuals may include: (1) providing guidance to healthy individuals to assist them in meeting their nutrient needs, (2) counseling those with special lifestyle considerations (e.g., athletes and vegetarians) or those requiring therapeutic diets, (3) formulating diets for research purposes, (4) developing food-based dietary guidance for individuals. In addition, providing consumer information on food and food supplement labels can assist consumers in planning their own diets.

The goal of planning diets for individuals is to have a low probability of inadequacy while minimising a potential risk of excess for each nutrient. For nutrients such as vitamins, minerals, and protein, this is done by ensuring that the usual intake meets the PRI or AI while not exceeding the UL. PRIs would be an overestimation for most individuals. For macronutrients which have a reference intake range, the usual intake of individuals should be between the lower and upper bounds of the reference range. For energy, the reference intake (average energy requirement) based on an individual's sex, age, height, weight, and physical activity level may be used as an initial planning goal; however, body weight must be monitored and intake adjusted as appropriate (IoM, 2005).

6.3. Labelling reference intake values

Dietary reference values are used for nutrition information on food and food supplements, as labelling reference intake values so that consumers can compare products and see how a food fits into their overall daily diet. In the SCF opinion on the revision of reference values for nutrition labelling (SCF, 2003), the PRI was suggested as a basis for Labelling Reference Intake (LRI) values. An 'overall' PRI was extracted, based upon PRI (RDA) values for adults taken from a number of European countries, as well as from values from the United States and FAO/WHO.

Also the population-weighted mean of the AR has been suggested for use in nutrition labelling as this lower value might be a better statistical approximation to the nutrient requirement for any one individual in the population (IoM; 2005; Tarasuk, 2006; Yates, 2006). Accepting the lower AR values or the higher PRI value as a basis for LRI will affect the use of nutrient content claims on foods that are linked to LRI, and therefore the number of foods that could claim being a 'source' of or 'high' in a certain nutrient.

However, the substantial differences in the LRI derived by taking the AR as the basic value, compared to the existing LRI, which are closer to existing RDA/PRI, might be confusing for the consumer, and for that reason the SCF Committee accepted the RDA/PRI as the basis for the LRI. Which value to take is not a scientific decision, but a management decision to be taken after careful consideration of all implications (EFSA, 2009).

6.4. Food-based dietary guidelines

As diets are composed of foods, food-based dietary guidelines (FBDG) are the form in which advice is provided to the consumer. FBDG are generally based upon scientific evidence on the relationship between diet and chronic disease risk, taking into account nutrient recommendations (WHO 2003). The FBDG are intended for use by individual members of the general public, and as a tool for use by policymakers, nutritionists, nutrition educators and healthcare providers, to help consumers in planning an overall healthy diet, while achieving an adequate nutrient intake (EFSA, 2009).

CONCLUSIONS

This Opinion focuses on the general principles for development and application of Dietary Reference Values (DRVs) - quantitative reference values on which to base nutrient and food recommendations for healthy individuals and populations which may be used for assessment and planning of diets.

Similarly to the earlier Scientific Committee on Food (SCF) report in 1993, the Panel proposes to derive the following Dietary Reference Values:

- *Population Reference Intakes (PRI)*: the level of (nutrient) intake that is adequate for virtually all people in a population group.
- *Average Requirement (AR)*: the level of (nutrient) intake that is adequate for half of the people in a population group, given a normal distribution of requirement.
- *Lower Threshold Intake (LTI)*: the level of intake below which, on the basis of current knowledge, almost all individuals will be unable to maintain “metabolic integrity”, according to the criterion chosen for each nutrient.

In addition, the Panel also proposes to derive the following Dietary Reference Values:

- *Adequate Intake (AI)*: the value estimated when a Population Reference Intake cannot be established because an average requirement cannot be determined. An Adequate Intake is the average observed daily level of intake by a population group (or groups) of apparently healthy people that is assumed to be adequate.
- *Reference Intake ranges for macronutrients (RI)*: the reference intake ranges for macronutrients, expressed as % of the energy intake. These apply to ranges of intakes that are adequate for maintaining health and associated with a low risk of selected chronic diseases.

The Panel will not address the Tolerable Upper Intake Level (UL) as this has been assessed previously. The Tolerable Upper Intake Level is the maximum level of total chronic daily intake of a nutrient (from all sources) judged to be unlikely to pose a risk of adverse health effects to humans.⁸

Some of the Dietary Reference Values - the Average Requirement, Population Reference Intake and Lower Threshold Intake - relate to nutrient requirements that are defined by specific criteria of nutrient adequacy. In defining nutrient requirements the selection of criteria to establish nutrient adequacy is an important step. For most nutrients a hierarchy of criteria for nutrient adequacy can be established, ranging from prevention of clinical deficiency to optimisation of body stores, or status. Which criterion, or combination of criteria, will be the most appropriate will be decided on a case-by-case basis.

⁸ An opinion on tolerable upper intake levels for vitamins and minerals was published in February 2006 (http://www.efsa.europa.eu/en/science/nda/nda_opinions.html).

Because of this, Dietary Reference Values are developed for different life stage and sex groups. The Panel proposes to define the age ranges used for each nutrient on a case-by-case basis depending on the available data. For the age group <6 months requirements are considered to be equal to the supply from breast-milk, except on a case-by-case basis where this does not apply. Separate reference values will be established for pregnant and lactating women, taking into account the additional nutrient requirement for the formation of new tissues, or to compensate for the nutrients lost to the body in the form of human milk, respectively, and considering the physiological adaptations that occur during these conditions.

Interpolation or extrapolation between population groups will be used in instances where no data are available for defined age and sex groups. Scaling methods using isometric (linear with body weight) or allometric (body weight to the power of a chosen exponent) or interpolation based on other non predefined parameters are being used. Which method is the most appropriate will be decided on a case-by-case basis.

Reference heights and weights are useful when more specificity about body size and nutrient requirements are needed than that provided by life stage categories. In the absence of more recent data, reference weights will be the same as in the SCF report, and for children <1 year, as established by the WHO for fully breastfed infants.

Dietary reference values can be used for different purposes, such as diet assessment and diet planning, both at the population and individual level, but also as a basis for reference values in food labelling, and in establishing food based dietary guidelines.

In dietary assessment of groups the Average Requirement can be used to estimate the prevalence of the risk of inadequate intakes of micronutrients (the Average Requirement cut-point method), if the distribution of nutrient intakes is normal, and intakes are independent from requirements. The Population Reference Intake should not be used for this purpose as this would result in overestimation of the proportion of the group at risk of inadequacy.

For macronutrients with a defined reference intake range, the distribution of usual intakes of individuals may be assessed to ascertain what proportion of the group lies outside the lower and upper limits of the reference range. In case of energy, the mean usual intake of energy of a defined group, relative to the average requirement, may be used in assessing the adequacy.

For assessment of adequacy of nutrient intakes in individuals dietary reference values are of limited use. Usual intakes below the Average Requirement are likely inadequate, and below the Lower Threshold Intake very probably inadequate, while chronic intakes above the Tolerable Upper Intake Level may be associated with an increased risk of adverse effects. For a valid assessment of the adequacy of an individual's usual intake, combined information with anthropometric, biochemical (status) and clinical data is needed.

In dietary planning for groups the usual intake distribution should be between the Average Requirement and Tolerable Upper Intake Level to avoid inadequate, respectively excessive intakes. For nutrients such as vitamins, minerals, and protein, the Population Reference Intake can be a practical starting point. However, target median intakes higher than the Population Reference Intake might be considered, especially in case of a skewed intake distribution. For macronutrients the distribution of usual intakes of individuals should be such as to minimise the proportion of the group that lies outside the lower and upper bounds of the reference range. For energy, the reference intake (estimated average energy requirement) of the group based on sex, age, height, weight, and physical activity level of the group may be used as a planning goal.

The goal of planning diets for individuals is to have a low probability of inadequacy while minimising a potential risk of excess for each nutrient. For nutrients such as vitamins, minerals, and protein, this is done by ensuring that the usual intake meets the Population Reference Intake or Adequate Intake while not exceeding the Tolerable Upper Intake Level. Population Reference Intakes would be an overestimation for most individuals. For macronutrients which have a Reference Intake range, the usual intake of individuals should be between the lower and upper bounds of the reference range. For energy, the reference intake (average energy requirement) based on an individual's sex, age, height, weight, and physical activity level may be used as an initial planning goal; however, body weight must be monitored and intake adjusted as appropriate.

REFERENCES

- AFSSA (Agence Française de Sécurité Sanitaire des Aliments), 2001. Apports nutritionnels conseillés pour la population française. Editions Tec&Doc, Paris, 111.
- Agutter PS and Wheatley DN, 2004. Metabolic scaling: consensus or controversy? *Theoretical Biology and Medical Modelling*, 1, 13.
- Asp N-G, Cummings JH, Howlett J, Rafter J, Riccardi G and Westenhoefer J, 2004. Process for the assessment of scientific support for claims on foods (PASSCLAIM) Phase Two: Moving Forward. *European Journal of Nutrition*, 43, Suppl. 2, 1-133.
- Asp N-G, Cummings JH, Mensink RP, Prentice A, Richardson DP and Saris WHM, 2003. Process for the assessment of scientific support for claims on foods (PASSCLAIM) Phase One: Preparing the Way. *European Journal of Nutrition*, 42, Suppl. 1 1-113.
- Beaton G, 1994. Criteria of an adequate diet. In: *Modern Nutrition in Health and Disease*. Eds Shils ME, Olson JA, Shike M. Lea & Febiger, Philadelphia, 1491-1505.
- Carriquiry AL, 1999. Assessing the prevalence of nutrient inadequacy. *Public Health Nutrition*, 2, 23-33.
- D-A-CH (Deutsche Gesellschaft für Ernährung - Österreichische Gesellschaft für Ernährung - Schweizerische Gesellschaft für Ernährungsforschung - Schweizerische Vereinigung für Ernährung), 2008. Referenzwerte für die Nährstoffzufuhr. Umschau Braus Verlag, Frankfurt am Main.
- Day NE, McKeown N, Wong MY, Welch A and Bingham S, 2001. Epidemiological assessment of diet: a comparison of a 7-day diary with a food frequency questionnaire using urinary markers of nitrogen, potassium and sodium. *International Journal of Epidemiology*, 30, 309-317.
- Day NE, Wong MY, Bingham S, Khaw KT, Luben R, Michels KB, Welch A and Wareham NJ, 2004. Correlated measurement error – implications for nutritional epidemiology. *International Journal of Epidemiology*, 33, 1373-1381.
- de Lauzon B, Volatier JL and Martin A, 2004. A Monte Carlo simulation to validate the EAR cut-point method for assessing the prevalence of nutrient inadequacy at the population level. *Public Health Nutrition*, 7, 893-900.
- Diplock A, Aggett P, Ashwell M, Bornet F, Fern E and Roberfroid M, 1999. Scientific Concepts of Functional Foods in Europe Consensus Document. *British Journal of Nutrition*, 81, S1-S27.
- DoH (Department of Health), 1991. Dietary Reference Values for Food Energy and Nutrients for the United Kingdom. Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy. HMSO, London.

- EFSA (European Food Safety Authority), 2009. Scientific Opinion of the Panel on Dietetic products, Nutrition and Allergies on a request from European Commission on the review of labelling reference intake values for selected nutritional elements. *The EFSA Journal*, 1008, 1-14.
- Eurodiet, 2000. Eurodiet core report: Nutrition and diet for healthy lifestyles in Europe. Science and policy implications. Available from:
http://ec.europa.eu/health/ph_determinants/life_style/nutrition/report01_en.pdf
- Feldman HA and McMahon TA, 1983. The 3/4 mass exponent for energy metabolism is not a statistical artifact. *Respiration Physiology*, 52, 149-163.
- GR (Gezondheidsraad), 2001. Dietary Reference Intakes: energy, proteins, fats and digestible carbohydrates. Publication no. 2001/19R, Health Council of the Netherlands, The Hague.
- Green R, Charlton R, Seftel H, Bothwell T, Mayet F, Adams B, Finch C and Layrisse M, 1968. Body iron excretion in man: a collaborative study. *American Journal of Medicine*, 45, 336-353.
- Hannon EM, Kiely M, Harrington KE, Robson PJ, Strain JJ and Flynn A, 2001. The North/South Ireland Food Consumption Survey: mineral intakes in 18-64-year-old adults. *Public Health Nutrition*, 4, 1081-1088.
- Health Council of the Netherlands, 2001. Dietary Reference Intakes: energy, proteins, fats and digestible carbohydrates. The Hague: Health Council of the Netherlands, 2006; publication no. 2001/19, 110 pp.
- Heaney RP, Recker RR and Saville PD, 1977. Calcium balance and calcium requirements in middle-aged women. *American Journal of Clinical Nutrition*, 30, 1603-1611.
- Hoffmann K, Boeing H, Dufour A, Volatier JL, Telman J, Virtanen M, Becker W and De Henauw S, 2002. Estimating the distribution of usual dietary intake by short-term measurements. *European Journal of Clinical Nutrition*, 56 Suppl 2, S53-62.
- IoM (Institute of Medicine), 1990. Nutrition during pregnancy. National Academies Press, Washington DC.
- IoM (Institute of Medicine), 1997. Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride. National Academies Press, Washington DC.
- IoM (Institute of Medicine), 1998. Dietary Reference Intakes for thiamin, riboflavin, niacin, vitamin B6, folate, vitamin B12, pantothenic acid, biotin and choline. National Academies Press, Washington DC.
- IoM (Institute of Medicine), 2000. Dietary Reference Intakes: Applications in Dietary Assessment. National Academies Press, Washington D.C.
- IoM (Institute of Medicine), 2002. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty acids, Cholesterol, Protein, and Amino Acids. National Academies Press, Washington D.C.
- IoM (Institute of Medicine), 2003a. Dietary Reference Intakes: Applications in Dietary Planning. National Academies Press, Washington D.C.
- IoM (Institute of Medicine), 2003b. Dietary Reference Intakes: Guiding principles for nutrition labelling and fortification. National Academies Press, Washington D.C.
- Jackson MJ, 1997. The assessment of bioavailability of micronutrients: introduction. *European Journal of Clinical Nutrition*, 51 Suppl 1, S1-2.
- King JC and Garza C, 2007. Harmonization of nutrient intake values. *Food Nutr Bull*, 28, S3-12.
- Kleiber M, 1932. Body size and metabolism. *Hilgardia*, 6, 315-351.
- Kleiber M, 1947. Body size and metabolic rate. *Physiological Reviews*, 27, 511-541.

- Matkovic V, Fontana D, Tominac C, Goel P and Chesnut CH, 3rd, 1990. Factors that influence peak bone mass formation: a study of calcium balance and the inheritance of bone mass in adolescent females. *American Journal of Clinical Nutrition*, 52, 878-888.
- Millward DJ, 1998. Metabolic demands for amino acids and the human dietary requirement: Millward and rRvers (1988) revisited. *Journal of Nutrition*, 128, 2563S-2576S.
- Millward DJ and Jackson AA, 2004. Protein/energy ratios of current diets in developed and developing countries compared with a safe protein/energy ratio: implications for recommended protein and amino acid intakes. *Public Health Nutrition*, 7, 387-405.
- Murphy SP, Guenther PM and Kretsch MJ, 2006. Using the dietary reference intakes to assess intakes of groups: pitfalls to avoid. *Journal of the American Dietetic Association*, 106, 1550-1553.
- NNR (Nordic Nutrition Recommendations), 2004. Integrating nutrition and physical activity. Nordic Council of Ministers, Copenhagen.
- NRC (National research Council), 1986. Nutrient Adequacy. Assessment Using Food Consumption Surveys. National Academic Press, Washington DC.
- Nusser SM, Carriquiry AL, Dodd KW and Fuller WA, 1996. A Semiparametric Transformation Approach to Estimating Usual Daily Intake Distributions. *Journal of the American Statistical Association*, 91, 1440-1449.
- O'Brien M, Kiely M, Harrington K, Robson P, Strain J and Flynn A, 2001. The North/South Ireland Food Consumption Survey: vitamin intakes in 18-64-year-old adults. *Public Health Nutrition*, 4, 1069-1080.
- Rubner M, 1883. Über den Einfluss der Körpergröße auf Stoff- und Kraftwechsel. *Zschr Biol*, 19, 535-562.
- SCF (Scientific Committee on Food), 1993. Report on nutrient and energy intakes for the European Community.
- SCF (Scientific Committee on Food), 2000. Opinion on guidelines for the development of tolerable upper intake levels for vitamins and minerals.
- SCF (Scientific Committee on Food), 2003. Opinion on the revision of reference values for nutrition labelling.
- Tarasuk V, 2006. Use of population-weighted Estimated Average Requirements as a basis for Daily Values on food labels. *American Journal of Clinical Nutrition*, 83, 1217S-1222S.
- USDA (U.S. Dept. of Agriculture) and HHS (U.S. Dept. of Health and Human Services), 2005. Nutrition and your health: dietary guidelines for Americans. U.S. Dept. of Agriculture, U.S. Dept. of Health and Human Services, Washington D.C.
- Waijers PM, Dekkers AL, Boer JM, Boshuizen HC and van Rossum CT, 2006. the potential of AGE MODE, an age-dependent model, to estimate usual intakes and prevalences of inadequate intakes in a population. *Journal of Nutrition*, 136, 2916-2920.
- West GB and Brown JH, 2005. The origin of allometric scaling laws in biology from genomes to ecosystems: towards a quantitative unifying theory of biological structure and organization. *Journal of Experimental Biology*, 208, 1575-1592.
- White CR and Seymour RS, 2003. Mammalian basal metabolic rate is proportional to body mass^{2/3}. *Proceedings of the National Academy of Sciences of the United States of America*, 100, 4046-4049.
- White CR and Seymour RS, 2005. Allometric scaling of mammalian metabolism. *Journal of Experimental Biology*, 208, 1611-1619.

- WHO (World Health Organization), 1998. Preparation and use of food-based dietary guidelines. Report of a joint FAO/WHO consultation. WHO technical report series n 880, WHO, Geneva.
- WHO (World Health Organization), 2006a. Multicentre Growth Reference Study Group. WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatrica*, Suppl. 450, 76-85.
- WHO (World Health Organization), 2006b. Guidelines on Food Fortification with Micronutrients (Allen L., de Benoist B., Dary O., Hurrell R. Eds).
- WHO/FAO (World Health Organization and Food and Agriculture Organization), 2003. Expert Report: Diet, nutrition and prevention of chronic diseases. Report of a Joint WHO/FAO Expert Consultation. WHO Technical Report Series 916.
- Yates AA, 2006. Which dietary reference intake is best suited to serve as the basis for nutrition labeling for daily values? *Journal of Nutrition*, 136, 2457-2462.

APPENDICES

APPENDIX 1 INTERPOLATION AND EXTRAPOLATION

Interpolation will be used in instances where no data are available for defined age and sex groups which permit proceeding according to sections 4.1 to 4.3. Interpolation between different population groups is the least reliable method to determine the nutrient requirement. Extrapolation from the known AR or AI of one population group to other groups for which no specific data exist can be performed based on differences in typical body weights or on differences in energy requirement. Extrapolation should be performed sex-specific if differences in nutrient requirement according to sex are known. The typical body weights for age groups are listed in Table 2 of section 5.6.

Depending on whether the requirement for the nutrient under consideration is associated with the metabolic rate or is not associated with the metabolic rate allometric or isometric scaling is performed.

In *isometric scaling* the AI or AR for the population group X under consideration is derived by multiplication of the known and sex-specific AI or AR of group Y with the quotient between the typical weight of group X and the weight of group Y:

[Formula 1]: $AI \text{ (or AR)}_X = AI \text{ (or AR)}_Y \times (\text{body weight}_X / \text{body weight}_Y)$.

Isometric scaling has, for example, been used for magnesium and fluoride.

Allometric scaling reflects that the metabolic rate of an organism is an exponential function of body mass (weight). The original finding that the metabolic rate of humans and animals was linearly related to body surface area (Rubner, 1883) and therefore, roughly to body weight^{2/3} was modified by Kleiber (1932 and 1947), who demonstrated that the logarithm of the basal metabolic rate of ten different mammalian species was linearly related to the logarithm of the body weight with a slope of 0.75. He coined the term metabolic body size for body weight^{0.75} and predicted that the requirement for nutrients should be proportional to the metabolic body weight if their rate of “destruction” or excretion was found to be proportional to metabolic rate. However, this rule only applied to mature organisms at indifferent environmental temperature and at rest. Although this rule has never directly been proven to be correct with respect to nutrient requirements and although the discussion whether the mass exponent between different species is the same as within a species and whether it is nearer 0.67 or nearer 0.75 has not yet been conclusively resolved (Feldman and McMahon, 1983; Agutter and Wheatley, 2004; White and Seymour, 2003 and 2005; West and Brown, 2005), scaling as the 0.75 power of body mass (weight) has been widely accepted in nutritional science. Compared to isometric scaling, allometric scaling from a higher to a lower body weight gives higher values, whereas allometric scaling from a lower to a higher body weight results in lower values.

[Formula 2]: $AI \text{ (or AR)}_X = AI \text{ (or AR)}_Y \times (\text{weight}_X / \text{weight}_Y)^{0.75}$

When scaling down from an adult AI or AR to children corrections for growth requirements have to be made in order to account for the nutrient amount required to be incorporated into newly formed tissue. One way to do this is to add an age specific growth factor based on the proportional increase in protein requirements for growth (as calculated from Tables 33 and 34 in the FAO/WHO/UNU report, 1985) to both the formula 1 and 2:

[Formula 3]: $AI \text{ (or AR)}_X = AI \text{ (or AR)}_Y \times (\text{weight}_X / \text{weight}_Y) \times (1 + \text{growth factor})$

[Formula 4]: $AI \text{ (or AR)}_X = AI \text{ (or AR)}_Y \times (\text{weight}_X / \text{weight}_Y)^{0.75} \times (1 + \text{growth factor})$.

The following growth factors have been applied (IoM, 1998; FAO/WHO, 2004):

<u>Age</u>	<u>Growth factor</u>
7 months - 3 years	0.30
4 - 8 years	0.15
9 -13 years	0.15
14 - 18 years males	0.15
14 - 18 years females	0.00

Another method of interpolation has been proposed and assumes that the nutrient requirement after the age of 0 to 5 months until the age of 18 years increases linearly with age and that the requirement between the age of 14 and 18 years is the same as that of persons 19 to 50 years old. Additionally, the following age correction factors (AF) are applied:

6-11 months	0.03
1- 3 years	0.14
4-8 years	0.38
9-13 years	0.69

Adequate Intake or the Population Reference Intake are then calculated as follows:

$$AI = AI_{[0-5 \text{ months}]} + [AF \times (AI_{[>14 \text{ year}]} - AI_{[0-5 \text{ month}]})]$$

or

$$PRI = PRI_{[0-5 \text{ months}]} + [AF \times (PRI_{[>14 \text{ year}]} - PRI_{[0-5 \text{ month}]})]$$

In this formula, “AI” stands for Adequate Intake and “PRI” for Population Reference Intake (Gezondheidsraad NL, 2002).

Extrapolation was used by the French expert Committee (AFSSA, 2001) as an alternative approach to interpolation in order to calculate reference values of vitamins for children. This approach assumes that both nutritional reference values at the ends of the range are valid, namely acceptable intakes for newborns (derived from the supply by maternal milk) and Population Reference Intakes for adults. Various representative parameters for children aged 0 to 18 years, provided by the National Institute of Statistics, were used for extrapolation: weight, height, body mass index, square height (representative of lean body mass), energy intake (as a marker of metabolic rates), and body surface area were tested. For each vitamin, only one parameter appeared to allow satisfactory extrapolation in both sexes, either starting from the adult values to reach newborn values and starting from newborn values to reach adult values: square height (for Vitamins, B₆, B₁₂, thiamine, biotin, folic acid) and energy intake (for vitamins C, A, E, riboflavin, pantothenic acid, niacin). This methodology does not require any a priori judgement on what could be the best equation for the derivation of values for children.

GLOSSARY / ABBREVIATIONS

AFSSA	Agence Française de Sécurité Sanitaire des Aliments
AI	Adequate Intake
AR	Average Requirement
CV	Coefficient of variation
D-A-CH	Nutrition Recommendations for Germany, Austria and Switzerland
DoH	Department of Health (United Kingdom)
DRV	Dietary Reference Value
EC	European Commission
EFSA	European Food Safety Authority
EU	European Union
FAO	Food and Agriculture Organisation
FBDG	Food-based dietary guidelines
FNB	U.S. Food and Nutrition Board
GR	Gezondheidsraad (The Netherlands Health Council)
IoM	Institute of Medicine (United States)
LTI	Lower Threshold Intake
NNR	Nordic Nutrition Recommendations
PRI	Population Reference Intakes
RI	Reference Intake ranges for macronutrients
SCF	Scientific Committee for Food
SD	Standard Deviation
UL	Tolerable Upper Intake Level
US	United States
WHO	World Health Organization