



Heterologous expression of polyketides in fungi and optimization by an in silico guided approach

Mølgaard, Louise; Hansen, Bjarne Gram; Nielsen, Jakob Blæsbjerg; Patil, Kiran Raosaheb; Thykær, Jette; Mortensen, Uffe Hasbro

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Mølgaard, L., Hansen, B. G., Nielsen, J. B., Patil, K. R., Thykær, J., & Mortensen, U. H. (2011). *Heterologous expression of polyketides in fungi and optimization by an in silico guided approach*. Poster session presented at Conference on Synthetic Biology and Cell Factories, Vejle, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Heterologous expression of polyketides in fungi and optimization by an *in silico* guided approach




Louise Mølgaard*, Bjarne G. Hansen, Jakob B. Nielsen, Kiran R. Patil, Jette Thykær & Uffe H. Mortensen

Center for Microbial Biotechnology, Department of Systems Biology, Technical University of Denmark, Kgs. Lyngby, Denmark - *lom@bio.dtu.dk


Polyketides are the source of the most potent antibiotics and anticancer agents available today. They constitute a large and diverse group of natural compounds produced by plants, fungi and bacteria. However, the productivity is often very low in the native producer. Thus, expression of polyketide biosynthesis genes in relevant cell factories has a great potential. Towards fulfilling this goal we have chosen the two fungal model species *Saccharomyces cerevisiae* and *Aspergillus nidulans*, where numerous genetic tools are available. The overall aim of the project is to construct microbial super hosts for polyketide production through the use of state of the art genetic engineering and *in silico* modeling tools.

Project strategy




Modeling

By using a stoichiometric model of *S. cerevisiae* and the algorithm OptGene, we were able to predict several gene deletions that should improve the production of 6-methylsalicylic acid (6-MSA)¹.




Strain Construction

6-MSA production was established in *S. cerevisiae* and *A. nidulans* followed by the predicted deletion of the gene encoding glutamate dehydrogenase 1 (*GDH1* and *gdhA*, respectively). This was coupled to the insertion of the respective *GDH2* and *gdhB*^{2, 3}.




The figure illustrates the method used for achieving 6-MSA production in both *A. nidulans* (left) and *S. cerevisiae* (right)^{2, 4}.



The gene targeting method used for deleting *gdhA* and *GDH1* in *A. nidulans* and *S. cerevisiae*, respectively³. For *A. nidulans* the deletion was combined with the insertion of *gdhB* whereas *GDH2* was overexpressed by a promoter exchange in *S. cerevisiae*.

Physiological Characterization

Removal of Gdh1 (or GdhA) decreased the growth rate of both *S. cerevisiae* and *A. nidulans*. However the yield of 6-MSA was improved. In *A. nidulans* the yield was increased by 63%.



The effects of the deletion of Gdh1/A on growth rate of both *S. cerevisiae* and *A. nidulans*.

The improved yield of 6-MSA due to the deletion of Gdh1/A.

Conclusion

The predictions obtained by modelling were validated in both *S. cerevisiae* and *A. nidulans*. The yield of 6-MSA was improved for both strains. Thus it has been shown that this method has great potential for the future design of efficient cell factories.

References

- [1] Patil, K. R., Rocha, I., Förster, J., Nielsen, J., **Evolutionary programming as a platform for *in silico* metabolic engineering**, BMC Bioinformatics, 2005, 6:308.
- [2] Hansen, B. G., Salomonsen, B., Nielsen, M. T., Nielsen, J. B., Hansen, N. B., Nielsen, K. F., Regueira, T. B., Nielsen, J., Patil, K. R., Mortensen, U. H., **Versatile Enzyme Expression and Characterization System for *Aspergillus nidulans*, with the *Penicillium brevicompactum* Polyketide Synthase Gene from the Mycophenolic Acid Gene Cluster as a Test Case**, Applied and Environmental Microbiology, 2011, 3044–3051.
- [3] Nielsen, J.B., Nielsen, M.L., Mortensen, U.H., **Transiently disrupted non-homologous end joining pathway in *Aspergillus nidulans* allows simple and efficient gene targeting. A Technological Advancement**, Fungal Genetics Biology, 2008, 45: 165-170..
- [4] Wattanachaisaerekul, S., Lantz, A. E., Nielsen, M. L., Nielsen, J., **Production of the polyketide 6-MSA in yeast engineered for increased malonyl-CoA supply**, Metabolic Engineering, 2008, 246-254.