



## Vaccine Induced Specific Protection Against Enteric Red Mouth Disease (ERM) Caused by *Yersinia Ruckeri* Serotype 1 Biotype 2

Deshmukh, Sidharta; Raida, Martin K.; Dalsgaard, Inger; Chettri, Jiwan K.; Buchmann, Kurt

*Published in:*  
Book of abstracts

*Publication date:*  
2011

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Deshmukh, S., Raida, M. K., Dalsgaard, I., Chettri, J. K., & Buchmann, K. (2011). Vaccine Induced Specific Protection Against Enteric Red Mouth Disease (ERM) Caused by *Yersinia Ruckeri* Serotype 1 Biotype 2. In P. W. Kania, & K. Buchmann (Eds.), *Book of abstracts* DAFINET. <http://www.dafinet.dk/DAFINET/Home.html>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## VACCINE INDUCED SPECIFIC PROTECTION AGAINST ENTERIC RED MOUTH DISEASE (ERM) CAUSED BY *YERSINIA RUCKERI* SEROTYPE 1 BIOTYPE 2

Sidharta Deshmukh<sup>1</sup><sup>a</sup>, Martin K Raida<sup>1</sup>, Inger Dalsgaard<sup>2</sup>,  
Jiwan K Chettri<sup>1</sup> and K. Buchmann<sup>1</sup>

<sup>1</sup>Laboratory of Aquatic Pathobiology, Department of Veterinary Disease Biology, University of Copenhagen, Denmark. <sup>2</sup>National Veterinary Institute, Technical University of Denmark.

In European fish farms there is evidence of enteric red mouth disease (ERM) outbreaks in previously vaccinated farmed fish. It has been suggested that the occurrence of a *Yersinia ruckeri* variant (biotype 2) may explain this situation. Recent development of commercial vaccines has included both biotype 1 and 2. In this study, the specificity of immune protection extended by three commercial vaccines viz; AQUAVAC ERM<sup>®</sup> Intervet Schering Plough (based on biotype 1 only), ERMOGEN VET<sup>®</sup> Novartis (based on biotype 1 only) and AQUAVAC RELERA<sup>®</sup> Intervet Schering Plough (based on both biotype 1 and 2) developed against ERM was investigated following intraperitoneal (IP) challenge with *Yersinia ruckeri* serotype 1 biotype 2. Fish were immersion vaccinated for 30 s and challenged 2, 4 and 6 months post vaccination. The onset and severity of various pathological lesions along with their disappearance during the course of disease was also carried out to evaluate the protective index conferred by three different vaccines. After IP challenge, the overall best relative percentage survival was observed in AQUAVAC RELERA<sup>®</sup> followed by ERMOGEN VET<sup>®</sup> with least survival rates in AQUAVAC ERM<sup>®</sup> among all vaccinated groups. Interestingly a marginal better immune protection was observed between AQUAVAC RELERA<sup>®</sup> and ERMOGEN VET<sup>®</sup> vaccinated group during the last two challenge trial. The onset and severity of pathological lesions observed during challenge 2 (i.e. 4 month post vaccination) suggested a beneficial efficacy shown by AQUAVAC RELERA<sup>®</sup> in terms of milder and lesser degree of certain pathological lesions like haemorrhages in or around the buccal cavity, base of fins and intestines, when compared to ERMOGEN VET<sup>®</sup>, AQUAVAC ERM<sup>®</sup> vaccinated group and *Yersinia ruckeri* (BT2) infected group.