



Finite groups in which some particular subgroups are TI-subgroups

Shi, Jiangtao; Zhang, Cui

Published in:
Miskolc Mathematical Notes

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Shi, J., & Zhang, C. (2013). Finite groups in which some particular subgroups are TI-subgroups. *Miskolc Mathematical Notes*, 14(3), 1037-1040. http://mat76.mat.uni-miskolc.hu/~mnotes/index.php?page=article&name=mmn_785

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



FINITE GROUPS IN WHICH SOME PARTICULAR SUBGROUPS ARE TI-SUBGROUPS

JIANGTAO SHI AND CUI ZHANG

Received 6 June, 2013

Abstract. We prove that G is a group in which all noncyclic subgroups are TI-subgroups if and only if all noncyclic subgroups of G are normal in G . Moreover, we classify groups in which all subgroups of even order are TI-subgroups.

2010 Mathematics Subject Classification: 20D10

Keywords: noncyclic subgroup, subgroup of even order, TI-subgroup, normal

1. INTRODUCTION

All groups in this paper are considered to be finite. Let G be a group and M a subgroup of G . It is known that M is called a TI-subgroup of G if $M \cap M^g = 1$ or M for any $g \in G$.

In [5], Walls studied groups in which all subgroups are TI-subgroups. In [1], Guo, Li and Flavell classified groups in which all abelian subgroups are TI-subgroups. As a generalization of [1], we showed in [3] that G is a group in which all nonabelian subgroups are TI-subgroups if and only if all nonabelian subgroups of G are normal in G , and we showed in [4] that G is a group in which all nonnilpotent subgroups are TI-subgroups if and only if all nonnilpotent subgroups of G are normal in G .

In this paper, as a further generalization of above references, we first classify groups in which all noncyclic subgroups are TI-subgroups. We call such groups NCTI-groups. For NCTI-groups, we have the following result, the proof of which is given in Section 2.

Theorem 1. *A group G is an NCTI-group if and only if all noncyclic subgroups of G are normal in G .*

Remark 1. It is easy to show that any NCTI-group is solvable but it might not be supersolvable. For example, the alternating group A_4 is an NCTI-group but A_4 is nonsupersolvable.

The first author was supported by NSFC, Grant No. 11201401 and 11361075.

We say that a group G has nontrivial even order if $|G|$ is even but $|G| \neq 2$. In the following, we will classify groups in which all subgroups of even order are TI-subgroups. We call such groups EOTI-groups. For EOTI-groups, we have the following result, the proof of which is given in Section 3.

Theorem 2. *A group G is a EOTI-group if and only if one of the following statements holds:*

- (1) *all subgroups of G of nontrivial even order are normal in G .*
- (2) *$G = Z_p \rtimes \langle g \rangle$ is a Frobenius group with kernel Z_p and complement $\langle g \rangle$, where p is an odd prime and $o(g) = 2n$ for some $n > 1$.*

Remark 2. We can easily get that any EOTI-group is solvable. But the alternating group A_4 shows that a EOTI-group might not be supersolvable.

Arguing as in proof of Theorem 2, we can obtain the following three results, which are generalizations of Theorem 1, [3] and [4] respectively. Here we omit their proofs.

Theorem 3. *A group G is a group in which all noncyclic subgroups of even order are TI-subgroups if and only if all noncyclic subgroups of G of even order are normal in G .*

Theorem 4. *A group G is a group in which all nonabelian subgroups of even order are TI-subgroups if and only if all nonabelian subgroups of G of even order are normal in G .*

Theorem 5. *A group G is a group in which all nonnilpotent subgroups of even order are TI-subgroups if and only if all nonnilpotent subgroups of G of even order are normal in G .*

2. PROOF OF THEOREM 1

Proof. The sufficiency part is evident, we only need to prove the necessity part.

Let G be an NCTI-group. Assume that G has at least one nonnormal noncyclic subgroup. Choose M as a nonnormal noncyclic subgroup of G of largest order. Then, for any noncyclic subgroup K of G , we have that K is normal in G if $M < K$.

We claim that $M = N_G(M)$. Otherwise, assume that $M < N_G(M)$. Let L be a subgroup with $M < L \leq N_G(M)$ such that M has prime index in L . By the choice of M , we have that L is normal in G . Since M is not normal in G , $N_G(M) < G$. Take $h \in G \setminus N_G(M)$. We have $M^h < L^h = L$. Since M is a normal maximal subgroup of L and $M^h \neq M$, we have $L = MM^h$. Note that M is a TI-subgroup of G , which implies that $M \cap M^h = 1$. Then $|L| = |MM^h| = |M||M^h|$. It follows that $|M| = |M^h| = \frac{|L|}{|M|} = |L : M|$ is a prime, this contradicts that M is noncyclic. Thus $M = N_G(M)$.

Then, by the hypothesis, $M \cap M^a = 1$ for every $a \in G \setminus N_G(M) = G \setminus M$. We have that G is a Frobenius group with M being a Frobenius complement. Since G

is an NCTI-group, it follows that all nonabelian subgroups of G are TI-groups. By [3], we have that all nonabelian subgroups of G are normal in G . Then M is abelian. It follows that every Sylow subgroup of M must be cyclic by [2, Theorem 10.5.6]. This implies that M is cyclic, a contradiction. Thus G has no nonnormal noncyclic subgroup.

□

3. PROOF OF THEOREM 2

Proof. (1) We first prove the necessity part. Let G be a EOTI-group. Assume that G has at least one nonnormal subgroup of nontrivial even order. Let Q be any nonnormal subgroup of G of nontrivial even order. Choose L as a nonnormal subgroup of G of nontrivial even order of largest order such that $Q \leq L$.

We claim that $L = N_G(L)$. Assume that $L < N_G(L)$. Let K be a subgroup with $L < K \leq N_G(L)$ such that L has prime index in K . By the choice of L , we have that K is normal in G . Let y be an element of $G \setminus N_G(L)$. Then $L \cap L^y = 1$ by the hypothesis. It follows that $K = LL^y$ since L is a normal maximal subgroup of K . Then $|L| = |L^y| = |K : L|$ is a prime, a contradiction. Thus $L = N_G(L)$.

It follows that G is a Frobenius group with complement L . Let N be the Frobenius kernel of G . We have $G = N \rtimes L$.

We claim that L is a maximal subgroup of G . If L is not a maximal subgroup of G . Let M be a maximal subgroup of G such that $L < M$. It is obvious that M also has nontrivial even order. By the choice of L , we have that M is normal in G . Since G is a Frobenius group, it follows that either $M \leq N$ or $N < M$. If $M \leq N$, then $L < N$, a contradiction. If $N < M$, then $G = N \rtimes L \leq M$, again a contradiction. Thus L is a maximal subgroup of G .

It follows that N is a minimal normal subgroup of G . Since N is nilpotent by [2, Theorem 10.5.6], we can assume that $N = Z_p^m$, where $m \geq 1$ is a positive integer. Let d be an element of L of order 2. Since G is a Frobenius group and N is abelian, we have $e^d = e^{-1}$ for every $1 \neq e \in N$. Thus $\langle e \rangle \rtimes \langle d \rangle$ is a subgroup of G .

We claim that $\langle e \rangle \rtimes \langle d \rangle$ is normal in G . Note that $|\langle e \rangle \rtimes \langle d \rangle| = 2p$, where p is an odd prime. If $\langle e \rangle \rtimes \langle d \rangle$ is not normal in G . Arguing as the subgroup Q , there exists a nonnormal subgroup T of G of nontrivial even order of largest order such that $\langle e \rangle \rtimes \langle d \rangle \leq T$. And we have that T is also a Frobenius complement of G . But $N \cap T \geq \langle e \rangle \neq 1$, a contradiction. Thus $\langle e \rangle \rtimes \langle d \rangle$ is normal in G .

Since $\langle e \rangle \rtimes \langle d \rangle \not\leq N = Z_p^m$, we have $Z_p^m < \langle e \rangle \rtimes \langle d \rangle$. It follows that $m = 1$. Thus $N = Z_p$ is cyclic. By the N/C-theorem, since $C_G(N) = N$ it follows that L is isomorphic to a subgroup of $\text{Aut}(Z_p) \cong Z_{p-1}$. Then L is a cyclic group of order $2n$ for some $n > 1$. Assume that $L = \langle g \rangle$. We have $G = Z_p \rtimes \langle g \rangle$, where p is an odd prime and $o(g) = 2n$ for some $n > 1$.

(2) Now we prove the sufficiency part.

(i) If all subgroups of G of nontrivial even order are normal in G , then G is obviously a EOTI-group.

(ii) Assume that $G = Z_p \rtimes \langle g \rangle$ is a Frobenius group with kernel Z_p and complement $\langle g \rangle$, where p is an odd prime and $o(g) = 2n$ for some $n > 1$. Let S be any subgroup of G of even order. It is obvious that $S \cap Z_p = 1$ or Z_p .

If $S \cap Z_p = 1$, we have $S \leq \langle g \rangle^f$ for some $f \in G$. Obviously S is not normal in G . Then $N_G(S) = \langle g \rangle^f$. We have $S \cap S^x \leq \langle g \rangle^f \cap (\langle g \rangle^f)^x = 1$ for every $x \in G \setminus \langle g \rangle^f = G \setminus N_G(S)$.

If $S \cap Z_p = Z_p$, then $Z_p < S$. We have $S = S \cap G = S \cap (Z_p \rtimes \langle g \rangle) = Z_p \rtimes (S \cap \langle g \rangle)$. Then S is normal in G .

It follows that S is always a TI-subgroup of G whenever $S \cap Z_p = 1$ or Z_p . Then G is also a EOTI-group. □

ACKNOWLEDGEMENT

The authors are grateful to the referee who gives valuable comments and suggestions.

REFERENCES

- [1] X. Guo, S. Li, and P. Flavell, "Finite groups whose Abelian subgroups are TI-subgroups," *J. Algebra*, vol. 307, no. 2, pp. 565–569, 2007.
- [2] D. J. Robinson, *A course in the theory of groups*, 2nd ed., ser. Graduate Texts in Mathematics. New York: Springer-Verlag, 1995, vol. 80.
- [3] J. Shi and C. Zhang, "Finite groups in which all nonabelian subgroups are TI-subgroups," *J. Algebra Appl.*, vol. 13, 1350074, 2014.
- [4] J. Shi and C. Zhang, "A note on TI-subgroups of a finite group," *Algebra Colloq.*, to appear.
- [5] G. Walls, "Trivial intersection groups," *Arch. Math.*, vol. 32, pp. 1–4, 1979.

Authors' addresses

Jiangtao Shi

Yantai University, School of Mathematics and Information Science, 264005 Yantai, China

E-mail address: shijt@math.pku.edu.cn

Cui Zhang

Technical University of Denmark, Department of Applied Mathematics and Computer Science, DK-2800 Lyngby, Denmark

E-mail address: cuizhang2008@gmail.com