



Resistive sensitivity functions for van der Pauw astroid and rounded crosses and cloverleaves

Koon, Daniel; Hansen, Ole

Published in:
American Physical Society. Bulletin

Publication date:
2014

[Link back to DTU Orbit](#)

Citation (APA):
Koon, D., & Hansen, O. (2014). Resistive sensitivity functions for van der Pauw astroid and rounded crosses and cloverleaves. *American Physical Society. Bulletin*, 59(1).

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract Submitted
for the MAR14 Meeting of
The American Physical Society

Resistive sensitivity functions for van der Pauw astroid and rounded crosses and cloverleaves¹ DANIEL KOON, St. Lawrence University, OLE HANSEN, Technical University of Denmark — We have calculated the sensitivity of van der Pauw resistances to local resistive variations for circular, square and astroid discs of infinitesimal thickness, as well as for the families of rounded crosses and cloverleaves, as a function of specimen parameters, using the direct formulas of our recent paper (Koon *et al.* 2013 *J. Appl. Phys.* **114** 163710) applied to “reciprocally dual geometries” (swapped Dirichlet and Neumann boundary conditions) described by Mareš *et al.* (2012 *Meas. Sci. Technol.* **23** 045004). These results show that (a) the product of any such sensitivity function times differential area, and thus (b) the ratio of any two sensitivities, is invariant under conformal mapping, allowing for the pointwise determination of the conformal mapping function. The family of rounded crosses, which is bounded in parameter space by the square, the astroid and an “infinitesimally thin” cross, seems to represent the best geometry for focusing transport measurements on the center of the specimen while minimizing errors due to edge- or contact-effects.

¹Made possible by an SLU Faculty research grant.

Daniel Koon
St. Lawrence University

Date submitted: 15 Nov 2013

Electronic form version 1.4