



The effect of whole-grain compared to refined wheat on the gut microbial composition and integrity in a colonic epithelial cell model following a 12-week energy-restricted dietary intervention in postmenopausal women

Christensen, Ellen Gerd; Licht, Tine Rask; Kristensen, M.; Bahl, Martin Iain

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Christensen, E. G., Licht, T. R., Kristensen, M., & Bahl, M. I. (2012). *The effect of whole-grain compared to refined wheat on the gut microbial composition and integrity in a colonic epithelial cell model following a 12-week energy-restricted dietary intervention in postmenopausal women*. Abstract from 7th Danish Conference on Biotechnology and Molecular Biology, Vejle, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The effect of whole-grain compared to refined wheat on the gut microbial composition and integrity in a colonic epithelial cell model following a 12-week energy-restricted dietary intervention in postmenopausal women

E. G. Christensen¹, T.R. Licht¹, M. Kristensen², M.I. Bahl^{1*}

(1) Gut Ecology, Division of Food Microbiology, DTU Food, Technical University of Denmark, Søborg, Denmark

(2) Department of Human Nutrition, University of Copenhagen, Denmark

*mbah@food.dtu.dk

Intake of whole-grain products are considered to decrease the risk of cardiovascular disease (CVD). This effect could potentially be linked to a prebiotic effect, hence positive modulation of the gut microbial composition or activity. Kristensen and coworkers recently conducted a study in postmenopausal women who were randomized to either whole-grain wheat (WW) (n=38) or refined wheat (RW) (n=34) consumption as part of an energy-restricted diet for 12-weeks following a 2-week run-in period with RW. Percentage fat mass as well as serum total and LDL cholesterol were found to differ between the two groups (Kristensen, et al, 2012). We used fecal samples from the same study to examine effects of WW and RW on the bacterial composition by quantitative PCR targeting the phylums *Bacteroidetes* and *Firmicutes*, and the genera *Bifidobacteria*, *Lactobacillus*, *Bacteroides*, and *Prevotella*, as well as the *Enterobacteriaceae* family. Potential bifidogenic effects were examined in depth by determining the levels of *B. bifidum*, *B. adolescentis*, *B. catenulatum*, and *B. longum*. Potential correlations between changes in microbiota and cardiovascular risk factors are to be investigated.

The composition of the gut microbiota may affect the intestinal integrity, which in this study was evaluated *in vitro* by determining transepithelial resistance (TER) across a Caco-2 cell monolayer. Fecal water collected after the run-in period and following the intervention period for 26 participants (WW; 15 participants, RW; 11 participants) were used to determine effects of WW, RW, and microbiota composition on TER. Preliminary results indicate that fecal water from WW and RW both before and after intervention in general had a positive effect on TER, however, there was no difference in TER between WW and RW. Potential correlations between microbial composition and effect on intestinal integrity are to be examined further.

References:

Kristensen, M., Toubro, S., Jensen, M.G., Ross, A.B., Riboldi, G., Petronio, M., Bügel, S., Tetens, I., Astrup, A. (2012) Wholegrain compared with refined wheat decreases the percentage of body fat following a 12-week, energy-restricted dietary intervention in postmenopausal women. *The Journal of Nutrition*, doi: 10.3945/jn.111.142315