



## Strong light-matter interaction in graphene - Invited talk

**Xiao, Sanshui**

*Publication date:*  
2014

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Xiao, S. (2014). *Strong light-matter interaction in graphene - Invited talk*. Abstract from 35th Progress In Electromagnetics Research Symposium , Guangzhou (Canton), China.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

---

# Strong Light-matter Interaction in Graphene

Sanshui Xiao<sup>1,2</sup>

<sup>1</sup>Department of Photonics Engineering, Technical University of Denmark, DK-2800 Kongens Lyngby, Denmark

<sup>2</sup>Centre for Nanostructured Graphene (CNG), Technical University of Denmark, DK-2800 Kongens Lyngby, Denmark

**Abstract**— Graphene has attracted lots of attention due to its remarkable electronic and optical properties, thus providing great promise in photonics and optoelectronics. However, the performance of these devices is generally limited by the weak light-matter interaction in graphene. The combination of graphene with noble-metal nanostructures is currently being explored for strong light-graphene interaction. We introduce a novel hybrid graphene-metal system for studying light-matter interactions with gold-void nanostructures exhibiting resonances in the visible range[1]. The hybrid system is further explored for sensing of Rhodamine 6G molecules with respect to the strong surface-enhanced Raman scattering. The interaction between graphene plasmon (supported by nanodot and antidot arrays) and the substrate phonons [2] is also experimentally demonstrated and structural control is used to map out the hybridization of plasmons and phonons, where the graphene is structured by the nanosphere lithography with structural control down to the sub-100 nanometer regime.

## REFERENCES

1. Zhu, X., Shi, L., Schmidt, M. S., Boisen, A., Hansen, O., Zi, J., Xiao, S., and Mortensen, N. A., “Enhanced light-matter interaction in graphene-coved gold nanovoid array,” *Nano Lett.*, Vol. 13, No. 10, 4090–4096, 2013.
2. Zhu, X., Wang, W., Yan, W., Larsen, M. B., Boeggild, P., Pedersen, T. G., Xiao, S., Zi, J., and Mortensen, N. A., “Plasmon-phonon coupling in large-area graphene dot and antidot arrays fabricated by nanosphere lithography,” *Nano Lett.*, in press, 2014.